

Anna Brzozowska
Joanna Kaluza
Kim T.B. Knoops
Lisette C.P.G.M. de Groot

Supplement use and mortality: the SENECA study

Received: 22 August 2007
Accepted: 2 April 2008
Published online: 14 April 2008

Prof. A. Brzozowska (✉) · J. Kaluza, PhD
Dept. Human Nutrition
Warsaw University of Life Sciences-SGGW
159c Nowoursynowska St.
02-776 Warsaw, Poland
Tel.: +48-22/5937-110
Fax: +48-22/5937-117
E-Mail: anna_brzozowska@sggw.pl

K.T.B. Knoops, PhD ·
Prof. L.C.P.G.M. de Groot
Division of Human Nutrition
Wageningen University
Wageningen, The Netherlands

Abstract *Background* It is hypothesis that in relatively healthy older people supplement usage can be consider as healthy life style habit and as such can positively influence longevity. *Aim of the study* To determine whether supplement use was associated with all-cause mortality in the participants of the SENECA study. *Methods* Baseline measurements were carried out in 1988/1989 among 75 to 80-year-old people living in 15 European small towns. All-cause mortality was followed up to April 30, 1999. Data from 920 men and 980 women who were ischemic heart diseases-, stroke- and cancer-free at baseline were included. The multivariate adjusted (for sex, age, years of education, physical activity, BMI, chronic diseases, Mediterranean Diet Score, alcohol use and the place of living) hazard ratio (HRs) and 95% confidence intervals (CIs) of mortality by use of any type of nutrient supplement and by particular nutrient supplement use were estimated by Cox proportional hazards regression

models. Results At baseline, 13% of participants used nutritional supplements, 19% of subjects were smokers. During 10 years of follow-up 445 men and 252 women died. Among non-smokers no significant associations between total supplement use and particular nutrient supplement use were observed. Among smokers use of any type of supplements (Multivariate HR: 1.52; 95%CI: 1.02–2.28), use of vitamin B₁ (Multivariate HR: 1.57; 95%CI: 1.00–2.48) and vitamin B₂ supplements (Multivariate HR: 1.60; 95%CI: 1.00–2.56) were associated with a significantly higher risk of all-cause mortality. The similar tendencies were observed among vitamin B₆ and vitamin C supplement users who were smokers. *Conclusions* Among smokers, participants of the SENECA study, supplement use increased all-cause mortality risk.

Key words elderly – Europe – supplement use – mortality – smoking

Introduction

Despite the wide use of vitamin and mineral supplements in different populations, the effects of such a

practice on health have not been well defined [20]. Several studies [16, 17, 19] reported that supplement usage is beneficial for primary and secondary prevention of some diseases and improves quality of life. Holmquist et al. [17] study indicated that using of low

doses of multivitamin supplements was relevant to lowering risk of myocardial infarction. Jacobs et al. [18, 19] observed significant decreased the risk of stomach cancer mortality among vitamin C users and colon cancer mortality among multivitamin supplements users. In contrast other studies [15, 18, 20, 28, 34 and 41] have failed to demonstrate the positive effect of supplementation in preventing age-related diseases.

The baseline round of the Survey in Europe on Nutrition and the Elderly, a Concerted Action (SENECA) revealed, on the basis of the blood analyses, a significant prevalence of deficiency for several nutrients including vitamin B₆ (up to 50% of subjects), vitamin B₁₂ (2–10%) and vitamin E (1–21%) among older people from small European towns [13]. To counteract low nutritional status only 13% of respondents reported having taken nutritional supplements [1]. The aim of this paper was to determine whether the use of supplements was associated with health benefits, measured as lowering all-causes mortality in the participants of SENECA study.

Subjects and methods

■ Study population

Data presented in this paper were collected at baseline (1988/1989) and final (1999) round of the SENECA survey. The population under study consisted of randomly selected, born between 1913 and 1918, habitants of 15 small towns in 11 European countries: Hamme (B), Roskilde (DK), Haguenau and Romans (F), Anogia and Archanes (GR), Monor (H), Padua, Fara Sabina, Magliano Sabina and Poggio Mirteto (I), Culemborg (NL), Vila Franca de Xira (P), Betanzos (E), Yverdon-les-Bains, Burgdorf and Bellinzona (CH), Marki (PL) [11].

In view of a poor prognosis and the high risk of death, subjects who suffered from ischemic heart diseases (376 subjects), stroke (63 subjects) and malignancy (46 subjects) were excluded. Finally, the analyses included 920 men and 980 women.

■ Data collection

Data were collected by interview at the participants' homes according to a standardised protocol [10]. Information about use of pharmaceutical preparations containing vitamins or/and minerals was collected with the question: "Do you use medicines, vitamins/minerals (tablets or injections) or tonics? If yes, please show them." Participants were also asked about daily doses. Having data on the composition of

supplements available for subjects the intake of 9 vitamins, calcium and iron were calculated on the local level.

Modified dietary history method was used to collect information about food consumption during the month before the interview. This method was validated against a 3-day weight record method in a subsample of 82 elderly subjects from 11 participating centres. Pearson's correlation coefficient for nutrient intake varied from 0.18 (vitamin A) to 0.79 (water) with a median coefficient of 0.58 [29]. The quality of diet was assessed by modified Mediterranean Diet Score (MDS) which comprised: ratio of MUFA to SFA; intake of: legumes, nuts, and seeds; cereal products and grains; fruits; vegetables and potatoes; meat and meat products; dairy products; and fish. The diet score varied from 0 (low quality diet) to 8 (high quality diet) [22].

Voorrips questionnaire was used to assess habitual physical activities of the last year. Items on household activities were estimated from very active to inactive (4–5 possible ratings). Sports and other activities were determined as type of activity; hours per day or week spend on it, and the period of the year in which the activity was normally performed. The value of Voorrips Score was calculated on bases on the net energetic cost of particular activities [40].

Non-smoking subjects were defined as never smoking or non-smoking since more than 15 years ago at baseline round of the study. Data about mortality were collected up to April 30, 1999.

■ Statistical analysis

All statistical analyses were done using the SPSS software (v.11.0). Because there were statistically significant interactions between smoking status of participants and use of any type of nutrient supplements as well as use of particular nutrient supplements, all the analyses were carried out separately for smokers and non-smokers.

The unadjusted Kaplan–Meier curves by supplement use and smoking status were fitted, and log-rank test was used to compare survival distribution between supplement users and non-users. Cox proportional hazards regression models were used to estimate the hazard ratios (HRs) and 95% confidence intervals (CIs) of mortality for users of any type of nutrient supplements as well as for users of particular nutrient supplements (vitamin B₁, B₂, B₆, B₁₂, C, A, D, E, folic acid, calcium and iron, separately) versus non-users. Multivariate HRs estimates and 95% CIs were adjusted for sex (men, women), age (continuous variable), years of education (continuous variable), physical activity (Voorrips Score: ≤8 points, >8 points),

Table 1 Description of the population (baseline study)

	Men (N = 920)	Women (N = 980)
Age (years) means \pm SD	73.3 \pm 1.8	73.2 \pm 1.9
Education (years) means \pm SD	8.6 \pm 4.1	7.5 \pm 3.6
Self-perceived health (%)		
Very poor and poor	6.7	10.8
Fair	24.7	31.5
Good and very good	68.3	56.7
Chronic disease (%)	64.5	76.3
Mediterranean diet score, modified (%)		
≤ 4 points	60.8	59.9
> 4 points	39.2	40.1
Smoking (%)	32.0	7.5
Alcohol use (%)	79.6	49.3
Supplement use (%)	10.9	14.8
Voorrips score (points) means \pm SD	18.5 \pm 13.4	11.8 \pm 9.5
Body mass index (kg/m ²) means \pm SD	26.6 \pm 3.8	27.0 \pm 4.7
< 20 (%)	3.2	5.4
20–25 (%)	30.5	30.0
25–30 (%)	48.9	39.4
> 30 (%)	17.4	25.2
Albumin (g/l) means \pm SD	41.9 \pm 3.2	41.6 \pm 3.0

BMI (≤ 25 kg/m², > 25 kg/m²), chronic diseases (yes, no), modified MDS (≤ 4 points, > 4 points), alcohol use (non-consumption, consumption) and the place of living (latitude $\leq 47^\circ$ North, $> 47^\circ$ North). The statistical significance was considered at $P < 0.05$.

Results

■ Characteristics of the cohort

At the beginning of the study the mean age of the subjects was 73.3 ± 1.8 years (Table 1). During the

baseline study, 11% of men and 15% of women used supplements, 32% of men and 8% of women were smokers. Fourteen percent of smoking men and 5% of women smoked more than 25 cigarettes or other tobaccos per day, 26% of smoking men and 16% of women smoked 15–24 cigarettes, 26% men and 35% of women smoked 5–14 cigarettes and other men and women smoked ≤ 4 cigarettes. A high percentage of study population suffered from chronic diseases (65% of men and 76% of women) as well as overweight and obesity (49 and 17% of men and 39 and 25% of women, respectively). The diets of approximately 40% of participants were scored above four points of a modified MDS.

During 10 years of SENECA study 445 men and 252 women died.

■ Intake of vitamins and minerals from supplements in relation to smoking status

A higher percentage of smoking men and smoking women used supplements (14.4 and 32.9%, respectively) in comparison to non-smoking subjects (9.2 and 13.4%, respectively). Vitamins B₁, B₂ and B₆ and C were used most often among both, smokers and non-smokers, while folic acid and calcium supplements were used by a lower number of subjects (Table 2).


The median intake of vitamins and iron from supplements was in the range of recommendations for these nutrients while median doses of supplemental calcium were very low. A part of non-smokers subjects used large doses of iron, the 90th percentile of supplemental intake of this nutrient amounted 67 mg/day, what exceeded the upper intake level (25 g/day) proposed in Nordic Recommendations [30].

Table 2 Intake of vitamins and minerals from supplements in relation to smoking status

Nutrient in supplements	Supplement users							
	Non-smoking (N = 176)				Smoking (N = 66)			
	%N	Percentiles			%N	Percentiles		
		10	50	90		10	50	90
Vitamin B ₁ (mg/d) ^a	57	1.2	2.0	63	71	1.0	2.0	20
Vitamin B ₂ (mg/d) ^a	56	1.5	2.0	26	67	1.0	2.0	8
Vitamin B ₆ (mg/d) ^a	60	0.8	2.0	36	69	0.5	2.0	10
Folic acid (μ g/d) ^a	29	100	100	400	36	50	100	200
Vitamin B ₁₂ (μ g/d) ^a	38	1.7	3.0	249	61	1.0	3.0	600
Vitamin C (mg/d) ^a	51	16	60	209	68	20	60	200
Vitamin A (μ g/d) ^a	44	200	1,200	3,000	59	300	1,000	4,000
Vitamin D (μ g/d) ^a	43	5	10	26	64	8	10	355
Vitamin E (mg/d) ^a	44	3	10	103	58	1	10	71
Calcium (mg/d)	35	3	162	1,000	23	5	97	700
Iron (mg/d) ^a	33	4	16	67	53	2.3	18	22

^aSignificant difference in prevalence of supplementation practice between non-smokers and smokers

Fig. 1 The unadjusted Kaplan–Meier survival curves for subjects from SENECA's baseline study by supplement use and smoking status



■ Supplement use and mortality

Among non-smokers unadjusted hazard ratio was a statistically significantly lower (28%) among supplement users than among non-users (Figure 1). The similar tendencies were observed among users of vitamins B₁, B₂ and B₆ and calcium. In contrast, among smokers there were tendencies towards an increase in mortality among users of any type of supplements (37%) as well as among vitamin B₁, B₂, B₆ and vitamin C supplement users (43–47%) versus non-users.

After multivariate adjustments (Table 3), among non-smokers none of the differences in all-cause mortality related to total or particular nutrients supplement use were statistically significant. Whereas among smokers the multivariate-adjusted hazard ratio for subjects who used any type of nutrient supplements compare to those who did not use them was significantly higher (52%, $P = 0.04$), for vitamin B₁ and B₂ supplements users HRs were also higher (57 and 60%, respectively, $P = 0.05$). The similar tendencies were observed among smokers—vitamin B₆ and vitamin C supplement users.

Discussion

Our study shows that supplement use is associated with mortality differently in smoking and non-smoking elderly and provide no evidence of a beneficial effect of supplement use on the risk of mortality during 10 years follow-up among SENECA's population. Slight adverse effects were found in smoking subjects. In smoking subjects the supplement use (any type or vitamin B₁, and B₂) was associated with slight adverse effects.

Table 3 Hazard ratios (HRs) and 95% confidence intervals (CIs) for all-cause mortality in SENECA-population by supplement use and smoking status unadjusted and adjusted for possible confounders

Users of nutrient supplements	Non-smoking (N = 1,527)		Smoking (N = 365)	
	HR	95% CI	HR	95% CI
Unadjusted				
Total ^a	0.72	0.53–0.97	1.37	0.97–1.94
Vitamin B ₁	0.74	0.50–1.09	1.45	0.98–2.13
Vitamin B ₂	0.74	0.45–1.02	1.44	0.96–2.15
Vitamin B ₆	0.67	0.45–1.00	1.43	0.96–2.12
Folic acid	0.73	0.42–1.26	1.14	0.66–1.97
Vitamin B ₁₂	0.97	0.62–1.49	1.25	0.82–1.92
Vitamin C	0.76	0.51–1.15	1.47	0.99–2.18
Vitamin A	0.75	0.48–1.16	1.21	0.78–1.87
Vitamin D	0.83	0.54–1.28	1.17	0.76–1.80
Vitamin E	0.86	0.56–1.30	1.13	0.72–1.78
Calcium	0.62	0.37–1.06	0.85	0.40–1.80
Iron	0.68	0.40–1.15	1.27	0.81–2.00
Adjusted^b				
Total ^a	0.80	0.57–1.11	1.52	1.02–2.28
Vitamin B ₁	0.80	0.52–1.23	1.57	1.00–2.48
Vitamin B ₂	0.71	0.45–1.11	1.60	1.00–2.56
Vitamin B ₆	0.71	0.46–1.09	1.52	0.96–2.42
Folic acid	0.77	0.42–1.42	1.19	0.64–2.20
Vitamin B ₁₂	1.12	0.70–1.79	1.22	0.74–2.00
Vitamin C	0.71	0.45–1.13	1.55	0.97–2.46
Vitamin A	0.82	0.50–1.34	1.22	0.73–2.03
Vitamin D	0.91	0.57–1.46	1.27	0.77–2.11
Vitamin E	0.97	0.62–1.51	1.31	0.78–2.21
Calcium	0.72	0.41–1.26	0.81	0.33–2.00
Iron	0.85	0.49–1.49	1.34	0.79–2.27

^aUsers of any type of nutrient supplements

^bAdjusted for sex, age, education, physical activity, body mass index, chronic diseases, Mediterranean Diet Score, alcohol use, latitude

Some meta-analysis of randomized controlled trials also showed no evidence of a protective effect of supplement use on mortality or morbidity [3, 5, 6, 8, 9]. No evidences of a protective effect were found in

the case of antioxidant or B-group vitamin supplements on the progression of atherosclerosis [6] and antioxidant supplements on gastrointestinal cancer [5], of vitamin C and vitamin E supplements use on various type of cancer [9], as well as usage of vitamin A, C, E and folate supplements on prevention of lung cancer [8]. Moreover, Bjelakovic et al. [4] who analyzed 68 randomized trials with 232,606 participants showed that treatment with antioxidant vitamins may increase all-cause mortality.

However, observational and experimental studies reported also inverse association between supplement use and mortality. Slightly decreased risk of colon cancer mortality among multivitamin users was observed with stronger association among moderate to heavy alcohol users. At least 15 years of follow-up was required to see the benefit of supplement use [19]. Low doses of multivitamin supplements were associated with substantially lower risk of nonfatal myocardial infarction in Stockholm Heart Epidemiology Program [17]. The results of SU.VI.MAX study indicated that nutritional doses of a combination of vitamin C, vitamin E, beta-carotene, selenium and zinc might have a protective effect on mortality in apparently healthy men in the age of 45–60 years [16].

Similarly as in our study, the adverse effect of supplement use on survival among smokers was reported by others [24, 32, 38, 41]. The β -Carotene and Retinol Efficacy Trial (CARET) was stopped 21 months earlier than planned because after 4 years the relative risks of death from any cause, from lung cancer and from cardiovascular diseases were higher among subjects who received combination of β -carotene and vitamin A than among the placebo group [32].

Watkins et al. [41] showed that the mortality from all cancers was significantly higher among smoking men who used multivitamins alone (HR: 1.13, 95%CI: 1.05–1.23) or in combination with vitamin A, C or E (HR: 1.16, 95%CI: 1.06–1.23); whereas among subjects who had never or had formerly smoked and used supplements hazard ratios were significantly lower, i.e. 0.86 (95%CI: 0.74–0.99) and 0.90 (95%CI: 0.82–0.98), respectively.

Moreover, among the participants of the Finnish ATBC Cancer Prevention Study α -tocopherol supplementation increased the risk for adenomas (HR: 1.66, 95%CI: 1.19–2.32), whereas β -carotene supplementation had no effect on the risk (HR: 0.98, 95%CI: 0.71–1.35) [25].

The mechanisms of the adverse effect of supplemental vitamins are still not fully explained. Probably, they could be related to pro-oxidative role of antioxidants when their doses are too high or as the results of interactions with components of cigarette

smoke. In our study the median intake of supplemental nutrients did not exceed significantly the recommended dietary intake among smokers, but the 90th percentile of intake exceeded three to ten times the RDA's for vitamin B₁, B₂, and C. High dose of those supplements probably had an adverse effect on health of participants who were smokers.

In humans different cellular markers of DNA damage were used to assess the pro-oxidative effects of vitamin C supplements in smokers. Welch et al. [42] observed a significant increase of 7-hydroxy-8-oxo-2'-deoxyguanosine concentrations after β -carotene supplementation in smokers, whereas in non-smokers the concentration of this marker decreased. Rehman et al. [36], who examined the effects of iron (14 mg/day) and vitamin C (either 60 or 260 mg/day) supplementation found that in subjects with higher initial level of plasma vitamin C total oxidative DNA damage in white blood cells significantly increased after supplementation. Whereas, in low plasma ascorbate subjects the supplementation was beneficial.

In clinical trial among smokers who regularly received 500 mg/day vitamin C a significant increase of plasma TBARS was shown, what indicated the increase of oxidative lipid damage [31].

In contrast, others studies were failed in founding the significant changes in DNA damage or LDL-lipid peroxidation after supplementation with vitamin C [2, 27], vitamin E [37], β -carotene [12, 35, 39] and vitamin combinations [33]. Moreover, also protective effect of supplement use on DNA damage was observed [7, 14, 23, 26].

Moreover, contrary to expectations high supplemental folate levels and folate intervention, after microscopic neoplastic foci are established in the colorectal mucosa, promote rather than suppress colorectal carcinogenesis. The results of animal studies suggest that folate possesses the dual modulators effect on carcinogenesis depending on the timing and dose of folate intervention. Folate deficiency has an inhibitory effect whereas folate supplementation has a promoting effect on progression of established neoplasms [21].

No data on the effect of B-group vitamins supplementation on mortality among smokers was found in the literature.

Data on vitamin/mineral supplement use presented in this paper were collected among free-living elderly, aged at the beginning of the study 70–75 years. At that time the prevalence of supplement use was rather small, i.e. 13%. Even the subjects were asked about current supplement use many of them were long-term users. Of the 1,043 participants who were reinterviewed in 1993 12% reported using supplements at baseline and after 5 years, 69% of participants

reported not using supplements at either time. A potential limitation of this study was the change in classification of 19% of the respondents after 5 years from the beginning of the study, i.e. 5% gave up this practice and 14% became supplement users. Our reference group therefore included some participants who become users after 5 years of observation. Another limitation was related to a statistical significant interaction between supplements use and smoking status of the participants. Dividing subjects into smokers and non-smokers groups made the subgroups small in number of cases and decreased the power of analysis.

In summary, supplementation practice had no beneficial effect on mortality among 70 to 75-year-old European participants of SENECA population. Our results indicated that some supplements should be used with special cautiously, and further studies are necessary, which also would allow for explanation of the mechanisms of harmful effects of supplements among smoking subjects.

■ **Acknowledgments** This study constitutes a part of the Healthy Ageing: a Longitudinal study in Europe (HALE) Project (QLK6-CT-2000-00211) and is funded by the European Union under Specific Programme "Quality of Life and Management of Living Resources".

References

1. Amorim Cruz JA, Moreiras O, Brzozowska A (1996) Longitudinal changes in the intake of vitamins and minerals of elderly Europeans. *Eur J Clin Nutr* 50(suppl 2):S77-S85
2. Anderson D, Philips BJ, Yu TW, Edwards AJ, Ayesh R, Butterworth KR (1997) The effects of vitamin C supplementation on biomarkers of oxygen radical generated damage in human volunteers with "low" or "high" cholesterol levels. *Environ Mol Mutagen* 30:161-174
3. Bjelakovic G, Nagorni A, Nikolova D, Simonetti RG, Bjelakovic M, Gluud C (2006) Meta-analysis: antioxidant supplements for primary and secondary prevention of colorectal cancer. *Aliment Pharmacol Ther* 24:281-291
4. Bjelakovic G, Nikolova D, Gluud L, Simonetti RG, Gluud C (2007) Mortality in randomized trial of antioxidant supplements for primary and secondary prevention. *JAMA* 297:842-857
5. Bjelakovic G, Nikolova D, Simonetti RG, Gluud C (2004) Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet* 364:1219-1228
6. Bleys J, Miller ER, Pastor-Barriuso R, Appel LJ, Guallar E (2006) Vitamin-mineral supplementation and the progression of atherosclerosis: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 84:880-887
7. Brown KM, Morrice PC, Duthie GG (1994) Vitamin E supplementation suppresses indexes of lipid peroxidation and platelet counts in blood of smokers and nonsmokers but plasma lipoprotein concentrations remain unchanged. *Am J Clin Nutr* 60:383-387
8. Cho E, Hunter DJ, Spiegelman D, Albanes D, Albanes D, Beeson WL, van den Brandt PA, Colditz GA, Festkanich D, Folsom AR, Fraser GE, Freudenheim JL, Giovannucci E, Goldbohm RA, Graham S, Miller AB, Rohan TE, Sellers TA, Virtamo J, Willett WC, Smith-Warner SA (2006) Intakes of vitamins A, C and E and folate and multivitamins and lung cancer: a pooled analysis of 8 prospective studies. *Int J Cancer* 118:970-978
9. Coulter ID, Hardy ML, Morton SC, Hilton LG, Tu W, Valentine D, Shekelle PG (2006) Antioxidants vitamin C and vitamin E for the prevention and treatment of cancer. *J Gen Intern Med* 21:735-744
10. de Groot L, van Staveren W (1988) Nutrition and the elderly. Manual of Operations, Euronut Report 11, Wageningen, The Netherlands
11. de Groot L, van Staveren W (1991) Description of survey towns and population. *Eur J Clin Nutr* 45(suppl 3):23-29
12. Gaziano JM, Hattar A, Flynn M, Johnson EJ, Krinsky NI, Ridker PM, Hennekens CH, Frei B (1995) Supplementation with β -carotene in vivo and in vitro does not inhibit low density lipoprotein oxidation. *Atherosclerosis* 112:187-195
13. Haller J, Lowik M, Ferry M, Ferro-Luzzi A (1991) Nutritional status: blood vitamins A, E, B₆, B₁₂, folic acid and carotene. *Eur J Clin Nutr* 45(suppl 3):63-82
14. Harats D, Ben-Naim M, Dabach Y, Hollander G, Havivi E, Stein O, Stein Y (1990) Effects of vitamin C and E supplementation on susceptibility of plasma lipoproteins to prevention induced by acute smoking. *Atherosclerosis* 85:47-54
15. Hennekens CH, Buring JE, Manson JE, Stampfer M, Rosner B, Cook NR, Belanger C, LaMotte F, Gaziano JM, Ridker PM, Willett W, Peto R (1996) Lack of effect of long-term supplementation with beta carotene on incidence of malignant neoplasms and cardiovascular disease. *N Engl J Med* 334:1145-1149
16. Hercberg S, Galan P, Preziosi P, Bertais S, Mennen LI, Malvy D, Roussel AM, Favier A, Briancon S (2003) Effects of long-term supplementation with nutritional doses of antioxidant vitamins and minerals on mortality and cancer: results of the SU.VI.MAX Study. *Ann Nutr Metab* 47:416
17. Holmquist C, Larsson S, Wolk A, de Faire U (2003) Multivitamin supplements are inversely associated with risk of myocardial infarction in men and women—Stockholm Heart Epidemiology Program (SHEEP). *J Nutr* 133:2650-2654
18. Jacobs EJ, Connell CJ, McCullough ML, Chao A, Jonas CR, Rodriguez C, Calle EE, Thun MJ (2002) Vitamin C, vitamin E, and multivitamin supplement use and stomach cancer mortality in the Cancer Prevention Study II Cohort. *Cancer Epidemiol Biomarkers Prev* 11:35-41
19. Jacobs EJ, Connell CJ, Patel AV, Chao A, Rodriguez C, Seymour J, McCullough ML, Calle EE, Thun MJ (2001) Multivitamin use and colon cancer mortality in Cancer Prevention study II cohort (United States). *Cancer Causes Control* 12:927-934
20. Kim I, Williamson DF, Byers T, Koplan JP (1993) Vitamin and mineral supplement use and mortality in a US cohort. *Am J Public Health* 83:546-550
21. Kim YI (2003) Role of folate in colon cancer development and progression. *J Nutr* 133(11 suppl 1):3731S-3739S

22. Knoops KTB, de Groot LCPGM, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, van Staveren W (2004) Mediterranean diet, lifestyle factors, and 10-years mortality in elderly European men and women: the HALE project. *JAMA* 292:1433–1439
23. Lee BM, Lee SK, Kim HS (1998) Inhibition of oxidative DNA damage, 8-OHdG, and carbonyl contents in smokers treated with antioxidants (vitamin E, vitamin C, β -carotene and red ginseng). *Cancer Lett* 132:219–227
24. Leppala JM, Virtamo J, Fogelholm R, Huttunen JK, Albanes D, Taylor PR, Heinonen OP (2000) Controlled trial of α -tocopherol and β -carotene supplements on stroke incidence and mortality in male smokers. *Arterioscler Thromb Vasc Biol* 20:230–235
25. Malila N, Virtamo J, Virtanen M, Albanes D, Tangrea JA, Huttunen JK (1999) The effect of α -tocopherol and β -carotene supplementation on colorectal adenomas in middle-age male smokers. *Cancer Epidemiol Biomarkers Prev* 8:489–493
26. Moller P, Viscovich M, Lykkesfeldt J, Loft S, Jesen A, Poulsen H (2004) Vitamin C supplementation decreases oxidative DNA damage in mononuclear blood cells of smokers. *Eur J Nutr* 43:267–274
27. Mulholland CW, Strain JJ, Trinick TR (1996) Serum antioxidant potential, and lipoprotein oxidation in female smokers following vitamin C supplementation. *Int J Food Sci Nutr* 47:227–231
28. Muntwyler J, Hennekens CH, Manson JE, Buring JE, Gaziano JM (2002) Vitamin supplement use in low-risk population of US male physicians and subsequent cardiovascular mortality. *Arch Intern Med* 162:1472–1476
29. Nes M, van Staveren WA, Zajkas G, Inelman EM, Morieras-Varela O (1991) Validity of the dietary history method in elderly subjects. *Eur J Clin Nutr* 45(suppl 3):97–104
30. Nordic Nutrition Recommendations (2004) Integrating nutrition, physical activity. Nordic Council of Ministers, Copenhagen
31. Nyyssönen K, Poulsen HE, Hayn M, Agerbo P, Porkkala-Sarataho E, Kaikkonen J, Salonen R, Salonen JT (1997) Effect of supplementation of smoking men with plain or slow release ascorbic acid on lipoprotein oxidation. *Eur J Clin Nutr* 51:154–163
32. Omenn GS, Goodman GE, Thornquist MD, Balmes J, Gullen MR, Glass A, Keogh JP, Meyskens FL, Valanis B, Williams JH, Barnhart S, Hammar S (1996) Effects of combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *N Engl J Med* 334:1150–1155
33. Preime H, Loft S, Nyyssönen K, Salonen JT, Poulsen HE (1997) No effect of supplementation with vitamin E, ascorbic acid, or coenzyme Q on oxidative damage estimated by 8-oxo-7,8-dihydro-2'-deoxyguanosine excretion in smokers. *Am J Clin Nutr* 65:503–507
34. Rautalahti MT, Virtamo J, Taylor PR, Heinonen OP, Albanes D, Haukka JK, Edwards BK, Karkkainen PA, Stolzenberg-Solomon RZ, Huttunen J (1999) The effects of supplementation with α -tocopherol and β -carotene on incidence and mortality of carcinoma of the pancreas in a randomized, controlled trial. *Cancer* 86:37–42
35. Reaven PD, Ferguson E, Navab M, Powell FL (1994) Susceptibility of human LDL to oxidative modification. Effects of variations in β -carotene concentration and oxygen tension. *Arterioscler Thromb* 14:1162–1169
36. Rehman A, Collis CS, Yang M, Kelly M, Diplock AT, Halliwell B, Rice-Evans C (1998) The effects of iron and vitamin C co-supplementation on oxidative damage to DNA in healthy volunteers. *Biochem Biophys Res Commun* 246:293–298
37. Reilly M, Delanty N, Lawson JA, Fitzgerald GA (1996) Modulation of oxidant stress in vivo in chronic cigarette smokers. *Circulation* 94:19–25
38. The Alpha-Tocopherol Beta Carotene Cancer Prevention Study Group (1994) The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med* 330:1029–1035
39. van Poppel G, Poulsen H, Loft S, Verhagen H (1995) No influence of β -carotene on oxidative DNA damage in male smokers. *J Natl Cancer Inst* 87:310–311
40. Voorrips LE, Ravelli ACJ, Dongelmans PCA, Deurenberg P, van Staveren WA (1991) A physical activity questionnaire for the elderly. *Med Sci Sports Exerc* 23:974–979
41. Watkins ML, Erickson JD, Thun MJ, Mulinaire J, Heath CW (2000) Multivitamin use and mortality in a large prospective study. *Am J Epidemiol* 152:149–162
42. Welch RW, Turley E, Sweetman SF, Kennedy G, Collins AR, Dunne A, Livingstone MB, McKenna PG, McKelvey-Martin VJ, Strain JJ (1999) Dietary antioxidant supplementation and DNA damage in smokers and non-smokers. *Nutr Cancer* 34:167–172