

Erratum to: Carbon and nitrogen isotopic ratios of urine and faeces as novel nutritional biomarkers of meat and fish intake

Gunter G. C. Kuhnle · Annemiek M. C. P. Joosen ·
Catherine J. Kneale · Tamsin C. O'Connell

Published online: 5 December 2012
© Springer-Verlag Berlin Heidelberg 2012

Erratum to: Eur J Nutr DOI 10.1007/s00394-012-0328-2

In the original publication, Table 2 was published incorrectly as the values for urine and faeces were inadvertently interchanged. The corrected table is given below.

The online version of the original article can be found under doi:[10.1007/s00394-012-0328-2](https://doi.org/10.1007/s00394-012-0328-2).

G. G. C. Kuhnle (✉)
Department of Food and Nutritional Sciences, University of
Reading, Whiteknights, PO Box 226, Reading RG6 6AP, UK
e-mail: g.g.kuhnle@reading.ac.uk

G. G. C. Kuhnle · A. M. C. P. Joosen
MRC Dunn Human Nutrition Unit, Wellcome Trust/MRC
Building, Cambridge, UK

G. G. C. Kuhnle
MRC Centre for Nutritional Epidemiology in Cancer Prevention
and Survival, Department of Public Health and Primary Care,
University of Cambridge, Cambridge, UK

C. J. Kneale · T. C. O'Connell
McDonald Institute for Archaeological Research,
University of Cambridge, Cambridge, UK

T. C. O'Connell
Department of Archaeology and Anthropology,
University of Cambridge, Cambridge, UK

Table 2 Isotope ratios ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of blood, faeces and urine (median and inter-quartile range)

Diet group	<i>n</i>	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)
Blood ^a			
Fish	13	−22.7 (−23.0 to −22.6)	8.5 (8.4–8.6)
Meat/fish	11	−23.0 (−23.2 to −22.6)	8.5 (8.3–8.7)
Meat	14	−22.8 (−23.1 to −22.5)	8.5 (8.3–8.7)
Vegetarian	–	–	–
Urine ^b			
Fish	14	−23.2 (−23.4 to −22.9)	6.7 (6.3–6.8)
Meat/fish	13	−24.0 (−24.3 to −23.5)	5.7 (5.2–6.0)
Meat	14	−24.3 (−24.4 to −23.8)	5.6 (5.5–5.9)
Vegetarian	4	−24.8 (−25.0 to −24.5)	3.5 (3.4–3.8)
Faeces ^b			
Fish	14	−25.8 (−26.2 to −25.5)	7.8 (7.3–8.1)
Meat/fish	13	−26.4 (−26.6 to −26.2)	7.0 (6.8–7.8)
Meat	14	−26.5 (−26.7 to −26.3)	6.9 (6.7–7.2)
Vegetarian	4	−27.2 (−27.4 to −27.0)	5.0 (4.5–5.4)

^a No significant differences between diets ($p > 0.6$ for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, Kruskal–Wallis test)

^b Significant differences between diets ($p < 0.0005$ for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, Kruskal–Wallis test)