

Mannich and *O*-Alkylation Reactions of Tetraalkoxyresorcin[4]arenes – The Use of Some Products in Ligand-Assisted Reactions

Benjamin R. Buckley,^[a] Jonathan Y. Boxhall,^[a] Philip C. Bulman Page,^[a] Yohan Chan,^[a] Mark R. J. Elsegood,^[a] Harry Heaney,^{*[a]} Kathryn E. Holmes,^[a] Matthew J. McIldowie,^[b] Vickie McKee,^[a] Matthew J. McGrath,^[a] Mauro Mocerino,^[b] Andrew M. Poulton,^[a] Edward P. Sampler,^[a] Brian W. Skelton,^[c] and Allan H. White^[c]

Keywords: Axial chirality / Chiral auxiliaries / Configuration determination / Diastereoselectivity / Mannich reactions / Resorcinarenes

In the original article,^[1] the structure of **7** (Figure 5 on page 5123) is inadvertently displayed again in Figure 8 on page 5125 instead of the structure of **20a** as stated in the caption. The correct Figure 8 is shown below.




Figure 8. X-ray structure of amide **20a**; the majority of the hydrogen atoms are omitted for clarity.

The Editors

- [1] B. R. Buckley, J. Y. Boxhall, P. C. B. Page, Y. Chan, M. R. J. Elsegood, H. Heaney, K. E. Holmes, M. J. McIldowie, V. McKee, M. J. McGrath, M. Mocerino, A. M. Poulton, E. P. Sampler, B. W. Skelton, A. H. White, *Eur. J. Org. Chem.* **2006**, 5117–5134.

Received: November 29, 2006
Published Online: January 16, 2007

[a] Department of Chemistry, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK
E-mail: h.heaney@lboro.ac.uk

[b] Department of Applied Chemistry, Curtin University of Technology, GPO Box U 1987, Perth WA 6845, Australia

[c] Department of Chemistry, University of Western Australia, Crawley WA 6009, Australia