

A Novel Route to a Bromo-Cyano-Substituted Azulene and Its Exploitation in the Construction of an Acetylenic Scaffold

Michael Åxman Petersen,^[a] Kristine Kilså,^[a] Anders Kadziola,^[a] and
Mogens Brøndsted Nielsen^{*[a]}

Keywords: Acetylenic scaffolding / Alkynes / Azulenes / Chromophores / Cross-coupling

A novel route to functionalized azulenes is devised from a dihydroazulene precursor. Thus, bromination of 1,1-dicyano-2-phenyl-1,8a-dihydroazulene followed by heating in the presence of bromide ions provides an efficient way to generate 3-bromo-1-cyano-2-phenylazulene. Formation of this somewhat unexpected product was confirmed by X-ray crystallographic analysis. It undergoes a palladium-catalyzed


cross-coupling reaction with trimethylsilylacetylene, affording a new azulene building block for acetylenic scaffolding. Oxidative homo-coupling hereof provides an azulene dimer, for which the optical and electrochemical properties are compared to the other azulenes.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2007)

Introduction

Derivatives of azulene (**1**) are interesting building blocks for advanced materials with electronic and photonic applications.^[1] The azulene system has a remarkable polarizability and a tendency to form a stabilized tropylium cation as well as a cyclopentadienyl anion, which may be enhanced by suitable functionalization by donor and acceptor groups.^[2] Cyano-substituted azulenes have proven as good electron acceptors that may be employed in organic metals. Thus, Hafner and co-workers^[3] showed that the electron donor tetrathiafulvalene forms a charge-transfer complex with 2,4,6,8-tetracyanoazulene (**2**). This compound was prepared from 2-cyanoazulene (**3**) by nucleophilic substitution reactions in the seven-membered ring, followed by hydrolysis and dehydrogenation steps. In contrast, the five-membered ring of azulene is reactive towards electrophiles, and mostly so at the 1- and 3-positions.^[4] For example, reaction of cyanogen bromide with azulene (**1**) in the presence of stannic chloride has provided 1-cyanoazulene (**4**).^[5] However, with a ten-fold excess of the cyanogen bromide–stannic chloride complex, 1,3-dibromoazulene and 1-bromo-3-cyanoazulene (**5**) were obtained.^[5] Interestingly, it was observed that compound **4** was inert to an excess of cyanogen bromide and stannic chloride and thus not able to act as a precursor for **5** under these conditions. Cyanoazulene derivatives such as **5** containing a reactive bromide substituent are potential precursors for larger conjugated electron-accepting scaffolds. Thus, it has recently been

shown that halide-functionalized azulenes undergo the Sonogashira cross-coupling reaction^[6] with alkynes and that the resulting ethynylazulenes are efficient building blocks for acetylenic scaffolding^[7] as well as for metathesis.^[8]




We became interested to exploit the possibility for dihydroazulenes, such as **6**, to act as precursors for cyano-bromo-substituted azulenes. Dihydroazulenes have attracted attention as photoswitches as they undergo, after light irradiation, a 10-electron *retro*-electrocyclization to the isomeric vinylheptafulvene compounds, which, in turn, undergo a thermal cyclization back to the dihydroazulene forms.^[9,10] Gierisch and Daub^[9] showed that dihydroazulenes could in fact also be converted into the corresponding 1-cyanoazulenes, albeit in low yields. We report here a new efficient synthetic procedure for obtaining a 3-bromo-1-cyano-substituted azulene from a dihydroazulene precursor and its further reactivity towards alkynes, providing new electron acceptors and chromophores.

Results and Discussion

Our synthesis starts from a 2-phenyl-substituted derivative of **6**, namely compound **7** (Scheme 1) that was readily

[a] Department of Chemistry, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen Ø, Denmark
E-mail: mbn@kiku.dk

Supporting information for this article is available on the WWW under <http://www.eurjoc.org> or from the author.




Scheme 1.

prepared according to a general literature protocol.^[11] We reckoned from calculations (vide infra) that bromination of this compound should give preferably the 7,8-dibromo compound **8**. Indeed, treating **7** with bromine gave **8** in quantitative yield.^[12] Single crystals of **8** were grown from CH_2Cl_2 /pentane and subjected to an X-ray crystallographic analysis, which confirmed the proposed structure (Figure 1). When a 0.05 M solution of **8** in CH_2Cl_2 was left overnight at 40 °C, it was converted to a mixture of the two azulenes **9** and **10**^[13] formed in a ratio of 4:7 and in a total yield of 61%. In contrast, heating simply the dihydroazulene **7** overnight at 40 °C in CH_2Cl_2 caused no azulene formation. Heating **8** in the presence of one equivalent of bromide (Bu_4NBr) gave solely compound **9** in a yield of 79%.^[14] Yet, running the reaction under more dilute conditions (0.009 M) and in the absence of Bu_4NBr produced solely the cyanoazulene **10** in a yield of 72%. While exploitation of 1-cyanoazulene (**4**) as a precursor for 1-bromo-3-cyanoazulene (**5**) was previously discarded,^[5] we find that treating azulene **10** with a small excess of Br_2 in CH_2Cl_2 resulted in clean conversion to the bromide **9**. These observations suggest that at least two mechanisms can be responsible for the formation of **9** from the precursor **8**: *i*) initial elimination of either HBr or HCN , followed by nucleophilic attack of bromide at C3 with expulsion of either cyanide or bromide ($\text{S}_{\text{N}}2'$), followed by a final elimination reaction, or *ii*) initial formation of **10** after two elimination reactions, followed by electrophilic substitution by attack of either Br_2 or BrCN . The structures of both **9** and **10** were confirmed by X-ray crystallography (Figure 1).^[15]

The formation of azulenes **9** and **10** from dibromide **8** can also be promoted by light irradiation. Thus, photolysis at 350 nm of a dilute sample of **8** (0.009 M) in dry CH_2Cl_2 (1.5 mL; without stabilizer) for 1 hour showed complete conversion to the cyanoazulene **10** (as judged from an ^1H NMR spectroscopic investigation of the reaction mixture after evaporation of the solvent), while photolysis for 1 hour of **8** (0.009 M) in the presence of Bu_4NBr (0.05 M) instead provided the bromide **9** (together with a minor unidentified by-product).

The first step of the bromination of dihydroazulene **7** was subjected to a computational study employing the Gaussian 03 program package^[16] in order to compare stabilities of possible cations. The isomeric ions were optimized at the semiempirical PM3 level, and on each structure a frequency

Figure 1. X-ray crystal structures of **8**, **9**, and **10**.


analysis was performed to secure that a real minimum had been obtained. Then single point energy calculations were performed at the B3LYP/6-311+G(2d,p) level. The relative energies (at 0 K) of the ions are depicted in Figure 2. As

Isomer	Relative energy (kcal mol ⁻¹)
"8-Br"	0
"7-Br"	+28.1
"6-Br"	+7.8
"5-Br"	+32.2
"4-Br"	+12.2
"3a-Br"	+28.2
"3-Br"	+25.5
"2-Br"	+22.0


Figure 2. Relative cation energies calculated at the B3LYP/6-311+G(2d,p) level on PM3-optimized structures.

expected from simple resonance formulas, an attack at C8 is the most favorable. Electrophilic attack of Br^+ at for example the 2- or 3-positions requires more than 20 kcalmol⁻¹ relative to attack at C8. For the subsequent attack of bromide, we find that the kinetic product of addition (assumed so from a proximity effect), namely the 7,8-dibromide **8** (7*S*,8*S*,8*aR* stereoisomer), is also the thermodynamic product as the 2,8-, 3*a*,8-, and 5,8-dibromides are more energetic by +17.9, +25.3, and +7.1 kcalmol⁻¹, respectively. All in all, the calculations substantiate the exclusive formation of the dibromide **8** from bromination of **7**.

The bromide **9** was subjected to a Sonogashira cross-coupling reaction^[6] with trimethylsilylacetylene employing the catalyst system of Hundertmark et al.^[17] (Scheme 2). The product **11**^[18] was desilylated with K_2CO_3 in MeOH/THF, and the terminal alkyne intermediate was then subjected to an oxidative Hay homo-coupling reaction^[19] to give the azulene dimer **12**.^[20] Single crystals of **12** were grown from toluene/ CHCl_3 , and the crystal structure is shown in Figure 3. Angles of 22° and 38° are observed between the phenyl and azulene rings.




Scheme 2.

Figure 3. Crystal structure of **12**.

The new azulene derivatives are strong chromophores as is evident from the UV/Vis absorption spectra shown in Figure 4. The azulene monomers **9** and **10** exhibit a bright violet and blue-violet color, respectively, in solution. They are only weakly fluorescent with structured emissions at 433 nm ($\phi_{\text{fl}} = 0.5\%$) for **9** and 431 nm ($\phi_{\text{fl}} = 0.4\%$) for **10** in CH_2Cl_2 ($\lambda_{\text{exc}} = 322$ nm). Compounds **11** and **12** are both

green in the solid state, which is also the case in solution for the dimer **12**, while the monomer **11** exhibits a bright blue color in solution. Cyclic voltammetry investigations in CH_2Cl_2 (0.1 M Bu_4NPF_6) show irreversible reductions at -1.23 (**9**), -1.38 (**10**), -1.80 (**11**), and -1.81 (**12**) V vs. Fc^+/Fc . Taking into account the electron-accepting nature of acetylenic scaffolds,^[21] it is somewhat surprising that compounds **11** and **12** are the poorest acceptors in the series.

Figure 4. UV/Vis absorption spectra in CH_2Cl_2 .

In conclusion, we have developed a new efficient synthesis of a 3-bromo-functionalized azulene. This compound is readily incorporated into new redox-active acetylenic chromophores. We are currently investigating the possibility for controlling the light-induced conversion of **8** into azulenes via a ring-opened intermediate.

Supporting Information (see also the footnote on the first page of this article): A Table containing a summary of general crystallographic data.

CCDC-626060 (for **8**), -626062 (for **9**), -626061 (for **10**), -626063 (for **12**) contain the detailed supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/datarequest/cif.

Acknowledgments

We thank the Danish Research Agency (grant #2111-04-0018) for financial support.

- [1] a) G. Iftime, P. G. Lacroix, K. Nakatani, A. C. Razus, *Tetrahedron Lett.* **1998**, 39, 6853–6856; b) S. Ito, S. Kikuchi, N. Morita, T. Asao, *J. Org. Chem.* **1999**, 64, 5815–5821; c) P. G. Lacroix, I. Malfant, G. Iftime, A. C. Razus, K. Nakatani, J. A. Delaire, *Chem. Eur. J.* **2000**, 6, 2599–2608; d) S. Ito, H. Inabe, N. Morita, K. Ohta, T. Kitamura, K. Imafuku, *J. Am. Chem. Soc.* **2003**, 125, 1669–1680; e) F. Wang, Y.-H. Lai, M. Y. Han, *Macromolecules* **2004**, 37, 3222–3230.
- [2] H. Möllerstedt, M. C. Piqueras, R. Crespo, H. Ottosson, *J. Am. Chem. Soc.* **2004**, 126, 13938–13939.
- [3] S. Schmitt, M. Baumgarten, J. Simon, K. Hafner, *Angew. Chem. Int. Ed.* **1998**, 37, 1077–1081.
- [4] H. Spiesecke, W. G. Schneider, *Tetrahedron Lett.* **1961**, 2, 468–472.
- [5] A. G. Anderson Jr, D. J. Gale, R. N. McDonald, R. G. Anderson, R. C. Rhodes, *J. Org. Chem.* **1964**, 29, 1373–1377.
- [6] K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* **1975**, 16, 4467–4470.
- [7] a) S. Ito, H. Inabe, T. Okujima, N. Morita, M. Watanabe, N. Harada, K. Imafuku, *J. Org. Chem.* **2001**, 66, 7090–7101; b)

- M. Lamberto, C. Pagba, P. Piotrowiak, E. Galoppini, *Tetrahedron Lett.* **2005**, 46, 4895–4899; c) K. K. H. Fabian, A. H. M. Elwahy, K. Hafner, *Eur. J. Org. Chem.* **2006**, 791–802.
- [8] A. Mikus, V. Sashuk, M. Kędziorek, C. Samojłowicz, S. Ostrowski, K. Grela, *Synlett* **2005**, 1142–1146.
- [9] S. Gierisch, J. Daub, *Chem. Ber.* **1989**, 122, 69–75.
- [10] L. Gobbi, P. Seiler, F. Diederich, *Angew. Chem. Int. Ed.* **1999**, 38, 674–678.
- [11] L. Gobbi, P. Seiler, F. Diederich, V. Gramlich, C. Boudon, J.-P. Gisselbrecht, M. Gross, *Helv. Chim. Acta* **2001**, 84, 743–777.
- [12] Dihydroazulene **7** (125 mg, 0.49 mmol) was dissolved in CH_2Cl_2 (4 mL) at -78°C under N_2 . Then a solution of Br_2 in CH_2Cl_2 (0.39 M, 1.25 mL) was slowly added (2 min). Stirring for 1 h at -78°C followed by evaporation in vacuo yielded **8** (205 mg, 100%) as a yellow-brown foam. An analytical sample was recrystallized from CH_2Cl_2 and pentane. Melting range 170 – 178°C , compound turns dark-violet above 95°C . $R_f = 0.36$ in EtOAc/heptane, 3:7 v/v, 2D-TLC showed breakdown of main product; decomposition product observed at $R_f = 0.075$. ^1H NMR (300 MHz, CDCl_3 , 25°C , 7.26 ppm): $\delta = 7.76$ (m, 2 H, Ar), 7.48 (m, 3 H, Ar), 6.98 (s, 1 H, CH), 6.27 [dd, $^3J(\text{H,H}) = 2.2$ Hz, $^3J(\text{H,H}) = 7.5$ Hz, 1 H, CH], 6.09 [dd, $^3J(\text{H,H}) = 7.5$ Hz, $^3J(\text{H,H}) = 12.2$ Hz, 1 H, CH], 5.91 [dd, $^3J(\text{H,H}) = 5.6$ Hz, $^3J(\text{H,H}) = 12.0$ Hz, 1 H, CH], 5.32 (m, 1 H, CH), 5.05 (m, 1 H, CH), 4.66 (br. s, 1 H, CH) ppm. ^{13}C NMR (75 MHz, CDCl_3 , 25°C , 77.0 ppm): $\delta = 144.5$, 139.6, 133.9, 130.3, 130.0, 129.3, 128.5, 126.4, 125.7, 121.0, 114.6, 111.7, 53.1, 51.5, 49.0, 44.6 ppm. UV/Vis (CH_2Cl_2): λ_{max} (ϵ) = 348 nm (26700).
- [13] Compound **10** was previously prepared in another manner: T. Nozoe, K. Takase, S. Fukuda, *Bull. Chem. Soc. Jpn.* **1971**, 44, 2210–2213.
- [14] Dibromide **8** (208 mg, 0.50 mmol) was dissolved in dry, unstabilized CH_2Cl_2 (10 mL) under N_2 atmosphere, whereupon NBu_4Br (171 mg, 0.53 mmol) was added. After stirring at 40°C for 16 h (caution: HCN evolution expected), the dark-purple reaction mixture was washed with water (2×30 mL), dried (MgSO_4), and the solvents evaporated in vacuo. Purification by flash column chromatography (EtOAc/heptane, 3:7, v/v, $R_f = 0.31$) yielded 3-bromo-1-cyano-2-phenylazulene **9** (121 mg, 79%), as a purple solid. M.p. 191 – 194°C . Crystals for X-ray crystallography were grown from CDCl_3 . ^1H NMR (300 MHz, 25°C , CDCl_3 , 7.26 ppm): $\delta = 8.65$ [d, $^3J(\text{H,H}) = 10.0$ Hz, 1 H, CH=], 8.62 [d, $^3J(\text{H,H}) = 10.8$ Hz, 1 H, CH=], 7.93–7.82 (m, 3 H, Ar, CH=), 7.68–7.50 (m, Ar, 5 H, CH=) ppm. ^{13}C NMR (75 MHz, 25°C , CDCl_3 , 77.0 ppm): $\delta = 150.7$, 143.5, 140.3, 139.2, 138.2, 136.5, 133.0, 130.2, 129.4, 128.6, 128.5, 128.2, 116.7, 104.4, 96.5 ppm; ES-TOF-MS (neg. mode): $m/z = 308$ [H^-M^-], 339 [MeOH^-M^-]. UV/Vis (CH_2Cl_2): λ_{max} (ϵ) = 269 (13000), 321 (46500), 359 (7500), 373 (7500), 588 (500) nm; fluorescence (CH_2Cl_2 , $\lambda_{\text{exc}} = 322$ nm): $\lambda_{\text{em}} = 414$ (sh), 433 (tail, $\phi = 0.5\%$) nm.
- [15] Compound **9** crystallizes in the acentric space group Cc but does so as a racemic twin with a ratio of 52:48. Moreover, the azulene ring displays disorder, so that the Br and CN substituents are interchanged in 17% of the unit cells. The azulene ring and the phenyl ring form an angle of 41° . Crystals of compound **10** contain two molecules in the asymmetric unit. One of the molecules is all planar (shown in Figure 1), induced by the crystal packing, and disordered. Here 38% of the molecules are rotated 180° with the phenyl ring overlapping the seven-membered ring of the azulene, and the five-membered rings of the azulene partially overlapping. The other molecule is well ordered and displays an angle of 33° between the phenyl ring and the azulene ring.
- [16] Gaussian 03, Revision B.03, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery Jr, T. Vreven, K. N. Kudin, J. C. Burant, J. M. Millam, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rega, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dannenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, J. Cioslowski, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, C. Gonzalez, and J. A. Pople, Gaussian, Inc., Pittsburgh PA, **2003**.
- [17] T. Hundertmark, A. F. Littke, S. L. Buchwald, G. C. Fu, *Org. Lett.* **2000**, 2, 1729–1731.
- [18] Compound **9** (200 mg, 0.65 mmol) was dissolved (with sonication) in dry THF (1.5 mL), dry toluene (1.5 mL), and $\text{NH}(\text{iPr})_2$ (0.2 mL) under argon. Then $[\text{Pd}(\text{PhCN})_2\text{Cl}_2]$ (33 mg, 0.09 mmol), CuI (20 mg, 0.1 mmol), and $\text{P}(\text{iBu})_3$ in hexane (10%, 0.25 mL, 0.08 mmol) were added. Trimethylacetylene (0.4 mL, 2.7 mmol) was added, and the reaction mixture was allowed to stir at room temp. for 3 h. The resulting dark blue solution was diluted with CH_2Cl_2 (25 mL) and washed with water (2×50 mL), dried (MgSO_4), and the solvents evaporated in vacuo. Purification by flash column chromatography (EtOAc/heptane, 2:8, v/v, $R_f = 0.4$) gave **11** as a green solid (150 mg, 79%). M.p. 112 – 113.5°C . ^1H NMR (300 MHz, CDCl_3 , 25°C , 7.26 ppm): $\delta = 8.71$ [d, $^3J(\text{H,H}) = 9.7$ Hz, 1 H, CH=], 8.63 [d, $^3J(\text{H,H}) = 9.7$ Hz, 1 H, CH=], 8.15 (m, 2 H, Ar), 7.85 [t, $^3J(\text{H,H}) = 9.8$ Hz, 1 H, CH=], 7.64–7.49 (m, 5 H, Ar, CH=), 0.30 (s, 9 H, CH_3) ppm. ^{13}C NMR (75 MHz, CDCl_3 , 25°C , 77.0 ppm): $\delta = 153.6$, 144.4, 144.1, 140.1, 138.1, 136.4, 133.3, 129.9, 129.6, 129.0, 129.0, 128.5, 117.2, 109.8, 102.3, 99.3, 95.6, 0.0 ppm. HR-ES-TOF-MS (pos. mode): $m/z = 348.1193$ [M^+Na^+]; calcd. for $\text{C}_{22}\text{H}_{19}\text{NNaSi}$: 348.1184. UV/Vis (CH_2Cl_2): λ_{max} (ϵ) = 273 (22000), 289 (22000), 328 (47000), 337 (51000), 372 (8000), 390 (br. sh, 7000), 601 (500) nm.
- [19] A. S. Hay, *J. Org. Chem.* **1962**, 27, 3320–3321.
- [20] Compound **11** (72 mg, 0.22 mmol) was dissolved in MeOH (3 mL) and THF (1.5 mL). Then K_2CO_3 (160 mg, 1.15 mmol) was added, and the reaction mixture was stirred for 1/2 h at room temp. and then diluted with CH_2Cl_2 . The organic phase was washed with water, dried (MgSO_4), and the solvents evaporated in vacuo. The dark-blue solid was dissolved in CH_2Cl_2 (15 mL), whereupon Hay catalyst [TMEDA (40 mg, 0.34 mmol) and CuCl (40 mg, 0.40 mmol) in CH_2Cl_2 (1 mL)] was added. After stirring for 3 1/2 h at room temp., the reaction mixture was washed with water, dried (MgSO_4), and the solvents evaporated in vacuo. Purification by flash column chromatography (CH_2Cl_2 , $R_f = 0.44$) followed by two procedures of precipitation, by dissolving in hot CHCl_3 followed by addition of heptane, yielded the dimer **12** (30 mg, 27%) as a green solid. M.p. 315 – 317°C (dec.). Crystals for X-ray crystallography were grown from toluene/ CHCl_3 . ^1H NMR (300 MHz, CDCl_3 , 25°C , 7.26 ppm): $\delta = 8.79$ [d, $^3J(\text{H,H}) = 10.0$ Hz, 2 H, CH=], 8.66 [d, $^3J(\text{H,H}) = 10.0$ Hz, 2 H, CH=], 8.13 [d, $^3J(\text{H,H}) = 7.6$ Hz, 4 H, Ar], 7.90 [t, $^3J(\text{H,H}) = 9.4$ Hz, 2 H, CH=], 7.69–7.52 (m, 10 H, CH=, Ar) ppm. ^{13}C NMR (75 MHz, CDCl_3 , 50°C , 77.0 ppm): $\delta = 140.5$, 138.4, 136.7, 129.8, 129.7, 129.4, 128.9 ppm, (missing signals due to poor solubility). MALDI-TOF-MS (neg. mode): $m/z = 504$ [M^-], 1008 [M^-M^-]. UV/Vis (CH_3Cl): λ_{max} (ϵ) = 278 (37500), 325 (71500), 345 (66500), 420 (16000), 454 (14000), 583 (1300) nm.
- [21] J.-P. Gisselbrecht, N. N. P. Moonen, C. Boudon, M. B. Nielsen, F. Diederich, M. Gross, *Eur. J. Org. Chem.* **2004**, 2959–2972.

Received: December 4, 2006

Published Online: February 9, 2007