


Novel Sorbicillin Derivatives with an Unprecedented Carbon Skeleton from the Sponge-Derived Fungus *Trichoderma* Species

Kerstin Neumann,^[a] Ahmed Abdel-Lateff,^[a] Anthony D. Wright,^[b] Stefan Kehraus,^[a]
Anja Krick,^[a] and Gabriele M. König^{*[a]}

Keywords: Natural products / Sorbicillin derivatives / Structure elucidation / Trichodermanones / Configuration determination

In Figure 1 on page 2268 of the original article,^[1] the structures of compounds **5** and **8** were inadvertently drawn incorrectly; the correct structures are given below.



Additionally, bislongiquinolide (**7**) was first isolated and named by Ayer et al.^[2]

The Authors

- [1] K. Neumann, A. Abdel-Lateff, A. D. Wright, S. Kehraus, A. Krick, G. M. König, *Eur. J. Org. Chem.* **2007**, 2268–2275.
[2] R. Andrade, W. A. Ayer, L. S. Trifonov, *Austr. J. Chem.* **1997**, *50*, 255.

Received: June 13, 2007
Published Online: July 10, 2007

[a] Institute for Pharmaceutical Biology, University of Bonn,
Nussallee 6, 53115 Bonn, Germany
E-mail: g.koenig@uni-bonn.de

[b] Australian Institute of Marine Science, PMB No. 3,
Townsville MC, Qld 4810, Australia