

A Convenient and General Method for the Synthesis of Indole-2,3-dicarboxylates and 2-Arylindole-3-carboxylates

Ilyas Ali Sayyed,^[a] Karolin Alex,^[a] Annegret Tillack,^[a] Nicolle Schwarz,^[a] Dirk Michalik,^[a] and Matthias Beller*^[a]

Keywords: Alkynes / Hydroamination / Indoles

A transition-metal-free, simple and efficient one-pot method for the synthesis of indole-2,3-dicarboxylates and 2-arylindole-3-carboxylates is described. The corresponding products are obtained by a domino hydroamination/Fischer indole cyclization in good-to-excellent yields from easily

available 1-alkyl-1-phenylhydrazines and acetylene carboxylates.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2007)

The catalytic addition of organic amines and their derivatives to alkenes and alkynes (hydroamination) to produce nitrogen-containing molecules is of significant importance for synthetic chemists in basic research as well as for those in the chemical industry.^[1] Today intermolecular hydroaminations are known to be catalyzed by a variety of transition metals (d- and f-block metals),^[2] alkali metals^[3] and Brønsted and Lewis acids.^[4] However, most of these catalysts are rather limited with respect to their substrate tolerance.

Various biologically active amine alkaloids, especially indoles, continue to attract the interest of organic chemists. Though many catalytic methods exist for the synthesis of indoles,^[5] still the most famous synthesis for indoles and their derivatives constitutes the Fischer indole synthesis.^[6] We have been interested for some time in the improvement and exploration of methodologies for the synthesis of indole heterocycles.^[7] For example, we developed a one-pot synthesis of tryptamines and tryptophols by a titanium-catalyzed hydrohydrazination of chloro- and silyloxy-substituted alkynes.^[7a,8] In continuation of this work, we became interested in the hydrohydrazination reaction of acetylenedicarboxylates, which are easily available. Some related carboxy-2,3-disubstituted indole derivatives are known to be potent inhibitors of thromboxane synthase,^[9] phospholipase-A₂,^[10] cyclooxygenase-2,^[11] steroid-5 α -reductase^[12] and glycine/NMDA antagonists.^[13] In addition to their biological activity, indole-2,3-dicarboxylate esters may serve as valuable synthetic intermediates for other indole derivatives and more complex indole heterocycles.^[14]

On the basis of our previous hydroamination protocols, we initially investigated the reaction of *N*-methyl-*N*-phenyl-

hydrazine (**1a**) with diethyl acetylenedicarboxylate (**2**) in the presence of 10 mol-% of Ti(NEt₂)₄ as the catalyst at 100 °C. Subsequent treatment of the reaction mixture with 3 equiv. of ZnCl₂ allowed the cyclization of the in-situ-generated hydrazone to give the corresponding diethyl 1-methylindole-2,3-dicarboxylate **3a** in 55% yield (Table 1, Entry 1).

Table 1. Reaction of 1-methyl-1-phenylhydrazine (**1a**) with diethyl acetylenedicarboxylate (**2**).^[a]

Entry	Hydrazine 1a [equiv.]	Alkyne 2 [equiv.]	Yield 3a [%]
1 ^[b]	1.0	1.0	55
2	1.0	1.0	63
3 ^[c]	1.0	1.0	30
4 ^[d]	1.0	1.0	60
5	1.5	1.0	66
6	1.0	1.5	78
7	1.0	2.0	87

[a] Reaction conditions: i) toluene, 100 °C, 24 h; ii) ZnCl₂ (3 equiv.), 100 °C, 24 h. [b] 10 mol-% Ti(NEt₂)₄, 20 mol-% 2,6-di-*tert*-butyl-4-methylphenol. [c] Reaction temperature 80 °C. [d] Reaction temperature 120 °C.























Interestingly, the same reaction occurred in 63% yield when performed in the absence of any titanium catalyst (Table 1, Entry 2). This observation is in agreement with the previous work of Acheson^[15] and Miki et al.^[16] who prepared selected dimethyl indole-2,3-dicarboxylates in yields of 13–62%. Interestingly, in 1935 Diels and Reese had already described the condensation of 1-benzyl-1-phenylhydrazine with acetylenedicarboxylate; however, no product yield of the respective indole-2,3-dicarboxylate was given.^[17]

[a] Leibniz-Institut für Katalyse e.V. an der Universität Rostock, Albert-Einstein-Str. 29a, 18059 Rostock, Germany
E-mail: matthias.beller@catalysis.de

Because of the limited information available, we decided to perform a systematic study to improve the yield of this domino hydroamination/cyclization sequence. Selected experiments are shown in Table 1. The use of an excess amount of phenylhydrazine **1a** (1.5 equiv.) improved the yield of **3a** slightly. In contrast, a considerable increase in

the yield of the corresponding indoledicarboxylate was observed when an excess of alkyne **2** was used. Hence, 1.5 and 2.0 equiv. of **2** gave 78 and 87% yield of **3a**, respectively. This is explained by oligomerization side reactions of **2**. We also examined the effect of temperature to improve the yield of the indole product. However, neither an increase in the

Table 2. Synthesis of indole-2,3-dicarboxylates.^[a]

Entry	Hydrazine	Alkyne	Product	Yield [%] ^[b]
1	 1a	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3a	87
2	 1b	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3b	67
3	 1c	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3c	60
4	 1d	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3d	65
5	 1e	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3e	60
6	 1f	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3f	55
7	 1g	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3g	75
8	 1h	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3h	40
9	 1i	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{CO}_2\text{Et}$	 3i	26
10	 1j	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{Ph}$	 3j	55
11	 1k	$\text{EtO}_2\text{C}-\text{C}\equiv\text{C}-\text{Ph}$	 3k	60


[a] Reaction conditions: i) hydrazine (1 equiv.), acetylenedicarboxylate (2 equiv.), toluene, 100 °C, 24 h; ii) ZnCl_2 (3 equiv.), 100 °C, 24 h.

[b] Yield of isolated products.

reaction temperature to 120 °C nor a lowering to 80 °C gave a better yield of **3a**. Also, attempts to improve the yield further by using different solvents such as benzene, methanol, THF, 1,4-dioxane and NMP were not fruitful. Noteworthy is that when ZnCl_2 was added in the beginning of reaction, the desired product was also obtained, albeit in lower yield.

As already described by Acheson et al.^[15] we were also able to isolate the intermediate hydrazone of the reaction between *N*-methyl-*N*-phenylhydrazine (**1a**) and acetylenedicarboxylate **2** in the absence of ZnCl_2 . After 24 h at 100 °C in toluene, a hydrazone/indole ratio of 83:17 was found. The hydrazone immediately underwent Fischer indole cyclization in the presence of the Lewis acid. At this point it is noteworthy that Schwesinger et al. demonstrated that the reaction of phenylhydrazine and acetylenedicarboxylate also yielded the corresponding hydrazone.^[18] However, as a result of the strong intramolecular hydrogen bond, subsequent Fischer indole cyclization to yield the indole-2,3-dicarboxylate was prevented. Apparently, the presence of two substituents at the hydrazine nitrogen atom facilitates the Fischer indole cyclization. Thus, we next applied the improved hydroamination–cyclization protocol to indole products **3a–k** (Table 2).

However, lower yields (26–40%) were obtained for the reaction of *N*-(4-bromophenyl)- and *N*-(4-chlorophenyl)-*N*-benzylhydrazine, presumably owing to the lower reactivity in the Fischer indole cyclization step. Notably, the methodology is also applicable to other activated acetylenedicarboxylic acid derivatives. Hence, the regioselective synthesis of 2-arylindolecarboxylate derivatives is possible from ethyl 3-phenyl-1-propyne-1-carboxylate in 55–60% yield (Table 2, Entries 10 and 11). To the best of our knowledge these are the first examples of such a direct synthesis of 2-arylindole-3-carboxylates. Moreover, by applying the more labile ethyl propiolate, 15% of the desired product was obtained under similar reaction conditions (Scheme 1). Here, the lower product yield can be explained by the increased propensity of ethyl propiolate to undergo self condensation.



Scheme 1. Reaction with ethyl propiolate.

In conclusion, we presented the synthesis of various diethyl indole-2,3-dicarboxylates by a domino hydroamination/Fischer indole cyclization synthesis. The corresponding products are obtained easily from commercially available substrates in general in good yield. We believe that this methodology constitutes the most convenient access to this class of compounds.

Experimental Section

Representative Procedure: An Ace pressure tube under an argon atmosphere was charged with 1-methyl-1-phenylhydrazine (0.366 g,

3.0 mmol), diethyl acetylenedicarboxylate (1.02 g, 6.0 mmol) and dry toluene (5 mL). The pressure tube was fitted with a Teflon cap and heated at 100 °C for 24 h in an oil bath. Then, the reaction mixture was cooled to r.t. and anhydrous ZnCl_2 (1.22 g, 9.0 mmol) was added. The reaction mixture was further heated at 100 °C for 24 h. The excess toluene was distilled off under reduced pressure, and the residue was purified by chromatography on silica gel (hexane/ethyl acetate, 9:1) to afford **3a** as a gummy liquid. Isolated yield: 0.717 g (87%). ^1H NMR (500 MHz, CDCl_3): δ = 8.13 (br. d, $^3J_{4,5}$ = 8.0 Hz, 1 H, 4-H), 7.37–7.33 (m, 2 H, 6,7-H), 7.28 (m, 1 H, 5-H), 3.83 [s, 3 H, Me(8)], 4.48 [q, 3J = 7.3 Hz, 2 H, CH_2 (10)], 4.38 [q, 3J = 7.2 Hz, 2 H, CH_2 (12)], 1.42 [t, 3J = 7.3 Hz, 3 H, Me(10)], 1.40 [t, 3J = 7.2 Hz, 3 H, Me(12)] ppm. ^{13}C NMR (125.8 MHz, CDCl_3): δ = 164.1 (C-11), 162.8 (C-9), 136.8 (C-7a), 135.0 (C-2), 125.4 (C-3a), 124.3 (C-6), 122.3, 122.5 (C-4,5), 110.0 (C-7), 108.0 (C-3), 62.3 [CH_2 (10)], 60.2 [CH_2 (12)], 31.3 [Me(8)], 14.0 [Me(10)], 14.4 [Me(12)] ppm (numbering according to Table 2, Entry 1). IR (neat): $\tilde{\nu}$ = 3055, 2982, 2938, 2905, 1733, 1717, 1700, 1615, 1539, 1471, 1444, 1412, 1379, 1273, 1247, 1210, 1157, 1103, 1034, 1014, 860, 788, 753, 742 cm^{-1} . MS (EI, 70 eV): m/z (%) = 275 (100) [$\text{M}]^+$, 230 (40), 229 (30), 203 (28), 202 (91), 200 (20), 158 (18), 157 (22), 89 (6). HRMS (EI): calcd. for $\text{C}_{15}\text{H}_{17}\text{NO}_4$ 275.1153; found 275.1152.

Acknowledgments

This work has been supported by the State of Mecklenburg-Western Pomerania, the BMBF (Bundesministerium für Bildung und Forschung), the Deutsche Forschungsgemeinschaft (Leibniz-price), and the Fonds der Chemischen Industrie (FCI). We thank all members of the analytical group for their support.

- [1] a) A. L. Odom, *Dalton Trans.* **2005**, 225–233; b) K. C. Hultsch, *Adv. Synth. Catal.* **2005**, 347, 367–391; c) S. Hong, T. J. Marks, *Acc. Chem. Res.* **2004**, 37, 673–686; d) M. Beller, J. Seayad, A. Tillack, H. Jiao, *Angew. Chem. Int. Ed.* **2004**, 43, 3368–3398; e) S. Doye, *Synlett* **2004**, 1653–1672; f) J. F. Hartwig, *Pure Appl. Chem.* **2004**, 76, 507–516; g) P. W. Roesky, T. E. Müller, *Angew. Chem. Int. Ed.* **2003**, 42, 2708–2710; h) M. Beller, C. Breindl, M. Eichberger, C. G. Hartung, J. Seayad, O. R. Thiel, A. Tillack, H. Trauthwein, *Synlett* **2002**, 1579–1594; i) M. Nobis, B. Drießen-Hölscher, *Angew. Chem. Int. Ed.* **2001**, 40, 3983–3985; j) T. E. Müller, M. Beller, *Chem. Rev.* **1998**, 98, 675–703.
- [2] a) A. V. Lee, L. L. Schafer, *Organometallics* **2006**, 25, 5249–5254; b) M. L. Buil, M. A. Esteruelas, A. M. Lopez, A. C. Mateo, *Organometallics* **2006**, 25, 4079–4089; c) A. M. Johns, M. Utsunomiya, C. D. Incarvito, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, 128, 1828–1839; d) K. Takaki, S. Koizumi, Y. Yamamoto, K. Komeyama, *Tetrahedron Lett.* **2006**, 47, 7335–7337; e) J.-J. Brunet, N. C. Chu, O. Diallo, *Organometallics* **2005**, 24, 3104–3110; f) K. Marcsekova, B. Wegener, S. Doye, *Eur. J. Org. Chem.* **2005**, 4843–4851; g) A. Tillack, V. Khedkar, H. Jiao, M. Beller, *Eur. J. Org. Chem.* **2005**, 5001–5012; h) L. T. Kaspar, B. Fingerhut, L. Ackermann, *Angew. Chem. Int. Ed.* **2005**, 44, 5972–5974; i) H. Wang, H.-S. Chan, Z. Xie, *Organometallics* **2005**, 24, 3772–3779; j) B. D. Ward, A. Maise-Francois, P. Mountford, L. H. Gade, *Chem. Commun.* **2004**, 704–705; k) L. Ackermann, L. T. Kaspar, C. Gschrei, *Org. Lett.* **2004**, 6, 2515–2518.
- [3] a) V. Khedkar, A. Tillack, C. Benisch, J. P. Melder, M. Beller, *J. Mol. Catal. A: Chem.* **2005**, 241, 175–183; b) K. Kumar, D. Michalik, I. Garcia Castro, A. Tillack, A. Zapf, M. Arlt, T. Heinrich, H. Böttcher, M. Beller, *Chem. Eur. J.* **2004**, 10, 746–757; c) J. Seayad, A. Tillack, C. G. Hartung, M. Beller, *Adv. Synth. Catal.* **2002**, 344, 795–813.

- [4] a) K. Motokura, N. Nakagiri, K. Mori, T. Mizugaki, K. Ebitani, K. Jitsukawa, K. Kaneda, *Org. Lett.* **2006**, 8, 4617–4620; b) A. A. M. Lapis, B. A. DaSilveira Neto, J. D. Scholten, F. M. Nachtigall, M. N. Eberlin, J. Dupont, *Tetrahedron Lett.* **2006**, 47, 6775–6779; c) L. L. Anderson, J. Arnold, R. G. Bergman, *J. Am. Chem. Soc.* **2005**, 127, 14542–14543.
- [5] For reviews, see: a) G. R. Humphrey, J. T. Kuethe, *Chem. Rev.* **2006**, 106, 2875–2911; b) G. W. Gribble, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1045–1075; c) T. L. Gilchrist, *J. Chem. Soc. Perkin Trans. 1* **1999**, 2848–2866; d) R. J. Sundberg, *Indoles*, Academic Press, San Diego, **1996**; e) C. J. Moody, *Synlett* **1994**, 681–688; f) G. W. Gribble, *Contemp. Org. Synth.* **1994**, 1, 145–172; g) U. Pindur, R. Adam, *J. Heterocycl. Chem.* **1988**, 25, 1–8.
- [6] a) R. R. Gupta, *Heterocyclic Chemistry*, Springer, New York, **1999**, vol. 2, p. 199; b) A. R. Jones, *Comprehensive Heterocyclic Chemistry* (Eds.: A. R. Katritzky, C. W. Rees), Pergamon Press, Oxford, **1984**, vol. 4, p. 334; c) M. Nakazaki, K. Yamamoto, *J. Org. Chem.* **1976**, 41, 1877–1877; d) R. J. Sundberg, *The Chemistry of Indoles*, Academic Press, New York, **1970**, p. 142.
- [7] a) V. Khedkar, A. Tillack, M. Michalik, M. Beller, *Tetrahedron* **2005**, 61, 7622–7631; b) A. Tillack, H. Jiao, I. Garcia Castro, C. G. Hartung, M. Beller, *Chem. Eur. J.* **2004**, 10, 2409–2420; c) A. Mobaligh, R. Jackstell, M. Beller, *Tetrahedron Lett.* **2004**, 45, 869–873; d) K. Kumar, A. Zapf, D. Michalik, A. Tillack, M. Arlt, M. Beller, *Org. Lett.* **2004**, 6, 7–10; e) M. Beller, C. Breindl, T. H. Riermeier, M. Eichberger, H. Trauthwein, *Angew. Chem. Int. Ed.* **1998**, 37, 3389–3391.
- [8] a) V. Khedkar, A. Tillack, M. Michalik, M. Beller, *Tetrahedron Lett.* **2004**, 45, 3123–3126; b) for the first example of a Ti-catalyzed hydrohydrazination of alkynes, see: C. Cao, Y. Shi, A. L. Odom, *Org. Lett.* **2002**, 4, 2853–2856.
- [9] S. S. Bhagvat, C. Gude, *Tetrahedron Lett.* **1994**, 35, 1847–1850.
- [10] M. Lehr, Ger. Offen. DE 4 338 770, **1995**; *Chem. Abstr.* **1995**, 123, 32958.
- [11] C. Lau, C. Black, D. Guay, J.-Y. Gauthier, Y. Leblanc, P. Roy, Y. Ducharme, P. Hamel, PCT Int. Appl. WO 96 37 467, **1996**; *Chem. Abstr.* **1997**, 126, 89261.
- [12] K. Sawada, S. Okada, P. Golden, N. Kayakiri, Y. Sawada, M. Hashimoto, H. Tanaka, *Chem. Pharm. Bull.* **1999**, 47, 481–491.
- [13] a) M. Rowley, P. D. Leeson, S. Grimwood, A. Foster, K. Saywell, *Bioorg. Med. Chem. Lett.* **1992**, 2, 1627–1630; b) R. Di Fabio, A. M. Capelli, N. Conti, A. Cugola, D. Donati, A. Feriani, P. Gastaldi, G. Gaviraghi, C. T. Hewkin, F. Micheli, A. Missio, M. Mugnaini, A. Pecunioso, A. M. Quaglia, E. Ratti, L. Rossi, G. Tedesco, D. G. Trist, A. Reggiani, *J. Med. Chem.* **1997**, 40, 841–850.
- [14] a) A. Penoni, G. Palmisano, G. Brogini, A. Kadowaki, K. M. Nicholas, *J. Org. Chem.* **2006**, 71, 823–825; b) A. Penoni, J. Volkmann, K. M. Nicholas, *Org. Lett.* **2002**, 4, 699–701; c) Y. Liu, G. W. Gribble, *Tetrahedron Lett.* **2001**, 42, 2949–2951; d) L. Biocchi, M. Giannangeli, *J. Heterocycl. Chem.* **1988**, 25, 1905–1909; e) D. Black, N. Kumar, L. C. H. Wong, *Aust. J. Chem.* **1986**, 39, 15–20; f) T. Sakakibara, Y. Tanaka, S. Yamasaki, *Chem. Lett.* **1986**, 797–800; g) J. Duflos, G. Queguiner, *J. Org. Chem.* **1981**, 46, 1195–1198.
- [15] R. M. Acheson, J. M. Vernon, *J. Chem. Soc.* **1962**, 84, 1148–1157.
- [16] Y. Miki, K.-I. Matsushita, H. Hibino, H. Shirokoshi, *Heterocycles* **1999**, 51, 1585–1592.
- [17] O. Diels, J. Reese, *Justus Liebigs Ann. Chem.* **1935**, 519, 147–157.
- [18] H. Schwesinger, A. Dalski, D. Sicker, H. Wilde, G. Mann, *J. Prakt. Chem.* **1992**, 334, 257–264.

Received: April 5, 2007

Published Online: July 10, 2007