

Regioselective Arylations of α -Amido Sulfones with Electron-Rich Arenes through Friedel–Crafts Alkylations Catalyzed by Ferric Chloride Hexahydrate: Synthesis of Unsymmetrical and Bis-Symmetrical Triarylmethanes

Ponnaboina Thirupathi^[a] and Sung Soo Kim^{*[a]}

Keywords: Iron / C–C coupling / α -Amido sulfones / Arenes / Heteroarenes / Triarylmethanes

In the original article,^[1] the authors failed to cite parallel reports of their own research. The following sentence should be inserted on the first page, second column, line 9:

Very recently we described the Friedel–Crafts alkylation of α -amido sulfones catalyzed by Amberlyst-15^[6j] and InBr₃.^[6k]

The authors apologize for any inconvenience this may have caused.

The Editor

[1] P. Thirupathi, S. S. Kim, *Eur. J. Org. Chem.* **2010**, 1798–1808.

[6] j) S. T. Kadam, P. Thirupathi, S. S. Kim, *Tetrahedron* **2009**, *65*, 10383–10389; k) P. Thirupathi, S. S. Kim, *J. Org. Chem.* **2009**, *74*, 7755–7761.

Received: January 28, 2010

Published Online: February 16, 2010

[a] Department of Chemistry, Inha University,
Incheon 402-751, South Korea
E-mail: sungsoo@inha.ac.kr