

Suzuki–Miyaura Cross-Coupling and Ring-Closing Metathesis: A Strategic Combination for the Synthesis of Cyclophane Derivatives

Sambasivarao Kotha^{*[a]} and Kalyaneswar Mandal^[a]

Keywords: Cyclophanes / Isomerization / Olefin metathesis / Palladium / Ruthenium / Suzuki–Miyaura cross-coupling

The synthesis of cyclophane derivatives through a sequence involving Suzuki–Miyaura cross-coupling between α,α' -dibromo-*m*-xylene and arylboronic acid derivatives, alkenylation and ring-closing metathesis has been achieved. One of the cyclophanes was obtained by tandem isomerization and

metathesis. Significant magnetic anisotropic effects on the intra-annular hydrogen atoms were observed.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2006)

Introduction

Ruthenium-catalysed metathesis and palladium-catalysed cross-coupling reactions have recently made a profound impact on carbon–carbon bond formation in organic synthesis. In particular, with the advent of the commercially available and well defined metal carbene complexes^[1] (Figure 1) of Grubbs (e.g., **1** and **2**) and Schrock (e.g., **3**), metathesis has attracted renewed interest and has been quickly accepted into the mainstream of organic synthesis.^[2] These catalysts function under mild reaction conditions and exhibit a wide range of functional group tolerance, and these advances have opened up a completely new set of possibilities in organic synthesis. Of the various modes of olefin metathesis, the intramolecular version (i.e., ring-closing metathesis, RCM) has become more popular than other metathesis processes. RCM is a powerful tool for the construction of small (≥ 5 members), medium and large carbocyclic (or heterocyclic) ring systems starting from acyclic precursors.^[3]

Similarly, of the various Pd-catalysed cross-coupling reactions, the Suzuki–Miyaura (SM) cross-coupling reaction is one of the most efficient methods for the construction of C–C bonds.^[4] The preferred status of the SM cross-coupling reaction over other Pd-catalysed cross-coupling reactions is not coincidental: its key advantages are its mild reaction conditions and the commercial availability of diverse boronic acids, which are also environmentally safer than other organometallic reagents. In addition, the handling and removal of boron-containing by-products are easier than those associated with the use of other organometallic reagents, especially in large-scale syntheses. The combina-




Figure 1. Commonly used olefin metathesis catalysts.

tion of these two powerful tools should therefore open up new and short synthetic routes to various complex targets.


A cyclophane is defined as a molecule containing a bridged aromatic ring and usually contains a molecular cavity of varied size. Cyclophanes, subjects of investigations since the 1950s, are useful model systems for studying through-space interactions,^[6] have also found many applications in supramolecular chemistry, metal ion transport and catalysis, and represent intricate resources for a variety of host–guest interactions,^[7] as a result of which there is a continuous need to develop new and simple strategies for their synthesis. Several such strategies have been developed over the years, but methods for the synthesis of cyclophanes based on the usage of metathesis, Suzuki–Miyaura (SM) cross-coupling or combinations of these two reactions are limited.

Very few groups have successfully employed metathesis as a key step for the synthesis of cyclophane derivatives,^[8] application of SM cross-coupling for the synthesis of cyclophanes is less explored than that of metathesis,^[9] whilst reports involving a combination of metathesis and SM coupling are rare in organic synthesis. Guan and co-workers were the first to use the combination of palladium-catalysed SM cross-coupling and RCM for the efficient synthesis of

[a] Department of Chemistry, Indian Institute of Technology, Bombay
Powai, Mumbai 400076, India
Fax: +91-22-2572-3480
E-mail: srk@chem.iitb.ac.in

m-terphenyl-based cyclophane, with the aid of Grubbs' 2nd generation catalyst **2**.^[10]

Here we report a simple and straightforward synthetic strategy for cyclophane derivatives using the SM cross-coupling reaction and RCM as key steps. The strategy adopted in our study is shown in Scheme 1. The preliminary results of this research were published elsewhere.^[5]




Scheme 1. Synthetic strategy for cyclophane synthesis.

Results and Discussion

In general, the majority of SM cross-coupling reactions described so far have been associated with the coupling of aryl halides with arylboronic acids. The use of benzyl bromides as SM coupling partners, however, is less well explored.^[11]

To test the feasibility of the first step in our strategy, various functionalized boronic acids were coupled with the two-armed benzyl bromide **4** (Scheme 2). We found that **4** would react with various boronic acids in the presence of the palladium catalyst [Pd(PPh₃)₄] to generate the required cross-coupling products in good yields (Table 1).^[12]




Scheme 2. The reaction was conducted with the boronic acid (1.5 equiv. for each bromine atom in the substrate), Pd(PPh₃)₄ (6–10 mol-%).

Table 1. Suzuki cross-coupling reactions between **4** and various functionalized boronic acids.

Entry	Boronic acids	R	Products	Yield ^[a] (%)
i	5a	<i>p</i> -OMePh	6a	32
ii	5b	<i>p</i> -CHOPh	6b	80
iii	5c	<i>p</i> -CNPh	6c	84 ^[b]
iv	5d	<i>p</i> -AcPh	6d	71 ^[b]

[a] Isolated yields. [b] Purification was carried out by repeated recrystallization after column chromatography.

This methodology was also extended to the three-armed benzyl bromide derivative **7** (Scheme 3, Table 2). Since C₃-symmetric molecules are valuable core units for dendrimer design and for the synthesis of useful ligands, this methodology is likely to find applications in these areas.



Scheme 3. The reaction was conducted with boronic acid (1.5 equiv. for each bromine atom in the substrate), Pd(PPh₃)₄ (10–15 mol-%).



Table 2. Suzuki cross-coupling reactions between **7** and various functionalized boronic acids.

Entry	Boronic acids	R	Products	Yield ^[a] (%)
i	5a	<i>p</i> -OMePh	8a	61
ii	5b	<i>p</i> -CHOPh	8b	70
iii	5c	<i>p</i> -CNPh	8c	82

[a] Isolated yields.

In an attempted synthesis of the target cyclophane **11**, compound **6a** was allylated by treatment with allyl bromide in the presence of indium (Scheme 4)^[13] and the diol **9** was then subjected to RCM in the presence of Grubbs' 1st generation catalyst **1**, giving a complex mixture of products. However, use of the 2nd generation catalyst **2** under high-dilution conditions (0.001 M) at room temp. delivered the cyclophane derivative **10** as a mixture of diastereomers (1:2; determined by ¹H NMR integration) in 47% yield.^[14] The lower yield could be accounted for by the formation of various oligomeric side products during the macrocyclization reaction, as reported earlier.^[2] The successful formation of **10** was indicated by its ¹H NMR spectroscopic data, whilst the molecular ion peak at *m/z* = 393 [*M* + Na] in the mass spectrum further confirmed the structure of the desired cyclophane derivative. Subsequent PCC oxidation of the above mixture gave cyclophane **11**, the structure of which, with the assigned *trans* geometry, was confirmed by X-ray crystallographic data.^[5] The crystal structure of **11** reveals that in the solid state arrangement, the ethylene bridge is disordered with an averaged population of 70:30, perhaps due to the availability of void space. Further, this disorder is dynamic in nature, as the crystals did not show any disorder when cooled to 133 K.

To generalize the method, we turned our attention to the preparation of the higher analogues of **11** as depicted in Scheme 5. The RCM precursor **12** was prepared by Grignard treatment of **6b** with 4-bromobut-1-ene in 71% yield, and exposure of the diolefinic compound **12** to Grubbs' 2nd generation catalyst **2** in dichloromethane under high-dilution conditions (0.001 M) at room temp. resulted in the formation of the cyclophane derivative **13** (1:1.7; mixture of two diastereomers as determined from the ¹H NMR integration ratio) as the major product. The successful formation of **13** was indicated by its ¹H NMR spectroscopic data, and the structure of the desired cyclophane derivative **13** was further confirmed by the molecular ion peak at *m/z* = 421 [*M* + Na] in the mass spectrum. To our surprise, a

Scheme 4. Synthesis of **11** with RCM as a key step.Scheme 5. Synthesis of higher analogues of **11**.

minor product was isolated along with the target product as a mixture of two diastereomers (1:2.2; determined from ^1H NMR integration ratio) in significant yield. The structure of this minor product was assigned as **14** and its unsymmetrical nature is supported by its ^1H NMR spectrum. The molecular ion peak at $m/z = 407$ [$M + \text{Na}$] in the mass spectrum further confirmed the assigned structure of compound **14**.

The unusual formation of compound **14** could be explained in terms of a tandem isomerization induced by the Grubbs' catalyst, followed by a RCM sequence through the intermediacy of **17** (Scheme 5).^[15] Finally, oxidation of the mixture of diastereomers gave the cyclophanedione as a single compound. Treatment of diols **13** and **14** with PCC in dichloromethane at room temp. for 2 h thus gave the cy-

clophane derivatives **15** and **16**, respectively, each as a single isomer in modest yield. The formation of compounds **15** and **16** was confirmed by their IR and ^1H NMR spectroscopic data, and the exclusive *trans* geometry of the cyclophane derivative **16** was assigned through the observed coupling constant ($J = 15.0$ Hz) of the olefinic proton in its ^1H NMR spectrum. Finally, the structures of **15** and **16** were further confirmed by high-resolution mass spectral analysis.

Interestingly, significant magnetic anisotropic effects on the intra-annular aromatic hydrogens (H_i) were observed in all the cyclophane derivatives, as shown in Figure 2. The H_i protons in the open-chain analogues **9** and **12** absorb in the region of $\delta = 7.02$ ppm, but in the cyclic forms (i.e., in the cyclophane derivatives **10**, **11** and **13–16**) the intra-annular




Figure 2. Magnetic anisotropic effects in various cyclophane derivatives.

hydrogen atoms (H_i) located between the planes of the two *para*-substituted benzene rings are found to be shifted moderately upfield. As the cavity size (or aliphatic chain length) in such phane systems decreased, the upfield shifts of H_i increased, due to the increase in shielding effects (Figure 2).

Conclusions

In summary, we have demonstrated a new approach for the synthesis of cyclophane derivatives through the use of a combination of SM cross-coupling and RCM as key steps. One of the cyclophane derivatives was obtained through a tandem isomerization and metathesis reaction. Significant magnetic anisotropic effects on the intra-annular hydrogen atoms were observed, so these macrocyclic compounds could be potential targets for charge density studies in addition to host–guest complex experiments.

Experimental Section

General Remarks: All reactions were monitored by thin layer chromatography (TLC) carried out on glass plates coated with Acme's silica gel GF 254 (containing 13% calcium sulfate as a binder). Visualization of the spots on TLC plates was achieved by exposure either to iodine vapour or to UV light. Flash chromatography was performed with Acme's silica gel (100–200 mesh). Petroleum ether refers to the fraction of boiling point 60–80 °C. All the commercial grade reagents were used without further purification. 1st and 2nd generation Grubbs' catalysts (**1** and **2**), α,α' -dibromo-*m*-xylene, 4-bromobut-1-ene and all the boronic acids used here were purchased from Aldrich Chemical Co., Inc. (Milwaukee, WI USA). Magnesium, indium and allyl bromide (s. d. fine-Chem Ltd.) were used as received. $\text{Pd}(\text{PPh}_3)_4$ was prepared by the reported procedure.^[16] Infrared spectra were recorded on a Nicolet Impact 400 FT-IR spectrometer in $\text{KBr}/\text{CHCl}_3/\text{CCl}_4$. ^1H NMR (300, 400 MHz) and ^{13}C NMR (75.4, 100.6 MHz) spectra were determined at room temperature on a Varian VXR 300 or an AX 400 mercury plus in CDCl_3 solutions. Coupling constants (J values) are given in Hertz (Hz). Chemical shifts are expressed in parts per million (ppm) downfield from tetramethylsilane (TMS) as internal reference. High-resolution mass spectra were determined on a Micromass Q-ToF spectrometer.

General Procedure for Coupling between Arylboronic Acids and α,α' -Dibromo-*m*-xylene (4**):** A mixture of α,α' -dibromo-*m*-xylene (**4**, 1 equiv.), the arylboronic acid (3 equiv.), $\text{Pd}(\text{PPh}_3)_4$ (6–10 mol-%) and Na_2CO_3 (4 equiv.) in water and THF (1:1) was heated at 80 °C. The reaction mixture was degassed with argon for 20 min prior to the addition of the catalyst. At the conclusion of the reaction (TLC monitoring), the reaction mixture was diluted with water and extracted with ethyl acetate, and the combined organic layer was washed with water and brine and dried with anhydrous MgSO_4 . The solvent was evaporated and the crude product was loaded onto a silica gel column. Elution of the column with EtOAc/petroleum ether gave the desired cross-coupling product.

1,3-Bis(4-methoxybenzyl)benzene (6a**):** 4-Methoxyphenylboronic acid (50 mg, 0.33 mmol) and aqueous Na_2CO_3 (2 M solution, 4 equiv.) were added to a solution of **4** (26 mg, 0.1 mmol) in dimethoxyethane (3 mL), and the resulting reaction mixture was degassed with argon for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (12 mg, 0.01 mmol, 10 mol-%) was then added, the reaction mixture was heated at re-

flux for 12 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with petroleum ether gave compound **6a** (10 mg, 32%) as a white solid. R_f = 0.5 (silica gel, EtOAc/petroleum ether 1:39); m.p. 73 °C (from hexane). ^1H NMR (300 MHz, CDCl_3): δ = 3.78 (s, 6 H, $2 \times \text{OCH}_3$), 3.88 (s, 4 H, $2 \times \text{CH}_2$), 6.82 (d, J = 8.8 Hz, 4 H, ArH), 6.98 (d, J = 8.1 Hz, 2 H, ArH), 7.01 (s, 1 H, ArH), 7.08 (d, J = 8.8 Hz, 4 H, ArH), 7.18 (t, J = 7.3 Hz, 1 H, ArH) ppm. ^{13}C NMR (75.4 MHz, CDCl_3): δ = 41.0, 55.3, 113.9, 126.6, 128.6, 129.5, 129.9, 133.4, 141.7, 158.0 ppm. UV: $\lambda_{\text{max}}(\text{CHCl}_3)/\text{nm}$ 279 ($\epsilon/\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1}$ 3908). HRMS (EI): calcd. for $\text{C}_{22}\text{H}_{22}\text{O}_2$: 318.1619; found: 318.1612.

1,3-Bis(4-formylbenzyl)benzene (6b**):** 4-Formylphenylboronic acid (170 mg, 1.14 mmol) and aqueous Na_2CO_3 (2 M solution, 4 equiv.) were added to a solution of **4** (100 mg, 0.38 mmol) in THF, and the resulting reaction mixture was degassed with argon for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (36 mg, 0.03 mmol, 8 mol-%) was then added, the reaction mixture was heated at reflux for 12 h and worked up as described in the General Procedure, and the crude product was purified by silica gel column chromatography. Elution of the column with EtOAc/petroleum ether (10%) gave compound **6b** as a colourless oil (95 mg, 80%). R_f = 0.38 (silica gel, EtOAc/petroleum ether 1:4). ^1H NMR (300 MHz, CDCl_3): δ = 4.02 (s, 4 H, $2 \times \text{CH}_2$), 7.01 (s, 1 H, ArH), 7.04 (d, J = 7.5 Hz, 2 H, ArH), 7.25 (t, J = 7.5 Hz, 1 H, ArH) 7.33 (d, J = 8 Hz, 4 H, ArH), 7.8 (d, J = 8.2 Hz, 4 H, ArH), 9.97 (s, 2 H, $2 \times \text{CHO}$) ppm. ^{13}C NMR (100.6 MHz, CDCl_3): δ = 42.1, 127.3, 129.2, 129.7, 129.8, 130.1, 134.9, 140.4, 148.3, 192.0 ppm. IR (neat): $\tilde{\nu}$ = 1699 ($\text{C}=\text{O}$) cm^{-1} . UV: $\lambda_{\text{max}}(\text{CHCl}_3)/\text{nm}$ 265 ($\epsilon/\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1}$ 17327). HRMS (Q-ToF): m/z calcd. for $\text{C}_{22}\text{H}_{18}\text{O}_2\text{Na}$ [$M + \text{Na}$]: 337.1204; found: 337.1208.

1,3-Bis(4-cyanobenzyl)benzene (6c**):** 4-Cyanophenylboronic acid (125 mg, 0.85 mmol) and aqueous Na_2CO_3 (2 M solution, 4 equiv.) were added to a solution of **4** (77 mg, 0.29 mmol) in THF, and the resulting reaction mixture was degassed with argon for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (27 mg, 0.02 mmol, 8 mol-%) was then added, the reaction mixture was heated at reflux for 12 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with EtOAc/petroleum ether (15%) gave compound **6c** as a white solid (76 mg, 84%). R_f = 0.20 (silica gel, EtOAc/petroleum ether 1:4); m.p. 114–115 °C (from DCM/hexane 1:9). ^1H NMR (300 MHz, CDCl_3): δ = 4.0 (s, 4 H, $2 \times \text{CH}_2$), 6.97 (s, 1 H, ArH), 7.02 (d, J = 7.3 Hz, 2 H, ArH), 7.23–7.27 (m, 5 H, ArH), 7.57 (d, J = 8 Hz, 4 H, ArH) ppm. ^{13}C NMR (100.6 MHz, CDCl_3): δ = 41.9, 110.1, 119.0, 127.3, 129.2, 129.6, 132.4, 139.9, 146.5 ppm. UV: $\lambda_{\text{max}}(\text{CHCl}_3)/\text{nm}$ 284 ($\epsilon/\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1}$ 5621). HRMS (Q-ToF): m/z calcd. for $\text{C}_{22}\text{H}_{17}\text{N}_2$ [$M + \text{H}$]: 309.1392; found: 309.1398.

1,3-Bis(4-acetylbenzyl)benzene (6d**):** 4-Acetylphenylboronic acid (93 mg, 0.57 mmol) and aqueous Na_2CO_3 (2 M solution, 4 equiv.) were added to a solution of **4** (50 mg, 0.19 mmol) in THF, and the resulting reaction mixture was degassed for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (21.8 mg, 10 mol-%) was then added, the reaction mixture was heated at reflux for 12 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with EtOAc/petroleum ether (25%) gave compound **6d** (46 mg, 71%). R_f = 0.4 (silica gel, EtOAc/petroleum ether 3:7); m.p. 81 °C (from DCM/hexane 1:9). ^1H NMR (300 MHz, CDCl_3): δ = 2.56 (s, 6 H, $2 \times \text{CH}_3$), 3.99 (s, 4 H, $2 \times \text{CH}_2$), 7.01 (s, 1 H, ArH), 7.02 (d, J = 7.9 Hz, 2 H, ArH), 7.20–7.26 (m, 5 H, ArH), 7.87 (d, J = 8.5 Hz, 4 H, ArH) ppm. ^{13}C NMR (75.4 MHz, CDCl_3): δ = 26.6, 41.9, 127.1, 128.7, 129.0, 129.1, 129.6, 135.3, 140.5, 146.7 ppm. IR (neat): $\tilde{\nu}$ = 1681 cm^{-1}

(C=O). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 274 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 18273). HRMS (Q-ToF): m/z calcd. for $\text{C}_{24}\text{H}_{22}\text{O}_2\text{Na}$ [$M + \text{Na}$]: 365.1517; found: 365.1515.

1,3,5-Tris(4-methoxybenzyl)-2,4,6-trimethylbenzene (8a): 4-Methoxyphenylboronic acid (89 mg, 0.59 mmol) and aqueous Na_2CO_3 (1.5 mL, 2 M solution, 6 equiv.) were added to a solution of **7** (52 mg, 0.13 mmol) in THF, and the resulting reaction mixture was degassed for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (22 mg, 0.02 mmol, 15 mol-%) was then added, the reaction mixture was heated at reflux for 24 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with EtOAc/petroleum ether (20%) gave compound **8a** (38 mg, 61%). $R_f = 0.25$ (silica gel, EtOAc/petroleum ether 1:3); m.p. 148 °C (from chloroform). ^1H NMR (300 MHz, CDCl_3): $\delta = 2.15$ (s, 9 H, $3 \times \text{ArCH}_3$), 3.77 (s, 9 H, $3 \times \text{OCH}_3$), 4.07 (s, 6 H, $3 \times \text{ArCH}_2$), 6.79 (d, $J = 9.0$ Hz, 6 H, ArH), 6.95 (d, $J = 8.7$ Hz, 6 H, ArH) ppm. ^{13}C NMR (75.4 MHz, CDCl_3): $\delta = 16.8$, 35.4, 55.3, 113.9, 128.8, 132.4, 134.6, 135.2, 157.7 ppm. UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 279 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 6734). HRMS (Q-ToF): m/z calcd. for $\text{C}_{33}\text{H}_{36}\text{O}_3\text{K}$ ($M + \text{K}$): 519.2302; found: 519.2318.

1,3,5-Tris(4-formylbenzyl)-2,4,6-trimethylbenzene (8b): 4-Formylphenylboronic acid (155 mg, 1.03 mmol) and aqueous Na_2CO_3 (2 M solution, 6 equiv.) were added to a solution of **7** (104 mg, 0.26 mmol) in THF, and the resulting reaction mixture was degassed for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (45 mg, 0.04 mmol, 15 mol-%) was then added, the reaction mixture was heated at reflux for 24 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with EtOAc/petroleum ether (20%) gave compound **8b** (87 mg, 70%). $R_f = 0.25$ (silica gel, EtOAc/petroleum ether 3:7); m.p. 198 °C (from chloroform). ^1H NMR (300 MHz, CDCl_3): $\delta = 2.13$ (s, 9 H, $3 \times \text{CH}_3$), 4.24 (s, 6 H, $3 \times \text{CH}_2$), 7.20 (d, $J = 8.1$ Hz, 6 H, ArH), 7.79 (d, $J = 8.1$ Hz, 6 H, ArH), 9.97 (s, 3 H, $3 \times \text{CHO}$) ppm. ^{13}C NMR (100.6 MHz, CDCl_3): $\delta = 17.1$, 36.8, 128.6, 130.3, 134.5, 134.8, 135.3, 147.9, 192.1 ppm. IR (neat): $\tilde{\nu} = 1690\text{ cm}^{-1}$ (C=O). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 293 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 13354). HRMS (Q-ToF): m/z calcd. for $\text{C}_{33}\text{H}_{31}\text{O}_3$ [$M + \text{H}$]: 475.2273; found: 475.2296.

1,3,5-Tris(4-cyanobenzyl)-2,4,6-trimethylbenzene (8c): 4-Cyanophenylboronic acid (75 mg, 0.51 mmol) and aqueous Na_2CO_3 (2 mL, 2 M solution) were added to a solution of **7** (51 mg, 0.13 mmol) in THF (15 mL), and the resulting reaction mixture was degassed for 20 min. $\text{Pd}(\text{PPh}_3)_4$ (22 mg, 0.02 mmol, 15 mol-%) was then added, the reaction mixture was heated at reflux for 24 h and worked up as described in the General Procedure, and the crude product was purified by column chromatography. Elution of the column with EtOAc/petroleum ether (20%) gave compound **8c** (49 mg, 82%) $R_f = 0.19$ (silica gel, EtOAc/petroleum ether 3:7); m.p. 232 °C (from chloroform). ^1H NMR (300 MHz, CDCl_3): $\delta = 2.09$ (s, 9 H, $3 \times \text{CH}_3$), 4.2 (s, 6 H, $3 \times \text{CH}_2$), 7.12 (d, $J = 8.1$ Hz, 6 H, ArH), 7.56 (d, $J = 8.1$ Hz, 6 H, ArH) ppm. ^{13}C NMR (100.6 MHz, CDCl_3): $\delta = 17.0$, 36.6, 110.2, 119.0, 128.6, 132.5, 134.2, 135.4, 145.9 ppm. IR (neat): $\tilde{\nu} = 2227\text{ cm}^{-1}$ (C $\equiv$ N). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 252 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 13345). HRMS (Q-ToF): m/z calcd. for $\text{C}_{33}\text{H}_{28}\text{N}_3$ [$M + \text{H}$]: 466.2283; found: 466.2262.

1,3-Bis[4-(1-hydroxybut-3-enyl)benzyl]benzene (9): Allyl bromide (32 mg, 0.26 mmol) and indium metal (26 mg, 0.23 mmol) were added to a solution of **6b** (32 mg, 0.1 mmol) in DMF (2 mL), and the resulting reaction mixture was allowed to stir at room temp. After completion of the reaction (15 min, TLC monitoring) the reaction mixture was quenched with HCl (50%), diluted with water (10 mL) and extracted with ethyl acetate (3×10 mL). The com-

bined organic layer was washed with water and brine and dried with MgSO_4 , the solvent was evaporated, and the crude product was loaded onto a silica gel column. Elution of the column with EtOAc/petroleum ether (20%) gave the desired product **9** (34 mg, 85%). $R_f = 0.27$ (silica gel, EtOAc/petroleum ether 1:4); m.p. 60–61 °C (from hexane). ^1H NMR (300 MHz, CDCl_3): $\delta = 2.04$ (d, $J = 4.4$ Hz, 2 H, $2 \times \text{OH}$), 2.47–2.53 (m, 4 H, $2 \times \text{CH}_2\text{CH}=\text{CH}_2$), 3.93 (s, 4 H, $2 \times \text{ArCH}_2$), 4.71 (t, $J = 6.2$ Hz, 2 H, $2 \times \text{CHOH}$), 5.12–5.2 (m, 4 H, $2 \times \text{CH}_2\text{CH}=\text{CH}_2$), 5.74–5.88 (m, 2 H, $2 \times \text{CH}_2\text{CH}=\text{CH}_2$), 7.0–7.02 (m, 3 H, ArH), 7.13–7.22 (m, 5 H, ArH), 7.27 (d, $J = 8.1$ Hz, 4 H, ArH) ppm. ^{13}C NMR (75.4 MHz, CDCl_3): $\delta = 41.6$, 43.9, 73.2, 118.5, 126.0, 126.8, 128.6, 129.0, 129.6, 134.6, 140.5, 141.3, 141.7 ppm. IR (neat): $\tilde{\nu} = 3398\text{ cm}^{-1}$ (OH). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 265 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 1956). HRMS (EI): m/z calcd. for $\text{C}_{28}\text{H}_{30}\text{O}_2\text{Na}$ [$M + \text{Na}$]: 421.2143; found: 421.2139.

1,3-Bis[4-(1-hydroxypent-4-enyl)benzyl]benzene (12): A small crystal of iodine was added to a suspension of magnesium (19 mg, 0.79 mmol) under argon in dry ether (3 mL). While the contents were stirred, 4-bromobut-1-ene (0.09 mL, 0.89 mmol, dissolved in 5 mL of dry ether) was added slowly. The iodine colour disappeared completely and the reaction mixture was then stirred at room temp. for 1 h. Compound **12** (80 mg, 0.25 mmol) in diethyl ether (5 mL) was then added slowly, the ice-cold temperature being maintained. The reaction mixture was then allowed to stir at room temp. for 24 h, quenched with saturated ammonium chloride, diluted with water (10 mL) and extracted with ethyl acetate (3×10 mL). The combined organic layer was washed with water and brine and dried with MgSO_4 , the solvent was evaporated, and the crude product was loaded onto a silica gel column. Elution of the column with EtOAc/petroleum ether (20%) gave compound **12** as a colourless liquid (77 mg, 71%). $R_f = 0.36$ (silica gel, EtOAc/petroleum ether 3:7). ^1H NMR (300 MHz, CDCl_3): $\delta = 1.73$ –1.93 (m, 6 H, $2 \times \text{OH}$, $2 \times \text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 2.05–2.20 (m, 4 H, $2 \times \text{CH}_2\text{CH}=\text{CH}_2$), 3.93 (s, 4 H, $2 \times \text{ArCH}_2$), 4.67 (dd, $J = 7.8$, 5.6 Hz, 2 H, $2 \times \text{CHOH}$), 4.98 (d, $J = 10.4$ Hz, 2 H, $2 \times \text{CH}=\text{CHH}$), 5.04 (d, $J = 16.8$ Hz, 2 H, $2 \times \text{CH}=\text{CHH}$), 5.79–5.89 (m, 2 H, $2 \times \text{CH}=\text{CH}_2$), 7.01 (d, $J = 6.4$ Hz, 2 H, $2 \times \text{ArH}$), 7.02 (s, 1 H, ArH), 7.17 (d, $J = 8.4$ Hz, 4 H, $4 \times \text{ArH}$), 7.20 (t, $J = 8.4$ Hz, 1 H, ArH), 7.25 (d, $J = 7.2$ Hz, 4 H, $4 \times \text{ArH}$) ppm. ^{13}C NMR (100.6 MHz, CDCl_3): $\delta = 30.3$, 38.2, 41.7, 74.0, 115.1, 126.2, 126.9, 128.8, 129.2, 129.8, 138.4, 140.7, 141.4, 142.5 ppm. IR (neat): $\tilde{\nu} = 3402\text{ cm}^{-1}$ (OH). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 265 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 1975). HRMS (Q-ToF): m/z calcd. for $\text{C}_{30}\text{H}_{34}\text{O}_2\text{Na}$ [$M + \text{Na}$]: 449.2457; found: 449.2471.

RCM of 1,3-Bis[4-(1-hydroxybut-3-enyl)benzyl]benzene (9) in the Presence of 2nd Generation Grubbs' Catalyst 2: Grubbs' catalyst **2** (7 mg, 10 mol-%) was added to a solution of compound **9** (32 mg, 0.08 mmol) in dry degassed DCM (80 mL, 1 mmolar) and the solution was stirred for 18 h at room temp. The reaction mixture was then concentrated, and the crude product was purified on a silica gel column. Elution of the column with EtOAc/petroleum ether (20%) gave compound **10** as a white solid (14 mg, 47%) and as a mixture of diastereomers (1:2; determined from the ^1H NMR integration ratio of the protons attached to the chiral carbon atom). The spectroscopic data are for the major isomer: $R_f = 0.23$ (silica gel, EtOAc/petroleum ether 3:7); m.p. 222–223 °C (from methanol). ^1H NMR (300 MHz, CDCl_3): $\delta = 1.74$ (d, $J = 4$ Hz, 2 H, $2 \times \text{OH}$), 2.50–2.55 (m, 4 H, $2 \times \text{CH}_2\text{CH}=\text{CH}$), 3.89 (s, 4 H, $2 \times \text{ArCH}_2$), 4.67–4.71 (m, 2 H, $2 \times \text{ArCHOH}$), 5.10 (t, $J = 3.7$ Hz, 2 H, $2 \times \text{CH}_2\text{CH}=\text{CH}$), 6.33 (s, 1 H, ArH), 6.97 (d, $J = 8.4$ Hz, 4 H, ArH), 7.07 (d, $J = 8.1$ Hz, 4 H, ArH), 7.14 (d, $J = 8.4$ Hz, 2 H, ArH), 7.26 (t, $J = 7.5$ Hz, 1 H, ArH) ppm. IR (neat): $\tilde{\nu} = 3392\text{ cm}^{-1}$ (OH). UV: $\lambda_{\max}(\text{CHCl}_3)/\text{nm}$ 262 ($\epsilon/\text{dm}^3\text{mol}^{-1}\text{cm}^{-1}$ 1700). HRMS

(Q-ToF): m/z calcd. for $C_{26}H_{26}O_2Na$ [$M + Na$]: 393.1831; found: 393.1823.

RCM of 1,3-Bis[4-(1-hydroxypent-4-enyl)benzyl]benzene (12) in the Presence of 2nd Generation Grubbs' Catalyst 2: Grubbs' catalyst **2** (11 mg, 10 mol-%) was added to a solution of compound **12** (50 mg, 0.12 mmol) in dry degassed DCM (100 mL, 1.26 mmolar) and the solution was stirred for 24 h at room temp. The reaction mixture was then concentrated, and the crude product was purified on a silica gel column. Elution of the column with EtOAc/petroleum ether (40%) gave compound **13** as a white solid (22 mg, 47%) and as a mixture of diastereomers (1:1.7; determined from the 1H NMR integration ratio of the protons attached to the chiral carbon atom). The spectroscopic data are for the major diastereomer of **13**: R_f = 0.4 (silica gel, EtOAc/petroleum ether 1:1); m.p. 213–214 °C (from methanol). 1H NMR (300 MHz, $CDCl_3$): δ = 1.48–2.0 [m, 10 H, $2 \times CH(OH)CH_2CH_2C=$], 3.90 (s, 4 H, $2 \times ArCH_2$), 4.53–4.59 (m, 2 H, $2 \times CHOH$), 5.34–5.38 (m, 2 H, $2 \times HC=$), 6.69 (s, 1 H, ArH), 6.99–7.32 (m, 11 H, ArH) ppm. IR (neat): $\tilde{\nu}$ = 3401 cm^{-1} (OH). UV: $\lambda_{max}(CHCl_3)/nm$ 263 ($\epsilon/dm^3 mol^{-1} cm^{-1}$ 2588). HRMS (Q-ToF): m/z calcd. for $C_{28}H_{30}O_2Na$ [$M + Na$]: 421.2144; found: 421.2157.

Continued elution of the column with the same solvent system gave **14** as a white solid (10 mg, 21%) and as a mixture of diastereomers (1:2.2; determined from the 1H NMR integration ratio of the protons attached to the chiral carbon atom). The spectroscopic data are for the major isomer of **14**: R_f = 0.35 (silica gel, EtOAc/petroleum ether 1:1); m.p. 218–220 °C (from methanol). 1H NMR (400 MHz, $CDCl_3$): δ = 1.57–1.68 [m, 4 H, $CH(OH)CH_2CH_2C=$ and $2 \times OH$], 2.31–2.38 [m, 2 H, $CH(OH)CH_2CH_2C=$], 2.50–2.57 [m, 2 H, $CH(OH)CH_2C=$], 3.91 (s, 2 H, $ArCH_2$), 3.92 (s, 2 H, $ArCH_2$), 4.52–4.58 (m, 2 H, $2 \times CHOH$), 5.27–5.30 (m, 2 H, $CH=CH$), 6.77 (s, 1 H, ArH), 6.99 (d, J = 8 Hz, 4 H, ArH), 7.04–7.07 (m, 4 H, ArH), 7.17 (d, J = 8 Hz, 2 H, ArH), 7.29 (t, J = 7.6 Hz, 1 H, ArH) ppm. IR (neat): $\tilde{\nu}$ = 3402 cm^{-1} (OH). UV: $\lambda_{max}(CHCl_3)/nm$ 263 ($\epsilon/dm^3 mol^{-1} cm^{-1}$ 2678). HRMS (Q-ToF): m/z calcd. for $C_{27}H_{28}O_2Na$ [$M + Na$]: 407.1987; found: 407.1979.

1,5(1,4),3(1,3)-Tribenzenacycloundecaphan-8-ene-6,11-dione (11): Compound **10** (12 mg, 0.03 mmol) in anhydrous DCM (3 mL) was added in one portion to a vigorously stirred suspension of PCC (15 mg, 0.07 mmol) in anhydrous DCM (10 mL). Stirring was continued at room temp. for 2 h. Anhydrous ether (50 mL) was added and decanted off, and the black residue was washed with further diethyl ether (3×10 mL). The combined ether extracts were concentrated, and the crude product was purified by silica gel column chromatography. Elution of the column with EtOAc/petroleum ether (10%) gave compound **11** (9 mg, 75%) as white, needle-like crystals. R_f = 0.77 (silica gel, EtOAc/petroleum ether 3:7); m.p. 174 °C (from DCM/hexane 1:4). 1H NMR (300 MHz, $CDCl_3$): δ = 3.61 (dd, J = 3.6, 1.5 Hz, 4 H, $2 \times COCH_2$), 3.91 (s, 4 H, $2 \times ArCH_2$), 5.75 (tt, J = 3.6, 1.5 Hz, 2 H, $2 \times CH=$), 6.01 (s, 1 H, ArH), 6.99 (d, J = 8.4 Hz, 4 H, ArH), 7.14 (d, J = 7.7 Hz, 2 H, ArH), 7.26 (t, J = 7.5 Hz, 1 H, ArH); 7.7 (d, J = 8.4 Hz, 4 H, ArH) ppm. ^{13}C NMR (75.4 MHz, $CDCl_3$): δ = 41.2, 43.9, 126.4, 128.1, 128.3, 129.0, 129.6, 133.5, 140.8, 146.2, 197.1 ppm. IR (neat): $\tilde{\nu}$ = 1676 cm^{-1} (C=O). UV: $\lambda_{max}(CHCl_3)/nm$ 258 ($\epsilon/dm^3 mol^{-1} cm^{-1}$ 14479). HRMS (EI): calcd. for $C_{26}H_{22}O_2$: 366.1620; found: 366.1620.

1,5(1,4),3(1,3)-Tribenzenacyclotridecaphan-9-ene-6,13-dione (15): Compound **13** (9 mg, 0.02 mmol) in anhydrous DCM (3 mL) was added in one portion to a vigorously stirred suspension of PCC (13 mg, 0.06 mmol) in anhydrous DCM (3 mL). Stirring was continued at room temp. for 2 h. Anhydrous diethyl ether (25 mL) was

added and decanted off, and the black residue was washed with further diethyl ether (3×5 mL). The combined ether extracts were concentrated, and the crude product was purified by silica gel column chromatography. Elution of the column with EtOAc/petroleum ether (10%) gave compound **15** (7 mg, 79%) as white, needle-like crystals. R_f = 0.35 (silica gel, EtOAc/petroleum ether 1:4); m.p. 159–160 °C (from DCM/hexane 1:4). 1H NMR (300 MHz, $CDCl_3$): δ = 2.33–2.40 (m, 4 H, $2 \times COCH_2CH_2$), 2.85 (dd, J = 8.1, 7.1 Hz, 4 H, $2 \times COCH_2$), 3.97 (s, 4 H, $2 \times ArCH_2$), 5.55 (t, J = 4.8 Hz, 2 H, $2 \times CH_2CH=$), 6.52 (s, 1 H, ArH), 7.08 (d, J = 8.4 Hz, 4 H, ArH), 7.17 (d, J = 6.9 Hz, 2 H, ArH), 7.29 (t, J = 7.5 Hz, 1 H, ArH), 7.73 (d, J = 8.4 Hz, 4 H, ArH) ppm. IR (neat): $\tilde{\nu}$ = 1679 cm^{-1} (C=O). UV: $\lambda_{max}(CHCl_3)/nm$ 255 ($\epsilon/dm^3 mol^{-1} cm^{-1}$ 5549). HRMS (Q-ToF): m/z calcd. for $C_{28}H_{27}O_2$ [$M + H$]: 395.2011; found: 395.2019.

1,5(1,4),3(1,3)-Tribenzenacyclododecaphan-8-ene-6,12-dione (16): Compound **14** (8 mg, 0.02 mmol) in anhydrous DCM (3 mL) was added in one portion to a vigorously stirred suspension of PCC (13 mg, 0.06 mmol) in anhydrous DCM (2 mL). Stirring was continued at room temp. for 2 h, anhydrous diethyl ether (25 mL) was added and decanted off, and the black residue was washed with further diethyl ether (3×5 mL). The combined ether extracts were concentrated, and the crude product was purified by silica gel column chromatography. Elution of the column with EtOAc/petroleum ether (10%) gave compound **16** (6 mg, 76%) as white, needle-like crystals. R_f = 0.34 (silica gel, EtOAc/petroleum ether 1:4); m.p. 165 °C (from DCM/hexane 1:4). 1H NMR (300 MHz, $CDCl_3$): δ = 2.53–2.59 (m, 2 H, $COCH_2CH_2$), 2.91–2.95 (m, 2 H, $COCH_2CH_2$), 3.48 (dd, J = 7.1, 1.2 Hz, 2 H, $COCH_2C=$), 3.92 (s, 2 H, $ArCH_2$), 3.96 (s, 2 H, $ArCH_2$), 5.37 (dtt, J = 15.3, 6.6, 1.5 Hz, 1 H, $CH_2CH_2CH=$), 5.67 (dtt, J = 15.6, 6.0, 1.5 Hz, 1 H, $COCH_2CH=$), 6.27 (s, 1 H, ArH), 6.96 (d, J = 8.7 Hz, 2 H, ArH), 7.08 (d, J = 8.4 Hz, 2 H, ArH), 7.12–7.20 (m, 2 H, ArH), 7.25–7.30 (m, 1 H, ArH), 7.63 (d, J = 8.4 Hz, 2 H, ArH), 7.72 (d, J = 8.1 Hz, 2 H, ArH) ppm. IR (neat): $\tilde{\nu}$ = 1676 cm^{-1} (C=O). UV: $\lambda_{max}(CHCl_3)/nm$ 255 ($\epsilon/dm^3 mol^{-1} cm^{-1}$ 13982). HRMS (Q-ToF): m/z calcd. for $C_{27}H_{24}O_2Na$ [$M + Na$]: 403.1674; found: 403.1676.

Acknowledgments

We thank the DST for the financial support and the SAIF, Mumbai, for spectroscopic data. KM thanks the CSIR, New Delhi for the award of a research fellowship.

- [1] a) P. Schwab, M. B. France, J. W. Ziller, R. H. Grubbs, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2039–2041; b) M. Scholl, S. Ding, C. W. Lee, R. H. Grubbs, *Org. Lett.* **1999**, *1*, 953–956; c) R. R. Schrock, J. S. Murdzek, G. C. Bazan, J. Robbins, M. DiMare, M. O'Regan, *J. Am. Chem. Soc.* **1990**, *112*, 3875–3886.
- [2] a) *Handbook of Metathesis* (Ed.: R. H. Grubbs), Wiley-VCH, Weinheim, **2003**, vols. 1–3. For reviews on olefin metathesis, see: b) S. K. Armstrong, *J. Chem. Soc., Perkin Trans. 1* **1998**, 371–388; c) A. Fürstner, *Angew. Chem. Int. Ed.* **2000**, *39*, 3012–3043; d) S. Kotha, N. Sreenivasachary, *Indian J. Chem. B* **2001**, *40*, 763–780; e) R. R. Schrock, A. H. Hoveyda, *Angew. Chem. Int. Ed.* **2003**, *42*, 4592–4633; f) S. J. Connon, S. Blechert, *Angew. Chem. Int. Ed.* **2003**, *42*, 1900–1923; g) R. H. Grubbs, *Tetrahedron* **2004**, *60*, 7117–7140; h) M. D. McReynolds, J. M. Dougherty, P. R. Hanson, *Chem. Rev.* **2004**, *104*, 2239–2258; i) K. C. Nicolaou, P. G. Bulger, D. Sarlah, *Angew. Chem. Int. Ed.* **2005**, *44*, 4490–4527.
- [3] For some related examples in this area, see: a) D. Banti, M. North, *Tetrahedron Lett.* **2003**, *44*, 8157–8160; b) A. G. J. Com-

- meureuc, J. A. Murphy, M. L. Dewis, *Org. Lett.* **2003**, *5*, 2785–2788; c) S. Content, C. J. Dutton, L. Roberts, *Bioorg. Med. Chem. Lett.* **2003**, *13*, 321–325; d) P. Beaulieu, W. W. Ogilvie, *Tetrahedron Lett.* **2003**, *44*, 8883–8885; e) A. G. M. Barrett, A. J. Hennessy, R. Le Vezouet, P. A. Procopiou, P. W. Seale, S. Stefaniak, R. J. Upton, A. J. P. White, D. J. Williams, *J. Org. Chem.* **2004**, *69*, 1028–1037; f) V. T. Nguyen, N. De Kimpe, *Tetrahedron Lett.* **2004**, *45*, 3443–3446; g) S. Kotha, K. Mandal, *Tetrahedron Lett.* **2004**, *45*, 2585–2588; h) J. S. Clark, M. C. Kimber, J. Robertson, C. S. P. McErlean, C. Wilson, *Angew. Chem. Int. Ed.* **2005**, *44*, 6157–6162; i) G. C. H. Chiang, A. D. Bond, A. Ayscough, G. Pain, S. Ducki, A. B. Holmes, *Chem. Commun.* **2005**, 1860–1862; j) V. T. H. Nguyen, E. Bellur, P. Langer, *Tetrahedron Lett.* **2006**, *47*, 113–116; k) M. Ball, B. J. Bradshaw, R. Dumeunier, T. J. Gregson, S. MacCormick, H. Omori, E. J. Thomas, *Tetrahedron Lett.* **2006**, *47*, 2223–2227; l) T. S. Cooper, B. Atrash, P. Sheldrake, P. Workman, E. McDonald, *Tetrahedron Lett.* **2006**, *47*, 2241–2243.
- [4] a) A. Suzuki, H. C. Brown, *Organic Synthesis via Boranes*, Aldrich Chemical Company, Inc, Milwaukee, USA, **2003**, vol. 3. For reviews on SM cross-coupling, see: b) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457–2483; c) S. Kotha, K. Lahiri, D. Kashinath, *Tetrahedron* **2002**, *58*, 9633–9695; d) F. Bellina, A. Carpita, R. Rossi, *Synthesis* **2004**, 2419–2440; e) M. Moreno-Manas, R. Pleixats, R. M. Sebastian, A. Vallribera, A. Roglans, *J. Organomet. Chem.* **2004**, *689*, 3669–3684; f) R. M. Kellogg, B. V. Syncom, *Chemtracts* **2004**, *17*, 451–455; g) L. Bai, J.-X. Wang, *Curr. Org. Chem.* **2005**, *9*, 535–553; h) O. Baudoin, *Eur. J. Org. Chem.* **2005**, 4223–4229.
- [5] For a preliminary communication, see: S. Kotha, K. Mandal, K. K. Arora, V. R. Pedireddi, *Adv. Synth. Catal.* **2005**, *347*, 1215–1218.
- [6] a) E. Weber, *Cyclophanes*, Springer-Verlag, Berlin, Germany, **1994**, vol. 172; b) F. Vögtle, *Cyclophane Chemistry: Synthesis Structures, and Reactions*, John Wiley & Sons, Chichester, UK, **1993**; c) F. Diederich, *Cyclophanes*, Royal Society of Chemistry, Cambridge, UK, **1991**; d) H. Takemura (Ed.), *Cyclophane Chemistry for the 21st Century*, Research Signpost, Trivandrum, India, **2002**.
- [7] V. Prautzsch, S. Ibach, F. Vögtle, *J. Inclusion Phenom. Macrocyclic Chem.* **1999**, *33*, 427–458.
- [8] a) A. J. Locke, C. Jones, C. J. Richards, *J. Organomet. Chem.* **2001**, *637–639*, 669–676; b) A. B. Smith, C. M. Adams, S. A. Kozmin, *J. Am. Chem. Soc.* **2001**, *123*, 990–991; c) A. B. Smith, S. A. Kozmin, C. M. Adams, D. V. Paone, *J. Am. Chem. Soc.* **2001**, *123*, 990–991; d) A. Fürstner, F. Stelzer, A. Rumbo, H. Krause, *Chem. Eur. J.* **2002**, *8*, 1856–1871; e) M. E. Layton, C. A. Morales, M. D. Shair, *J. Am. Chem. Soc.* **2002**, *124*, 773–775; f) J. Tae, Y. Yang, *Org. Lett.* **2003**, *5*, 741–744; g) D. Song, G. Blond, A. Fürstner, *Tetrahedron* **2003**, *59*, 6899–6904; h) V. Martinez, J. Blais, D. Astruc, *Angew. Chem. Int. Ed.* **2003**, *42*, 4366–4369; i) V. Martinez, J.-C. Blais, G. Bravic, D. Astruc, *Organometallics* **2004**, *23*, 861–874; j) M. D. Watson, F. Jäckel, N. Severin, J. P. Rabe, K. Müllen, *J. Am. Chem. Soc.* **2004**, *126*, 1402–1407; k) G. Erker, G. Kehr, R. Fröhlich, *J. Organomet. Chem.* **2004**, *689*, 1402–1412; l) D. Branowska, A. Rykowski, *Tetrahedron* **2005**, *61*, 10713–10718; m) T. Ueda, N. Kanomata, H. Machida, *Org. Lett.* **2005**, *7*, 2365–2368; n) D. Branowska, I. Buczek, K. Kalin'ska, J. Nowaczyk, A. Rykowski, *Tetrahedron Lett.* **2005**, *46*, 8539–8541; o) S. K. Collins, Y. El-Azizi, *Pure Appl. Chem.* **2006**, *78*, 783–789; p) Y. El-Azizi, A. Schmitzer, S. K. Collins, *Angew. Chem. Int. Ed.* **2006**, *45*, 968–973.
- [9] a) A.-C. Carbonnelle, J. Zhu, *Org. Lett.* **2000**, *2*, 3477–3480; b) G. J. Bodwell, J. Li, *Angew. Chem. Int. Ed.* **2002**, *41*, 3261–3262; c) G. J. Bodwell, J. Li, *Org. Lett.* **2002**, *4*, 127–130; d) B. B. Smith, D. E. Hill, T. A. Cropp, R. D. Walsh, D. Cartrette, S. Hipps, A. M. Shachter, W. T. Pennington, W. R. Kwochka, *J. Org. Chem.* **2002**, *67*, 5333–5337; e) W. Wang, J. Xu, Y.-H. Lai, *Org. Lett.* **2003**, *5*, 2765–2768; f) A. J. Roche, B. Canturk, *Org. Biomol. Chem.* **2005**, *3*, 515–519; g) R. Lépine, J. Zhu, *Org. Lett.* **2005**, *7*, 2981–2984.
- [10] D. H. Camacho, E. V. Salo, Z. Guan, *Org. Lett.* **2004**, *6*, 865–868.
- [11] For recent examples, see: a) M. J. Sharp, V. Snieckus, *Tetrahedron Lett.* **1985**, *26*, 5997–6000; b) G. J. Pernia, J. D. Kilburn, J. W. Essex, R. J. Mortishire-Smith, M. Rowley, *J. Am. Chem. Soc.* **1996**, *118*, 10220–10227; c) S. Chowdhury, P. E. Georghiou, *Tetrahedron Lett.* **1999**, *40*, 7599–7603; d) H. Juteau, Y. Gareau, M. Labelle, C. F. Sturino, N. Sawyer, N. Tremblay, S. Lamontagne, M. C. Carrière, D. Denis, K. M. Metters, *Bioorg. Med. Chem.* **2001**, *9*, 1977–1984; e) L. Botella, C. Najera, *Angew. Chem. Int. Ed.* **2002**, *41*, 179–181; f) L. Chahen, H. Doucet, M. Santelli, *Synlett* **2003**, 1668–1672; g) S. Langle, M. Abarbri, A. Duchêne, *Tetrahedron Lett.* **2003**, *44*, 9255–9258; h) S. M. Nobre, S. I. Wolke, R. G. da Rosa, A. L. Monteiro, *Tetrahedron Lett.* **2004**, *45*, 6527–6530; i) C.-P. Chang, Y.-L. Huang, F.-E. Hong, *Tetrahedron* **2005**, *61*, 3835–3839; j) R. Singh, M. S. Viciu, N. Kramareva, O. Navarro, S. P. Nolan, *Org. Lett.* **2005**, *7*, 1829–1832.
- [12] In entries ii, iii and iv (Table 1), the required cross-coupling product was contaminated with ca. 10% of the self-coupling product derived from the corresponding boronic acid (observed from the ¹H NMR peak intensity ratio) because of their similar polarity. Hence, the purification of these materials was carried out by either repeated column chromatography or repeated crystallizations after column chromatography.
- [13] Yield varies from 74 to 85%.
- [14] A similar yield was observed when the reaction was performed in presence of 7 mol-% catalyst in 0.001 M dilution, but a longer time (28 h) was necessary for the completion of the reaction. However, under low dilution conditions (0.003 M), the yield of **15** decreased to 33%.^[5]
- [15] Related examples of Grubbs' catalyst-induced isomerization, see: a) T. R. Hoye, M. A. Promo, *Tetrahedron Lett.* **1999**, *40*, 1429–1432; b) M. K. Gurjar, P. Yakambram, *Tetrahedron Lett.* **2001**, *42*, 3633–3636; c) S. S. Kinderman, J. H. van Maarseveen, H. E. Schoemaker, H. Hiemstra, P. J. T. Rutjes, *Org. Lett.* **2001**, *3*, 2045–2048; d) D. C. Braddock, A. J. Wildsmith, *Tetrahedron Lett.* **2001**, *42*, 3239–3242; e) C. Cadot, P. I. Dalko, J. Cossy, *Tetrahedron Lett.* **2002**, *43*, 1839–1841; f) D. C. Braddock, A. Matsuno, *Tetrahedron Lett.* **2002**, *43*, 3305–3308; g) A. E. Sutton, A. Benjamin, A. Seigal, D. F. Finnegan, M. L. Snapper, *J. Am. Chem. Soc.* **2002**, *124*, 13390–13391; h) M. Arisawa, Y. Terada, M. Nakagawa, A. Nishida, *Angew. Chem. Int. Ed.* **2002**, *41*, 4732–4734; i) B. Alcaide, P. Almendros, *Chem. Eur. J.* **2003**, *9*, 1259–1262; j) W. A. L. van Otterlo, E. L. Ngidi, C. B. de Koning, *Tetrahedron Lett.* **2003**, *44*, 6483–6486; k) B. Schmidt, *Eur. J. Org. Chem.* **2003**, 816–819; l) B. Schmidt, *Synlett* **2004**, 1541–1544; m) S. Kotha, K. Mandal, *Tetrahedron Lett.* **2004**, *45*, 1391–1394.
- [16] D. R. Coulson, *Inorg. Synth.* **1972**, *13*, 121–124.

Received: June 26, 2006

Published Online: September 28, 2006