


SPOTLIGHTS ...

Hollow Spheres

H. Xu, W. Wei, C. Zhang, S. Ding,
X. Qu, J. Liu, Y. Lu, Z. Yang*

Low-Temperature Facile Template Synthesis of Crystalline Inorganic Composite Hollow Spheres

Chem. Asian J.
DOI: 10.1002/asia.200700017




Nothing in it! Composite hollow spheres of TiO_2 (shown), BaTiO_3 , and SrTiO_3 with varied crystalline phases and compositions can be synthesized under mild conditions. The sulfonated polystyrene gel templates used ensure that the crystalline hollow spheres grow without damage to their shells.

FRET Sensors

J. L. Vinkenborg, T. H. Evers,
S. W. A. Reulen, E. W. Meijer,
M. Merkx*

Enhanced Sensitivity of FRET-Based Protease Sensors by Redesign of the GFP Dimerization Interface

ChemBioChem
DOI: 10.1002/cbic.200700109




Close encounters. Sensor proteins based on fluorescence resonance energy transfer (FRET) often display a modest change in emission ratio upon activation. Here, we show that promoting intramolecular interactions between donor and acceptor fluorescent domains is an attractive new strategy for increasing the ratiometric change in FRET-based protease sensors.

Franck–Condon Analysis

R. Brause, M. Santa, M. Schmitt,*
K. Kleinermanns*

Determination of the Geometry Change of the Phenol Dimer upon Electronic Excitation

ChemPhysChem
DOI: 10.1002/cphc.200700127



Analysing phenol: The structural changes of phenol upon electronic excitation are quantified experimentally and theoretically. A Franck–Condon analysis of the fluorescence emission spectra confirms the local nature of the excitation.


Cheminformatics


A. Bender,* J. Scheiber, M. Glick,
J. W. Davies, K. Azzaoui, J. Hamon,
L. Urban, S. Whitebread, J. L. Jenkins

Analysis of Pharmacology Data and the Prediction of Adverse Drug Reactions and Off-Target Effects from Chemical Structure

ChemMedChem
DOI: 10.1002/cmdc.200700026

Preclinical Safety Pharmacology attempts to anticipate adverse drug reactions (ADRs) during early phases of drug discovery by testing compounds in simple, in vitro binding assays. In this paper we describe the successful application of cheminformatics methods to predict adverse side effects of drugs to accelerate drug discovery and decrease late stage attrition in drug discovery projects.





An empty low-lying $3\sigma^*$ orbital is responsible for the catalytic activity of oxidoiron(IV) compounds in alkane hydroxylation. We study the influence of the ligand environment on the spin state and catalytic activity of simple systems, correlating these properties to the strength of the electron donation from the ligands, and in particular, we prove that systems with *weak* equatorial and axial donor ligands show superior catalytic activity.

High-Valent Oxidoiron(IV) Systems

L. Bernasconi,* M. J. Louwerse,
E. J. Baerends*

The Role of Equatorial and Axial Ligands in Promoting the Activity of Non-Heme Oxidoiron(IV) Catalysts in Alkane Hydroxylation

Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.200601238



Unusual cleavage: A series of mono-nuclear or dinuclear Fe^{II} -phenylpyruvate enolate complexes that contain tetradentate tris(6-methyl-2-pyridylmethyl)amine (6-Me₃-TPA) or tridentate benzyl bis(2-quinolinylmethyl)amine (Bn-BQA) ligands are reported. The phenylpyruvate complexes react with dioxygen to undergo oxidative C2–C3 bond cleavage of phenylpyruvate (see scheme).


Iron Complexes

T. K. Paine, J. England, L. Que, Jr.*

Iron-Catalyzed C2–C3 Bond Cleavage of Phenylpyruvate with O₂: Insight into Aliphatic C–C Bond Cleaving Dioxygenases

Chem. Eur. J.
DOI: 10.1002/chem.200601844

Contrasting images: A new T_1 contrast agent for magnetic resonance imaging (MRI) based on MnO nanoparticles reveals a bright signal enhancement and fine anatomic structures in the T_1 -weighted MR image of a mouse brain (see picture; left MRI, right MnO-enhanced MRI (MONEMRI)). Furthermore, MnO nanoparticles conjugated with a tumor-specific antibody were used for selectively imaging breast cancer cells in a metastatic brain tumor.



Imaging Agents

H. B. Na, J. H. Lee,* K. An,
Y. I. Park, M. Park, I. S. Lee,
D.-H. Nam, S. T. Kim, S.-H. Kim,
S.-W. Kim, K.-H. Lim, K.-S. Kim,
S.-O. Kim, T. Hyeon*

Development of a T_1 Contrast Agent for Magnetic Resonance Imaging Using MnO Nanoparticles

Angew. Chem. Int. Ed.
DOI: 10.1002/anie.200604775



On these pages, we feature a selection of the excellent work that has recently been published in our sister journals. If you are reading these pages on a computer, click on any of

the items to read the full article. Otherwise please see the DOIs for easy online access through Wiley InterScience.