

# The Diene Effect: The Design, Development, and Mechanistic Investigation of Metal-Catalyzed Diene-yne, Diene-ene, and Diene-allene [2+2+1] Cycloaddition Reactions

Mitchell P. Croatt<sup>[a]</sup> and Paul A. Wender<sup>\*[a]</sup>

**Keywords:** Cycloaddition / Pauson–Khand reaction / Rhodium / Heterogeneous catalysis / Allenes

Metal-catalyzed diene-yne, diene-ene, and diene-allene [2+2+1] cycloaddition reactions provide new methods for the facile construction of highly functionalized five-membered rings. These reactions can be conducted with a variety of substrate substitution patterns and functional groups and often in the absence of solvent. The special reactivity of dienes, a key to enabling or enhancing the effectiveness of the [2+2+1] and other reactions, is significantly different from that of alkynes, alkenes, or allenes. For example, the [2+2+1] reaction of a diene-yne is accelerated compared to that of the corresponding ene-yne. An even more dramatic "diene

effect" is found with diene-enes and diene-allenes. While bis-enes and ene-allenes are not reported to yield [2+2+1] cycloadducts, the related diene-enes and diene-allenes undergo efficient [2+2+1] cycloadditions, providing new routes to cyclopentanones and alkylidenecyclopentanones. Mechanistic studies suggest that the unique reactivity observed with dienes arises from their participation in the putative rate-determining reductive elimination step by providing an additional energy-lowering coordination site for the transition metal catalyst.

## 1. Introduction


The design and discovery of new reactions are of unique importance to the advancement of synthesis. New reactions not only offer enhanced process efficiencies and selectivities, they also serve as the starting point for the introduction of new synthetic strategies and the resultant realization of

greater step economy.<sup>[1]</sup> All of these factors are of critical importance to green chemistry and the ideal synthesis.<sup>[1]</sup> To appreciate this point, one need only consider the first synthesis of cyclooctatetraene by Willstätter in 1911, which was ten steps,<sup>[2]</sup> and the subsequent one-step synthesis reported by Reppe in 1948<sup>[3]</sup> based on a then new reaction, the [2+2+2+2] cyclotetramerization of acetylene. In addition to step savings and efficiency and selectivity advantages, new reactions often provide greener<sup>[4]</sup> options for route, reagent,<sup>[5]</sup> and starting material selection and frequently allow for reduction of waste stream (steps add solvent from reac-

[a] Department of Chemistry, Stanford University, Stanford, CA 94305, USA  
Fax: +1-650-725-0259  
E-mail: wenderp@stanford.edu



Mitchell P. Croatt earned his B.S. degree in chemistry from the University of Minnesota, Twin Cities in 2002. While at the University of Minnesota, Mitchell took part in undergraduate research dealing with the synthesis of iminosugars under the guidance of Prof. George A. O'Doherty. In 2007, Mitchell obtained his Ph.D. in organic chemistry at Stanford University under the direction of Prof. Paul A. Wender. His research accomplishments at Stanford include the design and development of new transition-metal-catalyzed reactions and the use of new reactions in target- and function-oriented syntheses. Mitchell is continuing his education with postdoctoral studies in Switzerland at ETH-Zurich in the group of Prof. Erick M. Carreira.



Paul A. Wender is a product of Wilkes, Yale (Ph.D. with Ziegler), and Columbia (postdoctoral studies with Stork) Universities and is currently the Bergstrom Professor of Chemistry with an appointment by courtesy with the Chemical and Systems Biology Department of the Stanford Medical School. His group is interested in advancing the science of synthesis and in addressing problems in chemistry, biology, medicine, and materials science. Projects in Prof. Wender's group are diverse and include the design and development of new reactions, complex molecule total synthesis, rational design and synthesis of medicinally significant molecules, and the design and development of new methods for drug delivery.

tions and purifications),<sup>[6]</sup> development time, and cost, collectively contributing to greater productivity (i.e., human economy). Prompted by these considerations, our group has directed attention to the design and development of new or novel reactions for the synthesis of commonly encountered ring systems based on metal-catalyzed 2-, 3-, and 4-component processes (i.e.,  $[m+n]$ ,  $[m+n+o]$ ,  $[m+n+o+p]$  cycloadditions wherein  $m$ ,  $n$ ,  $o$ , and  $p$  are the number of atoms in each component joined in forming a new ring). A conceptual theme in many of these studies has been to use a catalyst to convert a commonly available starting functionality (e.g., alkene, alkyne, allene, or diene, the last a focus of this review) to a reactive intermediate that could then be trapped by one or more commonly available 1-, 2-, 3-, 4-, or 5-atom systems to produce a multicomponent product, a form of controlled oligomerization.<sup>[7]</sup> For example, our design of metal-catalyzed  $[5+2]$  reactions<sup>[8]</sup> was based on reports that vinylcyclopropanes react with transition metals to produce acyclic dienes and cyclopentenones.<sup>[9]</sup> It was hypothesized that this process proceeded through a reactive metallacyclohexene intermediate that might be trapped by a 2-atom  $\pi$ -component to produce a metallacyclooctene, which upon reductive elimination would give a seven-membered ring, overall representing a new  $[5+2]$  cycloaddition reaction, a homolog of the Diels–Alder  $[4+2]$  cycloaddition.<sup>[10]</sup> This process has now been shown to succeed intra- and intermolecularly.<sup>[8]</sup> It also inspired the design of an analogous  $[6+2]$  reaction of vinylcyclobutanones with  $\pi$ -systems,<sup>[11]</sup> related studies on the  $[6+1]$  reactions of allenylcyclobutanes and carbon monoxide (CO),<sup>[12]</sup> and the  $[3+2]$  reactions of cyclopropenones and alkynes.<sup>[13]</sup>

A significant characteristic of the above  $[m+n]$  reactions is that they convert relatively simple starting materials into products of increased molecular complexity and, therefore, often added value. Obviously, additional value could be realized if more components could be incorporated into the overall cycloaddition process. Like most cycloadditions, the previously described  $[m+n]$  processes allow for the formation of, at most, two new bonds from two components ( $m$  and  $n$ ) through capture of an organometallic intermediate. Additional bonds, and therefore value, could be created by “recycling” intermediates, i.e., using the organometallic intermediate involved in the coupling of the first two components to incorporate additional components, thereby forming subsequent organometallic intermediates and ultimately a multicomponent product. For example, the  $[5+2]$  cycloaddition has been used to incorporate one or two equivalents of CO leading to the first  $[5+2+1]$ <sup>[14]</sup> and  $[5+1+2+1]$ <sup>[15]</sup> cycloaddition reactions. A related series of studies in our group started with investigations on the first intramolecular  $[4+4]$  cycloadditions of bis-dienes<sup>[16]</sup> and led to the first metal-catalyzed intramolecular  $[4+2]$  cycloadditions,<sup>[17]</sup> which in turn prompted studies on the interception of organometallic intermediates in the latter reaction with CO.<sup>[18]</sup> This resulted in the first  $[4+2+1]$  and  $[2+2+1]$ <sup>[19]</sup> reactions of dienes,  $\pi$ -systems, and CO, the topic of this microreview. Our treatment is organized into sections dealing with the design of the reactions (Section 2), reactions of dienes, CO,

and alkynes (Section 3), alkenes (Section 4), and allenes (Section 5), mechanistic investigations (Section 6), and concluding remarks (Section 7).

## 2. Design of the $[4+2+1]$ and $[2+2+1]$ Reactions of Dienes Tethered to $\pi$ -Systems

Similar to our trapping of intermediates in the  $[5+2]$  reaction with CO to produce a  $[5+2+1]$  reaction, it was proposed that organometallic intermediates in the metal-catalyzed  $[4+2]$  reaction could incorporate a molecule of CO to produce either a  $[2+2+1]$  or  $[4+2+1]$  reaction (Figure 1), the former being an unexplored variant of the Pauson–Khand reaction<sup>[20]</sup> and the latter an entirely new process.<sup>[21]</sup> Experimentally, this notion received encouraging support when the treatment of diene-ynone **7** with a rhodium(I) catalyst under an atmosphere of CO yielded compounds resulting from the  $[4+2]$  cycloaddition and the new  $[2+2+1]$  and  $[4+2+1]$  reaction pathways (Scheme 1).<sup>[19a,22]</sup>





Figure 1. New reaction design: proposed trapping of intermediates in the  $[4+2]$  reaction with CO.



Scheme 1. Preliminary trapping studies: diversion of the  $[4+2]$  cycloaddition leads to new  $[2+2+1]$  and  $[4+2+1]$  reactions.


## 3. The Dienyl-Pauson–Khand Reaction

### 3.1 The Intramolecular Dienyl-Pauson–Khand Reaction<sup>[19a]</sup>

#### 3.1.1 Reaction Optimization

By variation of the reaction conditions that concomitantly produced the  $[4+2]$  product **8**,  $[2+2+1]$  reaction prod-


uct **9** (dienyl-Pauson–Khand reaction), and [4+2+1] reaction product **10** (Scheme 1), conditions were found that selectively and efficiently produced cyclopentenone **9** (Scheme 2). Important observations from the optimization study were that the [4+2] reaction product **8** was essentially eliminated by lowering the reaction temperature from 40 °C to ambient temperature and that the [2+2+1] reaction was not significantly affected by increasing or decreasing the pressure of CO.



Scheme 2. The optimized dienyl-Pauson–Khand reaction.

### 3.1.2 Substrate Scope


Using conditions similar to those of Scheme 2, diene-ynes with various substitutions were examined, and excellent yields of the dienyl-Pauson–Khand products resulted (Scheme 3). Many substrates with terminal alkynes (e.g., **13**) produced only modest yields of cycloadducts; however, substrates with silyl-substituted alkynes (e.g., **15**), which can be used as surrogates for terminal alkynes, reacted much more efficiently. Significantly, *spiro*-fused ring system **18**, incorporating a quaternary center, can be formed diastereoselectively in excellent yield.



Scheme 3. Representative dienyl-Pauson–Khand reactions.

### 3.1.3 Diene and Alkene Comparison

To explore the relative reactivities of dienes, alkenes, and styrenes as 2-carbon components in the [2+2+1] process, the reactions of ene-ynes **21** and **23** were compared with that of diene-yne **19** under similar conditions (Scheme 4). Significantly, ene-ynes **21** and **23** required higher catalyst loadings, increased temperatures, and extended reaction times when compared to the reaction of diene-yne **19**. These reactions illustrate that the additional unsaturation of a diene and its nature (ene vs. arene) significantly influence the rate of these cycloadditions.



Scheme 4. Comparison of the [2+2+1] reactions of a diene-yne with ene-ynes.


## 3.2 The Intermolecular Dienyl-Pauson–Khand Reaction<sup>[19b]</sup>

The Pauson–Khand reaction is a powerful process for assembling five-membered rings, but in most cases the two  $\pi$ -systems must be tethered. The intermolecular Pauson–Khand reaction has potential to yield products of significantly increased molecular complexity since it combines three separate, simple components. However, systems that work well intramolecularly with a five-atom tether often do not succeed with longer tethers or when conducted intermolecularly. To achieve an intermolecular Pauson–Khand reaction, some additional reactivity is required in the form of strain,<sup>[23]</sup> chelation assistance,<sup>[24]</sup> or both.<sup>[23b]</sup> In fact, Narasaka noted in the first report of a rhodium(I)-catalyzed intermolecular Pauson–Khand reaction, “the use of more reactive alkenes is essential” for the reaction with his best yield (69%) obtained with norbornene.<sup>[25]</sup> Based on the increase in reactivity observed with dienes relative to alkenes in the intramolecular dienyl-Pauson–Khand reaction, it was hypothesized that a diene could be used as a more reactive  $\pi$ -system in the intermolecular Pauson–Khand reaction.


### 3.2.1 Reaction Optimization

As anticipated from the previously discussed work, dienes provided the necessary reactivity enhancement for the intermolecular (three-component) dienyl-Pauson–Khand reaction, affording cycloadducts in excellent yield.

This reaction was found to work best with  $[\text{RhCl}(\text{CO})_2]_2$  relative to other  $\text{Rh}^{\text{I}}$  catalysts in TCE (Scheme 5: 1,1,2,2-tetrachloroethane) or DCE (1,2-dichloroethane). Significantly, the intermolecular dieny-Pauson–Khand reaction can be conducted in the absence of solvent, providing cyclopentenone **27** in good yield. While the role of solvent in modulating heat transfer on a large scale reactions must be considered, solventless reactions offer a green approach to some problems, particularly with respect to atom economy,<sup>[6]</sup> since they avoid atom loss from solvent, typically the largest atom loss in most reactions. As yet unoptimized in this reaction is the need for excess diene, due in part to its volatility and in part to its competing oligomerization under the reaction conditions. Finally, the intermolecular dieny-Pauson–Khand reaction was found to be scalable with only a modest decrease in efficiency (Scheme 6).



Scheme 5. Solvent study with the intermolecular dieny-Pauson–Khand reaction.




Scheme 6. Scalability of the intermolecular dieny-Pauson–Khand reaction.

### 3.2.2 Alkyne Scope

Different alkynes were examined using 2,3-dimethyl-1,3-butadiene as the diene component and previously optimized conditions (Scheme 7). In most cases the reactions were high yielding and tolerated different functional groups. The reaction with 1,4-dichloro-2-butyne is less efficient but provides a product, compound **35**, with a remarkable array of differentiated functionalities, a valuable building block for synthesis.


### 3.2.3 Diene Scope

The initial diene examined in the intermolecular dieny-Pauson–Khand reaction was 2,3-dimethyl-1,3-butadiene, a choice that was advantageous because reactions with this diene are higher yielding than reactions with most of the other dienes examined (Scheme 8). This diene is probably optimal under the current conditions due to the slow rate of oligomerization of both the initial diene and the result-



Scheme 7. Alkyne scope in the dieny-Pauson–Khand reaction.

ant olefin of the product. It was observed that other dienes used in this intermolecular dieny-Pauson–Khand reaction are more prone to these side reactions. An exception, 2,3-dibenzyl-1,3-butadiene, was comparable in reactivity to 2,3-dimethyl-1,3-butadiene. Significantly, when only 1.2 equiv. of this nonvolatile diene were used with alkyne **28**, an 82% yield of cyclopentenone **43** was obtained.




Scheme 8. Diene scope in the dieny-Pauson–Khand reaction.



### 3.2.4 Diene and Alkene Comparison

Because the intermolecular dieny-Pauson–Khand reaction was high yielding when dienes were used in the place of an alkene, the reactivity difference between a diene and an alkene was evaluated. For this comparison, norbornene was chosen as the alkene since it was previously found to be one of the best alkenes in the intermolecular Pauson–Khand reaction.<sup>[22]</sup> Under identical reaction conditions, 2,3-dimethyl-1,3-butadiene produced a 98% yield of cyclopentenone **27**, whereas norbornene generated less than 5% of cycloadduct **45** at full conversion (Scheme 9). Furthermore, the reaction with norbornene required twice as much time to consume the starting materials as compared to the reaction with a diene.




Scheme 9. Comparative reactivities of 2,3-dimethyl-1,3-butadiene and norbornene.

## 4. Diene-ene [2+2+1] Reaction<sup>[19c]</sup>


### 4.1 Introduction and Initial Results

A longstanding limitation of the [2+2+1] reaction of  $\pi$ -systems with CO is the necessity of using an alkyne as one of the two-carbon components. Previously, there were only two methods for the coupling of two alkenes and CO: a multiple-step procedure using stoichiometric zirconium<sup>[26]</sup> and a procedure involving cyclocarbonylation of an alkene tethered to an allene (Scheme 10).<sup>[27]</sup> In the latter case, Itoh and co-workers report that an ene-type cyclization reaction occurs with all other allene-enes (discussed later in Scheme 20) instead of a [2+2+1] cycloaddition reaction,<sup>[28]</sup> an observation that has also been reported by Brummond and co-workers.<sup>[29]</sup>

Given the accelerating effect of the diene on the Pauson–Khand reaction involving a diene-yne (Scheme 4 and Scheme 9), the first metal-catalyzed [2+2+1] reaction of a diene-ene and CO was attempted (Scheme 11). Gratifyingly, the diene-ene [2+2+1] reaction proved to be an efficient process. For dienes with smaller substituents at the 2-position (such as **53**), the  $\beta,\gamma$ -olefin of the [2+2+1] reaction product isomerized to the more thermodynamically stable  $\alpha,\beta$ -unsaturated position.<sup>[22c]</sup> This was found to occur under the reaction conditions and on purification.




Scheme 10. Prior reports of [2+2+1] reactions to form cyclopentanones.



Scheme 11. Initial conditions for the diene-ene [2+2+1] reaction.

### 4.2 Reaction Optimization


A screening of catalysts, solvents, and temperatures resulted in the identification of the optimal conditions for this new process:  $[\text{RhCl}(\text{CO})_2]_2$  (5 mol-%) as catalyst in DCE (0.1 M) at 80 °C. Significantly, it was found that the concentration of the reaction could be increased to the point at which no solvent is required. In the case of diene-enes **51** and **53**, solventless reactions resulted in 83% and 89% yields of product, respectively (including isomerized enone **55**; Scheme 12). These two reactions were designed to be “green”, especially in terms of atom-economy,<sup>[23]</sup> because no solvent atoms were wasted, no excess reagent was used, and the only lost atoms were from the rhodium catalyst, which was used substoichiometrically. It is important to note that in this and other cases the minimal catalyst loading could not be determined, as it would require the use of larger scale reactions than that acceptable in an academic setting. Loadings between 1–10 mol-% allowed for accurate weighing without using large quantities of starting materials.



Scheme 12. Solvent-free diene-ene [2+2+1] reactions.


### 4.3 Substrate Scope

With the optimized conditions for the diene-ene [2+2+1] reaction determined, variations in substitutions, tethers, and tether lengths were examined (Scheme 13). The reaction can be used to form heterocycles by incorporating an ether or sulfonamide group in the tether (**56** and **58**), quaternary centers by having additional substitution on the alkene (**60**), and [4.3.0] bicyclic products by having an additional methylene unit in the tether (**62**). The excellent yields for these reactions are significant because the formation of heterocycles, quaternary centers, and [4.3.0] bicycles has historically been difficult to accomplish with many transition-metal-catalyzed reactions, especially Pauson–Khand-type reactions.



Scheme 13. Representative diene-ene [2+2+1] reactions.


Based on the success of the reaction of diene-ene **62** bearing a 4-atom tether, an intermolecular reaction was attempted using 2,3-dimethyl-1,3-butadiene, norbornene, and CO. Significantly, [2+2+1] reaction product **64** was formed in 73% yield with only the *exo*-product observed (Scheme 14).<sup>[30]</sup> This intermolecular reaction converts simple starting materials that are all commercially available into a product with a large increase in molecular complexity and therefore potential value.



Scheme 14. Intermolecular diene-ene [2+2+1] reaction.

### 4.4 Diene and Alkene Comparison

To determine if the diene is necessary for a [2+2+1] reaction to form a cyclopentanone, control reactions with bis-enes were examined (Scheme 15). The bis-enes were similar to a diene-ene with respect to the electronics or coordinating ability of the diene. Of mechanistic significance, none of these reactions produced [2+2+1] cycloadducts; only slow decomposition of starting materials was observed. For the diene-ene [2+2+1] reaction, in contrast to the dienyl-Pauson–Khand reaction, the diene does more than accelerate the reaction; the diene enables the [2+2+1] reaction to occur.



Scheme 15. Control reactions for the diene-ene [2+2+1] reaction.

### 4.5 Synthetic Utility

While examining the functionality produced by the diene-ene [2+2+1] reaction, it was noted that the core system of all medicinally used prostacyclin analogues contain a similar bicyclic structure (Figure 2). It was envisioned that reduction of the carbonyl group of a [2+2+1] product from the convex face would produce the stereocenters of such core structures. Additionally, the side chain could be attached via a cross-metathesis reaction.<sup>[31]</sup> Since remodulin (**71**) is one of the most successful prostacyclin analogues, and would pose a significant test for the diene-ene [2+2+1] reaction due to the bicyclo[4.3.0]nonane core, synthesis of its tricyclic core was chosen for study. The synthetic route used for large-scale production of remodulin,<sup>[32]</sup> reported in 2002, is 19 total steps (15 longest linear) and features a diastereoselective Pauson–Khand reaction using stoichiometric quantities of a cobalt catalyst.<sup>[33]</sup>






Figure 2. Structure of prostacyclin, carbacyclin, and medically examined prostacyclin analogues.

The diene-ene required for synthesizing remodulin using a [2+2+1] reaction was assembled in only six steps from 3'-hydroxy acetophenone (Scheme 16). In this case, the cationic catalyst previously examined with the dienyl-Pauson–Khand reaction in Section 3 (Scheme 17) proved superior for the construction of the tricyclic core of remodulin.




Scheme 16. Synthesis of diene-ene for remodulin. (a)  $\text{K}_2\text{CO}_3$ , acetone, NaI, allyl bromide, 95%; (b) 220 °C (microwave), 75% (2:1 mixture of desired to undesired regioisomer); (c) LiHMDS, acrolein, 89%; (d) TsOH, 95%; (e)  $\text{NaBH}_4$ ,  $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ , 89%; (f) TBSCl, imidazole, 64%.

With conditions determined for the diene-ene [2+2+1] reaction, a serial [2+2+1]/diastereoselective reduction was attempted (Scheme 18). Results showed that it was possible to form the tricyclic core of remodulin and install four of the five stereogenic centers in the molecule in a single operation. Importantly, this step converts a simple monocyclic starting material into an intermediate that can be used in the synthesis of remodulin or a variety of analogues, because the two sites required for diversification are readily functionalized from the completed core structure (mono-



Scheme 17. Screening of conditions for the [2+2+1] reaction.

substituted olefin for cross-metathesis, and phenol for alkylation). More importantly, this study illustrates the value of the diene-ene [2+2+1] reaction as a method for the step-economical synthesis of complex targets.




Scheme 18. Construction of tricyclic core of remodulin.


## 5. The Diene-allene [2+2+1] Cycloaddition Reaction<sup>[19d]</sup>

### 5.1 Introduction and Initial Results

There are many reports of allene-yne [2+2+1] reactions (Scheme 19)<sup>[34]</sup> and their usefulness has been illustrated in total syntheses of several natural products.<sup>[20a]</sup> Notably, when a rhodium(I) catalyst is used, the allene reacts selectively with the distal olefin.<sup>[35]</sup> This behavior is clearly process-dependent, based on earlier reports from our group in which diene-allenes reacted in a [4+2] reaction with either the distal or proximal olefin of the allene depending on whether a  $\text{Rh}^{\text{I}}$  or  $\text{Ni}^0$  catalyst is used.<sup>[17c]</sup> Despite the amount of research on the allenic Pauson–Khand reaction, there have not been any reports of an alkene reacting with a simple allene in a [2+2+1] fashion.<sup>[36]</sup> In fact, it has been reported independently by Brummond<sup>[27]</sup> and Itoh<sup>[26]</sup> that allenes tethered to alkenes preferentially and exclusively undergo an ene-type cyclization reaction, even under an atmosphere of CO (Scheme 20). Only in a single case was a [2+2+1] reaction of this type observed (discussed earlier in Scheme 10). Since a unique reactivity was observed for dienes with the diene-yne and diene-ene [2+2+1] reactions, a study of the diene-allene [2+2+1] reaction was undertaken.




Scheme 19. Reactions of allenes tethered to alkynes in the presence of CO.



Scheme 20. Reactions of allenes tethered to alkenes in the presence of CO.

Under the conditions optimal for the diene-ene [2+2+1] reaction, diene-allene **89** reacted with the proximal olefin of the allene in a [2+2+1] fashion (Scheme 21). Importantly, this reactivity is different from that of an allene-yne (Scheme 19) or allene-ene (Scheme 20). Diene-allene **89** also underwent a [4+2] reaction with the initial conditions; however, this type of reactivity was averted by lowering the reaction temperature from 80 °C to ambient temperature. In this case, a 99% yield of alkylidene cyclopentane **90** was produced.



Scheme 21. Diene-allene [2+2+1] reaction.

## 5.2 Reaction Optimization and Co-Catalysis of the Reaction

Examination of solvents for this reaction led to the realization of a large rate increase for reactions run in TFE (2,2,2-trifluoroethanol) relative to those run in DCE (Table 1). This acceleration appears to be a case of acid co-catalysis since the addition of various acids to DCE also increased the reaction rate (compare entry 1 with entries 5–10). Significantly, it was found that the rate enhancement is not linear with the  $\text{p}K_{\text{a}}$  of the acid used since the reaction is faster in TFE (aqueous  $\text{p}K_{\text{a}}$  of 12.5, entry 2) than in acetic acid (aqueous  $\text{p}K_{\text{a}}$  of 4.76, entry 8) or methanol (aqueous  $\text{p}K_{\text{a}}$  of 15.5, entry 3). Additionally, it was found that phenol (aqueous  $\text{p}K_{\text{a}}$  of 10), whose acidity is similar to TFE, had an analogous rate enhancement when used in the

reaction as an additive (entries 9 and 10). Although this acceleration appears to be a result of the  $\text{p}K_{\text{a}}$  of the additive needing to be in an appropriate range, it could also be due to the hydrogen-bond-donating ability of the acids.<sup>[37]</sup> It was previously reported that hydrogen-bond donation does not correlate directly with acidity<sup>[38]</sup> and that TFE is a better hydrogen-bond donor than methanol or acetic acid.<sup>[37]</sup> Based on these results, it is hypothesized that the acid co-catalyst hydrogen-bonds with a CO ligand or rhodium–acyl species and therefore increases the rate of the reaction by lowering the barrier for CO insertion or reductive elimination. Importantly, this study illustrates the significance of screening different acids with transition-metal-catalyzed cycloaddition reactions, something rarely evaluated. Benefiting from the acceleration, the diene-allene [2+2+1] reaction in TFE using only 0.1 mol-% catalyst was complete in eight hours at ambient temperature (97% yield of **90**, entry 11).

Table 1. Acidic acceleration of the diene-allene [2+2+1] reaction.

Entry	Solvent	Additive (equiv.)	Time [h]	Yield of <b>90</b> <sup>[a]</sup>
1	DCE	–	7	99%
2	TFE	–	0.25	92%
3	MeOH	–	12	96%
4	CH <sub>3</sub> CN	–	45	77%
5	DCE	TsOH (0.05)	1.3	81%
6	DCE	AcOH (6)	4	93%
7	DCE	AcOH (100)	2.5	96%
8	AcOH	AcOH (neat)	2	93%
9	DCE	PhOH (14)	0.25	94%
10	DCE	TFE (14)	2	95%
11 <sup>[b]</sup>	TFE	–	8	97%


[a] Isolated yield; [b] 0.1 mol-% catalyst, 0.5 M.

## 5.3 Substrate Scope

Under the optimal reaction conditions {1–2.5 mol-% [RhCl(CO)<sub>2</sub>]<sub>2</sub> in TFE or DCE at ambient temperature with 1 atm of CO}, different diene-allenes were examined. Similar to the other [2+2+1] reactions, the diene-allene [2+2+1] reaction tolerates a wide range of substitutions and tethers without significant decreases in efficiency (Scheme 22). It was observed that reactions were much higher yielding with substitution at the terminus of the allene, although the reaction of monosubstituted allene **94** was successful when slightly modified conditions were used. While the reaction of a substrate with methyl substitution at the terminus of



the allene (**96**) was minimally selective, the reaction of a substrate with a larger *tert*-butyl group (**98**) produced a single diastereomer in 83 % yield.



Scheme 22. Representative diene-allene [2+2+1] reactions.

Importantly, dienes tethered to allenes allow for access to a different mode of reactivity when compared with reactions of alkynes or alkenes tethered to allenes (Scheme 19 and Scheme 20, respectively). Additionally, the reaction conditions are very mild (generally ambient temperature with 1 atm of CO) and produce products with a high level of molecular complexity.

## 6. Mechanistic Investigation of the [2+2+1] Reactions of Dienes Tethered to $\pi$ -Systems

### 6.1 Introduction

Due to the reactivity enhancement observed with dienes in the [2+2+1] reactions,<sup>[39]</sup> a study was initiated to address the source of this effect and understand why the [4+2+1] reaction is generally not observed.<sup>[40]</sup> The proposed mechanism for the metal-catalyzed [2+2+1] reaction of a diene tethered to a  $\pi$ -system involves five main steps: (1) catalyst coordination,<sup>[41]</sup> (2) oxidative cyclization, (3) CO insertion, (4) reductive elimination, and (5) CO coordination (Figure 3). Each of these steps can be connected to additional steps (e.g. the conversion of **A** to **C**); however, for the sake

of clarity, these five main steps will be discussed. Since changes in CO pressure do not have a significant effect on the rate of the reaction,<sup>[19]</sup> it is likely that CO coordination is not the rate-determining step (RDS). Therefore, it was decided to determine which of the other four steps is the RDS.

### 6.2 *cis*-Diene Results

To obtain information about the energy barriers for the steps described earlier, *cis*-dienes tethered to the respective  $\pi$ -systems were initially studied (Figure 4). If the energy barrier for ring expansion (**I** to **C**) is larger than the barrier for CO insertion and subsequent reductive elimination, *cis*-diene **100** should produce cycloadduct **101** wherein the vinyl substituent is epimeric to previously synthesized cycloadduct **3**. Alternatively, if ring expansion to **C**<sup>[42]</sup> is facile and reversible, the geometry of the diene will rapidly be scrambled and the same product will be formed irrespective of the initial stereochemistry of the diene. It should be noted that cycloadducts corresponding to **3** are thermodynamically favored, but there is no interconversion between the two epimeric [2+2+1] reaction products since formation of the  $\alpha,\beta$ -unsaturated system is a thermodynamic well.

When *cis*-diene-yne **102** and *cis*-diene-ene **103** were used in the [2+2+1] reaction, cycloadducts identical to those obtained from the corresponding *trans*-diene starting materials were generated and at approximately the same rates (Scheme 23). Importantly, the *trans*-diene starting materials were observed in both cases; however, a *cis*-diene without a tethered  $\pi$ -system (**104**) did not isomerize under the reaction conditions (Scheme 24). These results support a mechanism in which catalyst coordination and oxidative cyclization are rapid, reversible, and not the RDS. Thus, the remaining options for the RDS are CO insertion or reductive elimination.

### 6.3 Stoichiometric Use of Catalyst

In order to obtain more mechanistic information, a stoichiometric amount of catalyst was used in an attempt to observe intermediates in the diene-yne and diene-ene [2+2+1] reactions. Although the observation of intermediates does not imply that they are necessarily on the reaction pathway,<sup>[43]</sup> and often their presence indicates that they are not along the pathway,<sup>[44]</sup> it was envisioned that the observation of a metallacycle without CO inserted would substantiate the hypothesis that CO insertion or reductive elimination is the RDS. Alternatively, observation of an intermediate with a rhodium-acyl group would provide strong evidence against CO insertion being the RDS. For this reason, if a rhodium-acyl species is observed, it is likely that reductive elimination is the RDS.

Initially, diene-yne **11** was examined using the standard catalyst,  $[\text{RhCl}(\text{CO})_2]_2$ , in a stoichiometric quantity (Scheme 25).<sup>[45]</sup> Based on the <sup>1</sup>H and 2D NMR spectroscopic experiments, two main intermediates were observed




Figure 3. Proposed mechanistic overview of the [2+2+1], [4+2+1], and [4+2] reactions.





Figure 4. Reaction pathways for *cis*-dienes and  $\pi$ -systems.

(Figure 5). These intermediates were converted into [2+2+1] reaction product **12** over time or upon heating. Unfortunately, the characterization of these intermediates proved difficult due to the low signal to noise ratio for  $^{13}\text{C}$  NMR spectroscopic experiments.




Scheme 23. Results of a *cis*-diene-yne and a *cis*-diene-ene.

Scheme 24. Reactivity of *cis*-diene without a tethered  $\pi$ -system.





To facilitate the characterization of the intermediates, a reaction was run with  $^{13}\text{C}$ -labeled CO on the catalyst



Scheme 25. Diene-yne [2+2+1] reaction with stoichiometric catalyst loading.

Figure 5. <sup>1</sup>H spectra using stoichiometric catalyst loading: (a) starting diene-yne; (b) intermediates formed; (c) further reaction forms product.

(Scheme 26). Significantly, this led to the determination that both of the intermediates in the diene-yne [2+2+1] reaction had inserted CO as evidenced by the shift and coupling in the <sup>13</sup>C NMR spectra (Figure 6).<sup>[46]</sup> As discussed earlier,

Scheme 26. Diene-yne [2+2+1] reaction with <sup>13</sup>C-labeled catalyst.Figure 6. <sup>13</sup>C NMR spectrum for a diene-yne reaction.Scheme 27. Diene-ene [2+2+1] reaction with <sup>13</sup>C-labeled catalyst.Figure 7. <sup>13</sup>C NMR spectrum for a diene-ene reaction.

this does not mean that these exact intermediates lie on the reaction pathway, but the evidence suggests that CO insertion and the steps prior to it are not the RDS.


The reaction of diene-ene **53** with a stoichiometric amount of the  $^{13}\text{C}$ -labeled catalyst was subsequently examined (Scheme 27). Similar to the diene-yne reaction, the  $^{13}\text{C}$  NMR spectrum showed two rhodium-acyl intermediates (Figure 7). Therefore, since the rate of forming the CO-inserted intermediates is faster than the rate of product formation, the evidence again suggests that CO insertion is not the RDS.

#### 6.4 Conclusions for the Mechanism of the [2+2+1] Reactions of Dienes Tethered to $\pi$ -Systems

Based on the empirical data, it was previously determined that CO coordination is not the RDS. From the reactions run with *cis*-dienes tethered to  $\pi$ -systems, it was found that catalyst coordination and oxidative cyclization are rapid and reversible for the diene-yne and diene-ene reactions. Finally, from the reactions using a stoichiometric amount of catalyst with  $^{13}\text{C}$ -labeled CO, it was determined that the rate of CO insertion for the two intermediates in the diene-yne and diene-ene reactions was faster than the rate of product formation. Therefore, under these conditions, the evidence suggests that *reductive elimination is the rate-determining step for the diene-yne and diene-ene [2+2+1] reactions*.

Since reductive elimination appears to be the RDS, two hypotheses can be made as to why a [2+2+1] reaction is favored over a [4+2+1] reaction. First, reductive elimination with an exocyclic vinyl group would provide the catalyst with bidentate coordination to stabilize the transition state as compared to single ligation for the [4+2+1] reaction (Scheme 28). A similar result has been observed with the [5+2] reaction whereby a “migratory reductive elimination” was found to be much lower in energy when the rhodium could bind to another olefin within the molecule.<sup>[47]</sup> Another possibility is that the reductive elimination occurs with 1,3-transposition of the olefin. In other words, intermediate **110** directly forms **111** (Scheme 29) in a similar mechanism as an allylation of a ketone with an allyl-metal species. Either of these hypotheses can be used to explain why the diene is important since the olefin is intricately in-

volved in both of the mechanisms. These hypotheses are also consistent with the observed absence of reactivity when the diene of the starting material is replaced with an isolated alkene or styrene.



Scheme 29. Allylic transposition and reductive elimination.


## 7. Conclusions

The mechanism of the [4+2] cycloaddition of dienes and  $\pi$ -systems has been used as a conceptual starting point for the design of an unexplored variant of the Pauson–Khand [2+2+1] reaction involving dienes and of a new [4+2+1] reaction. The intramolecular metal-catalyzed [4+2] reaction introduced in our earlier work<sup>[17]</sup> is a very efficient process, often serving as an alternative or preferred option to a Diels–Alder process. Significantly, capture of intermediates along the [4+2] reaction pathway with CO has been possible using commercially available, bench-stable rhodium(I) catalysts, resulting in diene-yne, diene-ene, and diene-allene [2+2+1] reactions. It is also noteworthy that capture can be conducted with other  $\pi$ -systems.<sup>[48]</sup> The [2+2+1] reactions discussed herein convert relatively simple starting materials into mono- or bicyclic molecules with significantly increased molecular complexity including three new bonds, up to four new stereocenters, and functional groups including ketones and olefins that allow for facile subsequent functionalization. Mechanistic studies suggest that with the diene-yne and diene-ene [2+2+1] reactions the rate-determining step is reductive elimination. It is therefore hypothesized that the additional olefin of the diene is beneficial for these reactions because it provides for bidentate binding to the catalyst during the reductive elimination. Thus, as has even been observed in other systems with different catalysts,<sup>[39]</sup> the diene exhibits unique reactivity compared to an isolated alkene even though one of the  $\pi$ -systems of the diene is unchanged by the reaction. This observation (diene effect) should have widespread use in other cycloaddition processes, especially those for which simple monoalkenes are poor substrates.

## Acknowledgments

This research was supported by a grant from the National Science Foundation (CHE-0450638). Fellowship support from Eli Lilly and Co. (M. P. C.) is gratefully acknowledged.

- [1] a) P. A. Wender, B. L. Miller, *Nature* **2009**, *460*, 197–201; b) P. A. Wender, M. P. Croatt, B. Witulski, *Tetrahedron* **2006**, *62*, 7505–7511; c) P. A. Wender, F. C. Bi, G. G. Gamber, F. Gosselin, R. D. Hubbard, M. J. C. Scanio, R. Sun, T. J. Wil-



Scheme 28. Assisted reductive elimination.



- liams, L. Zhang, *Pure Appl. Chem.* **2002**, *74*, 25–31; d) P. A. Wender, S. T. Handy, D. L. Wright, *Chem. Ind. (London)* **1997**, 766–769; e) P. A. Wender, B. L. Miller, *Organic Synthesis: Theory and Applications*, Vol. 2 (Ed.: T. Hudlicky), JAI Press, Greenwich, **1993**, pp. 27–66.
- [2] R. Willstätter, E. Waser, *Ber. Dtsch. Chem. Ges.* **1911**, *44*, 3423–3445.
- [3] W. Reppe, O. Schlichting, K. Klager, T. Toepel, *Justus Liebigs Ann. Chem.* **1948**, *560*, 1–92.
- [4] a) For a thematic issue on green chemistry, see: *Chem. Rev.* **2007**, *107*, 2167–2820; b) for the principles of green chemistry, see: P. T. Anastas, J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University Press, New York, **1998**, p. 30; c) for more information, visit the ACS Green Chemistry Institute at <http://www.chemistry.org/portal/a/c/s/1/acsgdisplay.html?DOC=greenchemistryinstitute/index.html>.
- [5] For a recent discussion of redox economy, see: N. Z. Burns, P. S. Baran, R. W. Hoffmann, *Angew. Chem. Int. Ed.* **2009**, *48*, 2854–2867.
- [6] a) For recent and seminal discussions of atom economy, see: C.-J. Li, B. M. Trost, *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 13197–13202 and b) B. M. Trost, *Science* **1991**, *254*, 1471–1477.
- [7] D. A. Nicewicz, A. D. Satterfield, D. C. Schmitt, J. S. Johnson, *J. Am. Chem. Soc.* **2008**, *130*, 17281–17283.
- [8] a) For the first [5+2] reaction of a VCP, see: P. A. Wender, H. Takahashi, B. Witulski, *J. Am. Chem. Soc.* **1995**, *117*, 4720–4721; b) for the first intermolecular [5+2] reaction, see: P. A. Wender, H. Rieck, M. Fuji, *J. Am. Chem. Soc.* **1998**, *120*, 10976–10977; c) for the first [5+2] reaction with an alkene, see: P. A. Wender, C. O. Husfeld, E. Langkopf, J. A. Love, N. Pleuss, *Tetrahedron* **1998**, *54*, 7203–7220; d) for the first [5+2] reaction with allenes, see: P. A. Wender, F. Glorius, C. O. Husfeld, E. Langkopf, J. A. Love, *J. Am. Chem. Soc.* **1999**, *121*, 5348–5349; e) for the first intermolecular [5+2] reaction with allenes, see: H. A. Wegner, A. de Meijere, P. A. Wender, *J. Am. Chem. Soc.* **2005**, *127*, 6530–6531; f) for the first [5+2] reaction with cyclopropyl imines, see: P. A. Wender, T. M. Pedersen, M. J. C. Scanio, *J. Am. Chem. Soc.* **2002**, *124*, 15154–15155.
- [9] a) I. Ryu, K. Ikura, Y. Tamura, J. Maenaka, A. Ogawa, N. Sonoda, *Synlett* **1994**, 941–942; b) M. F. Salomon, R. G. Salomon, *J. Chem. Soc., Chem. Commun.* **1976**, 89–90; c) H. W. Voight, J. A. Roth, *J. Catal.* **1974**, *33*, 91–97; d) R. G. Salomon, M. F. Salomon, J. L. C. Kachinski, *J. Am. Chem. Soc.* **1977**, *99*, 1043–1054; e) M. P. Doyle, D. Van Leusen, *J. Am. Chem. Soc.* **1981**, *103*, 5917–5919; f) M. P. Doyle, D. Van Leusen, *J. Org. Chem.* **1982**, *47*, 5326–5338.
- [10] For computational analysis of the mechanism of the [5+2] reaction using density functional theory, see: Z.-X. Yu, P. A. Wender, K. N. Houk, *J. Am. Chem. Soc.* **2004**, *126*, 9154–9155.
- [11] P. A. Wender, A. G. Correa, Y. Sato, R. Sun, *J. Am. Chem. Soc.* **2000**, *122*, 7815–7816.
- [12] a) For a [6+1] reaction allenylcyclobutanes, see: P. A. Wender, N. M. Deschamps, R. Sun, *Angew. Chem. Int. Ed.* **2006**, *45*, 3957–3960; b) for the carbonylative rearrangement of allenic ethers, see: P. A. Wender, N. M. Deschamps, R. Sun, *Can. J. Chem.* **2005**, *83*, 838–842.
- [13] P. A. Wender, T. J. Paxton, T. J. Williams, *J. Am. Chem. Soc.* **2006**, *128*, 14814–14815.
- [14] P. A. Wender, G. G. Gamber, R. D. Hubbard, L. Zhang, *J. Am. Chem. Soc.* **2002**, *124*, 2876–2877.
- [15] P. A. Wender, G. G. Gamber, R. D. Hubbard, S. M. Pham, L. Zhang, *J. Am. Chem. Soc.* **2005**, *127*, 2836–2837.
- [16] P. A. Wender, N. C. Ihle, *J. Am. Chem. Soc.* **1986**, *108*, 4678–4679.
- [17] a) P. A. Wender, T. E. Jenkins, *J. Am. Chem. Soc.* **1989**, *111*, 6432–6434; for related subsequent reports by Livinghouse and Wender using alkenes and allenes, see: b) R. S. Jolly, G. Luedtke, D. Sheehan, T. Livinghouse, *J. Am. Chem. Soc.* **1990**, *112*, 4965–4966; c) P. A. Wender, T. E. Jenkins, S. Suzuki, *J. Am. Chem. Soc.* **1995**, *117*, 1843–1844.
- [18] For the trapping of intermediates in the [4+2] reaction with an alkyne to achieve a [4+2+2] reaction, see: P. A. Wender, J. P. Christy, *J. Am. Chem. Soc.* **2006**, *128*, 6530–6531.
- [19] a) For the intramolecular diene-yne [4+2+1] and [2+2+1] reactions, see P. A. Wender, N. M. Deschamps, G. G. Gamber, *Angew. Chem. Int. Ed.* **2003**, *42*, 1853–1857; b) for the intermolecular dienyl-Pauson–Khand reaction, see: P. A. Wender, N. M. Deschamps, T. J. Williams, *Angew. Chem. Int. Ed.* **2004**, *43*, 3076–3079; c) for the diene-ene [2+2+1] reaction, see: P. A. Wender, M. P. Croatt, N. M. Deschamps, *J. Am. Chem. Soc.* **2004**, *126*, 5948–5949; d) for the diene-allene [2+2+1] reaction, see: P. A. Wender, M. P. Croatt, N. M. Deschamps, *Angew. Chem. Int. Ed.* **2006**, *45*, 2459–2462.
- [20] For recent reviews including work from the groups of Brummond, Buchwald, Carretero, Chung, Cook, Greene, Jeong, Krafft, Livinghouse, Mitsudo, Murai, Mukai, Narasaka, Perias, Periasamy, Schore, Rautenstrauch, and Yamaguchi, see: a) T. Shibata, *Adv. Synth. Catal.* **2006**, *348*, 2328–2336; b) D. Strübing, M. Beller, *Top. Organomet. Chem.* **2006**, *18*, 165–178; c) J. Blanc-Urgoiti, L. Anorbe, L. Perez-Serrano, G. Dominguez, J. Perez-Castells, *J. Chem. Soc. Rev.* **2004**, *33*, 32–42; d) S. E. Gibson, A. Stevenazzi, *Angew. Chem. Int. Ed.* **2003**, *42*, 1800–1810; e) B. E. Hanson, *Comments Inorg. Chem.* **2002**, *23*, 289–318; f) K. M. Brummond, J. L. Kent, *Tetrahedron* **2000**, *56*, 3263–3283; g) A. J. Fletcher, S. D. R. Christie, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1657–1668; h) L. Yet, *Technical Report No. 18*, Vol. 4, Albany Molecular Research Inc., New York, **2000**.
- [21] a) For an example using a stoichiometric organometallic reagent as one of the components in a [4+2+1] reaction, see: J. W. Herndon, G. Chatterjee, P. P. Patel, J. J. Matsi, S. U. Turner, J. J. Harp, M. D. Reid, *J. Am. Chem. Soc.* **1991**, *113*, 7808–7809; b) for a metal-catalyzed [4+2+1] reaction of a diene-yne and trimethylsilyldiazomethane subsequently reported, see: J. Montgomery, Y. Ni, *J. Am. Chem. Soc.* **2004**, *126*, 11162–11163.
- [22] a) For subsequent reports of the dienyl-Pauson–Khand reaction, see: M.-C. P. Yeh, W.-C. Tsao, J.-S. Ho, C.-C. Tai, D.-Y. Chiou, L.-H. Tu, *Organometallics* **2004**, *23*, 792–799; b) K. H. Park, S. Y. Choi, S. Y. Kim, Y. K. Chung, *Synlett* **2006**, *4*, 527–532; c) for an initially reported [3+2+1] reaction that was corrected to be a diene-ene [2+2+1] reaction, see: Y. K. Chung, S. I. Lee, J. H. Park, S.-G. Lee, *J. Am. Chem. Soc.* **2004**, *126*, 10190 [*J. Am. Chem. Soc.* **2004**, *126*, 2714–2715].
- [23] a) For the first report, see: I. U. Khand, G. R. Knox, P. L. Pauson, W. E. Watts, *J. Chem. Soc., Chem. Commun.* **1971**, 36; b) for a report of an enantioselective intermolecular Pauson–Khand reaction, see: T. Shibata, K. Takagi, *J. Am. Chem. Soc.* **2000**, *122*, 9852–9853; c) for a review of components in the intermolecular Pauson–Khand reaction, see: S. E. Gibson, N. Mainolfi, *Angew. Chem. Int. Ed.* **2005**, *44*, 3022–3037.
- [24] a) For a seminal contribution, see: M. E. Krafft, *J. Am. Chem. Soc.* **1988**, *110*, 968–970; b) for a more recent example, see: K. Itami, K. Mitsudo, K. Fujita, Y. Ohashi, J. Yoshida, *J. Am. Chem. Soc.* **2004**, *126*, 11058–11066.
- [25] T. Kobayashi, Y. Koga, K. Narasaka, *J. Organomet. Chem.* **2001**, *624*, 73–87.
- [26] a) D. R. Swanson, C. J. Rousset, E. Negishi, T. Takahashi, T. Seki, M. Saburi, Y. Uchida, *J. Org. Chem.* **1989**, *54*, 3521–3523; b) E. Negishi, D. Choueiry, T. B. Nguyen, D. R. Swanson, N. Suzuki, T. Takahashi, *J. Am. Chem. Soc.* **1994**, *116*, 9751–9752.
- [27] K. Itoh, T. Makino, *J. Org. Chem.* **2004**, *69*, 395–405.
- [28] The authors report *trans*-fusion of the rings of **50** based on NOESY crosspeaks; however, the spectroscopic data for **50** match those of **55**, which appears to have *cis*-fusion of the rings based on strong 1D nOe enhancement between the two angular hydrogen atoms. Importantly, the diastereotopic methyl peaks from the malonate tether in **49** appear as a 6-hydrogen singlet in the 1H NMR spectrum although the 1H NMR spectrum of **50** and all analogous products formed from the diene-ene



- [2+2+1] reaction with a malonate tether have significant separation of the two methyl peaks. Based on this data, it appears that **50** is incorrectly assigned and is actually *cis*-fused and that **49** is correctly assigned as the *trans*-fused bicycle.
- [29] K. M. Brummond, H. Chen, B. Mitasev, A. D. Casarex, *Org. Lett.* **2004**, *6*, 2161–2163.
- [30] N. G. Rondan, M. N. Paddon-Row, P. Caramella, J. Mareda, P. H. Mueller, K. N. Houk, *J. Am. Chem. Soc.* **1982**, *104*, 4974–4976.
- [31] N. A. Sheddin, J. Mulzer, *Org. Lett.* **2006**, *8*, 3101–3104.
- [32] a) R. M. Moriarty, N. Rani, L. A. Enache, M. S. Rao, H. Batra, L. Guo, R. A. Penmasta, J. P. Staszewski, S. M. Tuladhar, O. Prakash, D. Crich, A. Hirtopeanu, R. Gilardi, *J. Org. Chem.* **2004**, *69*, 1890–1902; b) R. M. Moriarty, H. Batra, WO 02053517, EP 1355872, US 680922; c) R. M. Moriarty, R. Penmasta, L. Guo, M. S. Rao, J. P. Staszewski, WO 9921830, EP1025083, CA 2307163.
- [33] It should be noted that in ref.<sup>[32b]</sup>, 10 mol-% catalyst was reported to have a 73% yield on the gram-scale, but ref.<sup>[32a]</sup> was run with stoichiometric catalyst loading on a 4.9 kilogram scale.
- [34] For a review of the allenic Pauson–Khand reaction including significant contributions from the groups of Brummond, Cazes, Livinghouse, Alcaide, Mukai, Cook, and Shibata, see: B. Alcaide, P. Almendros, *Eur. J. Org. Chem.* **2004**, 3377–3383.
- [35] For a rationale for the olefin selectivity in the Rh<sup>I</sup>-catalyzed allenic Pauson–Khand reaction based on DFT calculations, see: A. S. Bayden, K. M. Brummond, K. D. Jordan, *Organometallics* **2006**, *25*, 5204–5206.
- [36] a) For the first [2+2+1] reaction of a sulfonyl allene and an alkene, see: F. Inagaki, C. Mukai, *Org. Lett.* **2006**, *8*, 1217–1220; b) for a recent [2+2+1] reaction of sulfonyl allenes with a tethered allene, see: F. Inagaki, S. Narita, T. Hasegawa, S. Kitagaki, C. Mukai, *Angew. Chem. Int. Ed.* **2009**, *48*, 2007–2011.
- [37] M. J. Kamlet, J.-L. M. Abboud, M. H. Abraham, R. W. Taft, *J. Org. Chem.* **1983**, *48*, 2877–2887.
- [38] R. W. Taft, M. J. Kamlet, *J. Am. Chem. Soc.* **1976**, *98*, 2886–2894.
- [39] a) For examples of a diene acting as a two-carbon component but accelerating the reaction relative to an isolated alkene, see: M. Gulías, J. Durán, F. López, L. Castedo, J. L. Mascareñas, *J. Am. Chem. Soc.* **2007**, *129*, 11026–11027; b) H. Du, B. Zhao, Y. Shi, *J. Am. Chem. Soc.* **2007**, *129*, 762–763.
- [40] For a recent report which examined the mechanism of the diene-ene [2+2+1] reaction with DFT calculations, see: W. H. Pitcock Jr, R. L. Lord, M.-H. Baik, *J. Am. Chem. Soc.* **2008**, *130*, 5821–5830.
- [41] This step would additionally involve conversion of the dimeric rhodium catalyst into two monomeric rhodium catalysts. This is presumably facile in the presence of dienes.
- [42] This structure could be an intermediate or the transition state between two intermediates with rhodium coordinated to either the alpha or beta face of the alkene.
- [43] J. Halpern, T. Okamoto, A. Zakhariev, *J. Mol. Catal.* **1976**, *2*, 5–68.
- [44] G. O. Spessard, G. L. Miessler in *Organometallic Chemistry*, Prentice-Hall Inc., New Jersey, **1997**, pp. 277–283.
- [45] The intramolecular dienyl-Pauson–Khand reaction was explored mostly using a different catalyst system, RhCl(CO)(PPh<sub>3</sub>)<sub>2</sub>/AgSbF<sub>6</sub>, but to compare all of the reactions of dienes tethered to  $\pi$ -systems, a single catalyst system was chosen to study, [RhCl(CO)<sub>2</sub>]<sub>2</sub>.
- [46] For related examples of rhodium–acyl shifts in <sup>13</sup>C NMR and Rh–C coupling constants, see: a) S. C. van der Slot, P. C. J. Kamer, P. W. N. M. van Leeuwen, J. A. Iggo, B. T. Heaton, *Organometallics* **2001**, *20*, 430–441; b) M. A. Huffman, L. S. Liebeskind, *Organometallics* **1992**, *11*, 255–266; c) J. M. Brown, A. G. Kent, *J. Chem. Soc., Chem. Commun.* **1982**, 723–725.
- [47] a) Y. Wang, J. Wang, J. Su, F. Huang, L. Jiao, Y. Liang, D. Yang, S. Zhang, P. A. Wender, Z.-X. Yu, *J. Am. Chem. Soc.* **2007**, *129*, 10060–10061; b) Z.-X. Yu, P. H.-Y. Cheong, P. Liu, C. Y. Legault, P. A. Wender, K. N. Houk, *J. Am. Chem. Soc.* **2008**, *130*, 2378–2379; c) P. Liu, P. H.-Y. Cheong, Z.-X. Yu, P. A. Wender, K. N. Houk, *Angew. Chem. Int. Ed.* **2008**, *47*, 3939–3941.
- [48] a) S. Gilbertson, B. DeBoef, *J. Am. Chem. Soc.* **2002**, *124*, 8784–8785; b) P. A. Evans, J. E. Robinson, E. W. Baum, A. N. Fazal, *J. Am. Chem. Soc.* **2003**, *125*, 14648 [*J. Am. Chem. Soc.* **2002**, *124*, 8782–8783]; c) P. A. Evans, E. W. Baum, *J. Am. Chem. Soc.* **2004**, *126*, 11150–11151; d) P. A. Evans, E. W. Baum, E. W. Fazal, N. Aleem, M. Pink, *Chem. Commun.* **2005**, *1*, 63–65; e) M.-H. Baik, E. W. Baum, M. C. Burland, P. A. Evans, *J. Am. Chem. Soc.* **2005**, *127*, 1602–1603; f) ref.<sup>[18]</sup>; g) B. DeBoef, W. R. Counts, S. R. Gilbertson, *J. Org. Chem.* **2007**, *72*, 799–804.

Received: August 15, 2009

Published Online: October 14, 2009