

Erratum

Differential changes in induced seizures after hippocampal treatment of rats with an antisense oligodeoxynucleotide to the GABA_A receptor $\gamma 2$ subunit [Eur. J. Pharmacol., 340 (1997) 153–160]¹

Jesper Karle^{a,*}, Peter Laudrup^b, Frank Sams-Dodd^b, Jens D. Mikkelsen^b, Mogens Nielsen^a

^a *The Research Institute of Biological Psychiatry, St. Hans Hospital, DK-4000 Roskilde, Denmark*

^b *H. Lundbeck, Department of Psychopharmacology, Ottiliavej 9, DK-2500 Valby, Denmark*

In the above-mentioned article Table 1, which is shown below, was inadvertently omitted. The Publisher apologises for any inconvenience caused.

Table 1

Results of the s.c. pentylenetetrazol test: seizure scores from two independent experiments are shown

	Median seizure score	25% quartile	75% quartile
Experiment 1			
Naive	4	4	4
Mismatch	3 ^a	1	3.75
Antisense	5 ^b	5	6
Experiment 2			
Naive	3	3	4.75
Mismatch	3	1.25	3
Antisense	6	2.25	6

^a $P < 0.05$ vs. naive; $n = 7$.

^b $P < 0.05$ vs. mismatch and vs. naive; $n = 7$ (Mann–Whitney U -test).

* Corresponding author. Department of Psychiatry, Rigshospitalet (National Hospital), Blegdamsvej 9, DK-2100 Copenhagen Ø, Denmark. Tel.: +45-35-456154; fax: +45-35-456218.

¹ PII of original article: S0014299997014234.