

Erratum

Erratum to: “Enhanced δ -opioid receptor-mediated antinociception in μ -opioid receptor-deficient mice” [Eur. J. Pharmacol. 387 (2000) 163–169]¹

Chunyuan Qiu^{a,b}, Ichiro Sora^c, Ke Ren^a, George Uhl^c, Ronald Dubner^{a,*}

^a Department of Oral and Craniofacial Biological Sciences, University of Maryland Dental School,
Room 5E-08, 666 West Baltimore Street, Baltimore, MD 21201, USA

^b Department of Anesthesiology, School of Medicine, University of Maryland, Baltimore, MD 21201, USA

^c NIDA-IRP and Department of Neurology and Neuroscience, Johns Hopkins University School Medicine, Baltimore, MD 21224, USA

Received 12 August 1999; received in revised form 5 November 1999; accepted 12 November 1999

The correct Fig. 1 of the above-mentioned article is given below.




Fig. 1. Baseline paw withdrawal latencies. There were significant differences in paw withdrawal latencies between μ -opioid receptor knockout ($-/-$), heterozygous ($+/-$) and wild-type ($+/+$) mice when the data from males and females were separated or combined. μ -opioid receptor knockout males and females had significantly lower paw withdrawal latencies than wild-type mice. $**P < 0.01$, $***P < 0.001$; μ -opioid receptor knockout female mice and the combined group had significantly lower paw withdrawal latencies than heterozygous mice, $\#P < 0.05$. The numbers below the histogram bars indicate the number of mice per group.

* Corresponding author. Tel.: +1-410-706-0860; fax: +1-410-706-0865.

E-mail address: rnd001@dental.umaryland.edu (R. Dubner).

¹ PII of original article S001429999008134.