

Corrigendum to “Neuropharmacological profiles of a novel atypical antipsychotic, NRA0562, in rats” [Eur. J. Pharmacol. 423 (2001) 27–33][☆]

Naoya Kawashima^{*}, Takeo Funakoshi, Tomohiro Omura, Shigeyuki Chaki,
Kazuya Kameo, Shigeru Okuyama

*CNS Diseases Research, Medicinal Pharmacology Laboratory, Medicinal Research Laboratories, Taisho Pharmaceutical Co., Ltd.,
1-403 Yoshino-cho, Saitama, Saitama 330-8530, Japan*

The authors regret that in the above-mentioned article errors were made.

The following points need to be considered: Figs. 3 and 4 should change place with each other. Fig. 3 should show

recording data of electrophysiology and Fig. 4 should show graphs. The legends are correct. The correct figures plus legends are given on the following page.

[☆] PII of original article S0014-2999(01)01085-8

^{*} Corresponding author.

E-mail address: n.kawashima@po.rd.taisho.co.jp (N. Kawashima).




Fig. 3. Reversal by NRA0562 of the inhibitory effects of methamphetamine (1 mg/kg, i.v.) on A10 (A) and A9 (B) dopamine neurons.




Fig. 4. Comparison of the potential of NRA0562 (A) and risperidone (B) to reverse the inhibitory effects of methamphetamine (1 mg/kg, i.v.) on A10 (O) and A9 (●) dopamine neurons. The results are presented as means with vertical lines showing S.E. ($n = 6$). Starting doses of NRA0562 and risperidone were 0.1 and 0.01 mg/kg, i.v., respectively. ** $P < 0.01$ vs. pre-drug value (Dunnett's test). ## $P < 0.01$ vs. methamphetamine + vehicle-treated group (Dunnett's test).