

Erratum

Erratum to “Functional role of nicotinic acetylcholine receptors in apoptosis in HL-60 cell line” [Eur. J. Pharmacol. 482 (2003) 25–29][☆]

Delphine Gimonet^{a,1}, Régis Grailhe^{b,2}, Paul Coninx^{a,1}, Frank Antonicelli^{c,3},
Bernard Haye^{c,3}, Françoise Liautaud-Roger^{a,*}

^a *Institut Jean-Godinot, Secteur Prévention, BP 171, 1 Avenue du général Koenig, 51056, Reims Cedex, France*

^b *Laboratoire Récepteur et Cognition, Institut Pasteur, 25 rue du Docteur Roux, 75015, Paris, France*

^c *FRE CNRS 2534, Faculté des Sciences, Université de Reims, Moulin de la Housse, BP 1039, 51687 Cedex 2, France*

The publisher regrets that Fig. 4 in the above-mentioned article was erroneously published. Herewith, the correct Fig. 4 is given.




Fig. 4. Effect of increasing concentrations of three agonists (X-axis) on bleomycin (BLM)-induced apoptosis (Y-axis). Only the agonist specific for $\alpha 7$ -nicotinic acetylcholine receptor (i.e. 3-[2,4-dimethoxybenzylidene]anabaseine (GTS-21)) produced an increase of apoptosis (** $P < 0.01$). Error bars represent the 95 confidence interval. The insert at the top indicates the time course of the experiment.

[☆] doi of original article 10.1016/j.ejphar.2003.09.050.

* Corresponding author. Tel.: +33-3-26-50-42-87; fax: +33-3-26-50-43-98.

E-mail addresses: rgrailhe@pasteur.fr (R. Grailhe), frank.antoncelli@univ-reims.fr (F. Antonicelli), francoise.roger@reims.fnclcc.fr (F. Liautaud-Roger).

¹ Tel.: +33-3-26-50-42-87; fax: +33-3-26-50-43-98.

² Tel./fax: +33-1-45-68-88-46.

³ Tel.: +33-3-26-91-34-35; fax: +33-3-26-91-31-68.