



Erratum

Erratum to “Isobolographic analysis of the antinociceptive interactions between ketoprofen and paracetamol”

[European Journal of Pharmacology 557/2-3 (2007) 141–146]

Hai-Xia Qiu, Jin Liu, Hui Kong, Yan Liu, Xing-guo Mei *

In Table 2 on page 143 the γ -value for the 2:1 combination should be 0.586 instead of 0.850. The correct Table 2 is shown below:

Table 2
Effect of ketoprofen, paracetamol and their fixed-ratio combinations in the mouse abdominal constriction test.

Paracetamol/ketoprofen combinations	ED _{50,mix} (mg/kg)	ED _{50,add} (mg/kg)	γ
Ketoprofen alone	37.60 ± 1.708	–	–
Paracetamol alone	129.35 ± 7.264	–	–
2:1	41.84 ± 4.392 ^a	71.33 ± 2.196	0.586
3:1	37.60 ± 3.556 ^a	80.34 ± 1.778	0.427
4:1	37.60 ± 2.940 ^a	86.93 ± 1.470	0.456

Data are presented as ED₅₀ ± S.E.M. Statistical analysis was performed with Student's *t*-test. ^a*P* < 0.01 vs. the respective additive group. γ < 1 indicates supra-additivity (synergy).

DOI of original article: [10.1016/j.ejphar.2006.11.017](https://doi.org/10.1016/j.ejphar.2006.11.017).

* Corresponding author.

E-mail addresses: mountee@gmail.com (H.-X. Qiu), xg_mei@yahoo.com (X. Mei).