

delde planten (Fig.1). Vers gewichten van zieke bladeren waren kleiner dan die van onbehandelde planten (Fig.2). Het watergehalte van zieke bladeren werd lager dan dat van onbehandelde bladeren (Fig.3). De uredosporenproduktie was bij hogere grondwaterpotentiaal groter dan bij lagere potentiaal (Fig.4).

References

- Bever, W. M., 1937. The influence of stripe rust on growth, water economy, and yield of wheat and barley. *J. agric. Res.* 54: 375-385.
- Johston, C. O. & Miller, E. C., 1934. Relation of leaf rust infection to yield, growth, and water economy of two varieties of wheat. *J. agric. Res.* 49: 955-981.
- Mehta, Y. R. & Zadoks, J. C., 1970. Uredospore production and sporulation period of *Puccinia recondita* f. sp. *trititica* on primary leaves of wheat. *Neth. J. Pl. Path.* 76: 267-276.
- Monteith, J. L. & Owen, P. C., 1958. A thermocouple method for measuring relative humidity in the range 95% to 100%. *J. scient. Instrum.* 35: 443-446.
- Murphy, H. C., 1935. Effect of crown rust infection on yield and water requirement of oats. *J. agric. Res.* 1: 387-411.
- Parodi, P. P. & Bitzer, M. J., 1969. Determinación simultánea de absorción de agua y transpiración de trigo (*Triticum aestivum* L.) infectadas y libres de polvillo de la hoja (*Puccinia recondita* Rob. ex Desm. f. sp. *tritici*). *Agricultura téc.* 29: 191-194.
- Slatyer, R. O. & Taylor, S. A., 1960. Terminology in plant water and soil water relations. *Nature, Lond.* 187: 922-924.

Addresses

- M. C. Cowan: Pye Research Centre, Stowmarket (Suffolk), England.
- J. C. Zadoks: Laboratorium voor Fytopathologie, Binnenhaven 9, Wageningen, the Netherlands.

Publications received

- I. D. Firman: A list of fungi and plant parasitic bacteria, viruses and nematodes in Fiji. Commonwealth Mycological Institute, Kew, Surrey, England, 1972, pp. 36. Price £ 0.60.
- W. T. H. Peregrine & M. A. Siddiqi: A revised and annotated list of plant diseases in Malawi. Commonwealth Mycological Institute, Kew, Surrey England, 1972, pp. 51. Price £ 0.80.