

Addresses

D. Mulder: Laboratorium voor Fytopathologie, Landbouwhogeschool, Binnenhaven 9, Wageningen, the Netherlands.

L. Herrera and R. Garcia: Laboratorio de Phytopatología, Universidad de Santa Clara, Santa Clara, Cuba.

Book review

F. J. Gommers: Nematicidal principles in Compositae. Diss., Rijksuniversiteit Groningen 1973. Also published as: Mededelingen van de Landbouwhogeschool Wageningen 73-17: 71 pp.

After earlier work carried out by others mainly at Wageningen, the Netherlands, Gommers investigated the suppressing properties of members of the Compositae on nematode populations at the Groningen and Wageningen universities. The plant parasite *Pratylenchus penetrans* is particularly sensitive to various Compositae. This is related to its endoparasitic behaviour. In most experiments this species served as the test organism. Nematodes may be killed by Compositae after having entered the roots. There are no indications that nematicidal compounds released into the soil play a role. Some ectoparasitic nematode species multiply strongly on certain Compositae suppressing *P. penetrans* populations.

In a series of field and glasshouse experiments several additional Compositae with suppressing activity were found. Taxonomic relationships became evident: species with suppressing capacity were frequently detected among the tribes Heliantheae and Helenieae, and not, for instance, among the tribe Anthemideae.

From Compositae different fractions with in vitro nematicidal activity could be isolated. An interesting discovery is that the in vitro activity of a number of these compounds (including α -terthienyl) is strongly enhanced by daylight. The near ultra-violet wavelengths were shown to be responsible for this effect. At the same wavelengths these compounds also show the greatest absorption. In vitro experiments with α -terthienyl in light and darkness showed that the compound is released from the nematodes more slowly than it is taken up. *P. penetrans* from roots of *Tagetes* were also rapidly killed when exposed to light of the same wavelengths.

H. Hoestra

K. F. Baker & R. J. Cook: Biological control of plant pathogens (with a foreword by S. D. Garrett). W. H. Freeman and Company, San Francisco, USA 1973: 433 pp.; 58 figs.; subject index; cloth bound. Price \$ 12.50, £ 5.90.

This book deals primarily with soil-borne diseases with only one chapter on pathogens of aerial parts. Much less is known of biological control of the latter than of soil-borne diseases.

The authors treat the subject clearly and systematically. After some chapters on general subjects such as biological balances, and the general nature of biological control, the roles of the pathogen, the antagonist, the host and the physical environment are discussed, each in a separate chapter. Much emphasis is laid on antagonistic micro-organisms. The authors present strong evidence that antagonists play an important role in limiting diseases, and might be used in more cases to control soil-borne diseases. They indicate approaches and techniques for this, but do not suggest that biological control is easy to achieve.

The book contains a wealth of well presented information and is pleasant to read. Not all views presented will be shared by all readers. The authors are aware of this and in their preface state that they will be content if, among other things, the book 'even irritates someone into proving us wrong'. In my opinion, many readers will be convinced that the time has come for a new interest in biological control of soil-borne plant diseases.

H. Hoestra