

## **Volume contents**

### **Volume 107, No. 1 January 2001**

#### **Special Issue**

#### **Induced Resistance to Plant Diseases**

**Papers presented at an International Symposium held in Corfu 22–27 May, 2000**

|                                                                                                                                                                                    |                |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Foreword</b>                                                                                                                                                                    | <b>vii</b>     |
| <b>Inducing resistance: a summary of papers presented at the First International Symposium on Induced Resistance to Plant Diseases, Corfu, May 2000</b>                            | <b>1–6</b>     |
| R. Hammerschmidt, J.-P. Métraux, L.C. van Loon                                                                                                                                     |                |
| <b>Concepts and direction of induced systemic resistance in plants and its application</b>                                                                                         | <b>7–12</b>    |
| J. Kuć                                                                                                                                                                             |                |
| <b>Systemic acquired resistance and salicylic acid: current state of knowledge</b>                                                                                                 | <b>13–18</b>   |
| J.-P. Métraux                                                                                                                                                                      |                |
| <b>Induced disease resistance in plants by chemicals</b>                                                                                                                           | <b>19–28</b>   |
| M. Oostendorp, W. Kunz, B. Dietrich, T. Staub                                                                                                                                      |                |
| <b><math>\beta</math>-Aminobutyric acid-induced resistance in plants</b>                                                                                                           | <b>29–37</b>   |
| G. Jakab, V. Cottier, V. Toquin, G. Rigoli, L. Zimmerli, J.-P. Métraux, B. Mauch-Mani                                                                                              |                |
| <b>Application of rhizobacteria for induced resistance</b>                                                                                                                         | <b>39–50</b>   |
| G.W. Zehnder, J.F. Murphy, E.J. Sikora, J.W. Kloepper                                                                                                                              |                |
| <b>Rhizobacteria-mediated induced systemic resistance: triggering, signalling and expression</b>                                                                                   | <b>51–61</b>   |
| C.M.J. Pieterse, J.A. Van Pelt, S.C.M. Van Wees, J. Ton, K.M. Léon-Kloosterziel, J.J.B. Keurentjes, B.W.M. Verhagen, M. Knoester, I. Van der Sluis, P.A.H.M. Bakker, L.C. Van Loon |                |
| <b>Heritability of rhizobacteria-mediated induced systemic resistance and basal resistance in <i>Arabidopsis</i></b>                                                               | <b>63–68</b>   |
| J. Ton, S. Davison, L.C. Van Loon, C.M.J. Pieterse                                                                                                                                 |                |
| <b>The hypersensitive response and its role in local and systemic disease resistance</b>                                                                                           | <b>69–78</b>   |
| E. Kombrink, E. Schmelzer                                                                                                                                                          |                |
| <b>Plant–pathogen interactions: genetic and comparative analyses</b>                                                                                                               | <b>79–84</b>   |
| A.H. Ellingboe                                                                                                                                                                     |                |
| <b>The relationship between pathogen-induced systemic resistance (ISR) and multigenic (horizontal) resistance in plants</b>                                                        | <b>85–93</b>   |
| S. Tuzun                                                                                                                                                                           |                |
| <b>Bacterial lipopolysaccharides and plant–pathogen interactions</b>                                                                                                               | <b>95–102</b>  |
| M.-A. Newman, J.M. Dow, M.J. Daniels                                                                                                                                               |                |
| <b>Signal interactions in induced resistance to pathogens and insect herbivores</b>                                                                                                | <b>103–111</b> |
| R.M. Bostock, R. Karban, J.S. Thaler, P.D. Weyman, D. Gilchrist                                                                                                                    |                |
| <b>Priming as a mechanism in induced systemic resistance of plants</b>                                                                                                             | <b>113–119</b> |
| U. Conrath, O. Thulke, V. Katz, S. Schwindling, A. Kohler                                                                                                                          |                |
| <b>Signal transduction in resistance to plant viruses</b>                                                                                                                          | <b>121–128</b> |
| A.M. Murphy, A. Gilliland, C.E. Wong, J. West, D.P. Singh, J.P. Carr                                                                                                               |                |

|                                                                                               |                |
|-----------------------------------------------------------------------------------------------|----------------|
| <b>Oxycom™ under field and laboratory conditions increases resistance responses in plants</b> | <b>129–136</b> |
| Y.C. Kim, K.A. Blee, J. Robins, A.J. Anderson                                                 |                |
| <b>The ecological concept of costs of induced systemic resistance (ISR)</b>                   | <b>137–146</b> |
| M. Heil                                                                                       |                |
| <b>Instructions to authors</b>                                                                | <b>147–151</b> |

## Volume 107, No. 2 February 2001

### Mini review

|                                                         |                |
|---------------------------------------------------------|----------------|
| <b>Hrp pilus – reaching through the plant cell wall</b> | <b>153–160</b> |
| M. Romantschuk, E. Roine, S. Taira                      |                |

### Research articles

|                                                                                                                                                                                                                              |                |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>A greenhouse test for screening sugar beet (<i>Beta vulgaris</i>) for resistance to <i>Rhizoctonia solani</i></b>                                                                                                         | <b>161–166</b> |
| O.E. Scholten, L.W. Panella, T.S.M. De Bock, W. Lange                                                                                                                                                                        |                |
| <b>The effect of temperature on variation in transmission of a BYDV PAV-like isolate by clones of <i>Rhopalosiphum padi</i> and <i>Sitobion avenae</i></b>                                                                   | <b>167–173</b> |
| I.N. Smyrnioudis, R. Harrington, M. Hall, N. Katis, S.J. Clark                                                                                                                                                               |                |
| <b><i>In planta</i> – complementation of <i>Clavibacter michiganensis</i> subsp. <i>sepedonicus</i> strains deficient in cellulase production or HR induction restores virulence</b>                                         | <b>175–182</b> |
| R. Nissinen, S. Kassuwi, R. Peltola, M.C. Metzler                                                                                                                                                                            |                |
| <b>Identification of the Chinese box orange (<i>Severinia buxifolia</i>) as an alternative host of the bacterium causing citrus Huanglongbing</b>                                                                            | <b>183–189</b> |
| T.-H. Hung, M.-L. Wu, H.-J. Su                                                                                                                                                                                               |                |
| <b>Resistance reaction of conifer species (European larch, Norway spruce, Scots pine) to infection by selected necrotrophic damping-off pathogens</b>                                                                        | <b>191–207</b> |
| M. Kacprzak, F.O. Asiegbu, G. Daniel, J. Stenlid, M. Mañka, M. Johansson                                                                                                                                                     |                |
| <b>Detection and indentification of European stone fruit yellows and other phytoplasmas in wild plants in the surroundings of apricot chlorotic leaf roll-affected orchards in southern France</b>                           | <b>209–217</b> |
| W. Jarausch, B. Jarausch-Wehrheim, J.L. Danet, J.M. Broquaire, F. Dosba, C. Saillard, M. Garnier                                                                                                                             |                |
| <b>Induced resistance to cyst and root-knot nematodes in cereals by DL-<math>\beta</math>-amino-<i>n</i>-butyric acid</b>                                                                                                    | <b>219–227</b> |
| Y. Oka, Y. Cohen                                                                                                                                                                                                             |                |
| <b>Survival of <i>Botrytis cinerea</i> in southeastern Spanish greenhouses</b>                                                                                                                                               | <b>229–236</b> |
| R. Raposo, V. Gomez, T. Urrutia, P. Melgarejo                                                                                                                                                                                |                |
| <b>Identification of pathogenic races 0, 1B/C, 5, and 6 of <i>Fusarium oxysporum</i> f. sp. <i>ciceris</i> with random amplified polymorphic DNA (RAPD)</b>                                                                  | <b>237–248</b> |
| M. del Mar Jiménez-Gasco, E. Pérez-Artés, R.M. Jiménez-Díaz                                                                                                                                                                  |                |
| <b>Molecular characterization of the incompatible interaction of <i>Vitis vinifera</i> leaves with <i>Pseudomonas syringae</i> pv. <i>pisi</i>: Expression of genes coding for stilbene synthase and class 10 PR protein</b> | <b>249–261</b> |
| N. Robert, J. Ferran, C. Breda, P. Coutos-Thévenot, M. Boulay, D. Buffard, R. Esnault                                                                                                                                        |                |

### Volume 107, No. 3 March 2001

|                                                                                                                                                                                                                    |                |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>A new natural planthopper vector of stolbur phytoplasma in the genus <i>Pentastiridius</i> (Hemiptera: Cixiidae)</b>                                                                                            | <b>263–271</b> |
| F. Gatineau, J. Larrue, D. Clair, F. Lorton, M. Richard-Molard, E. Boudon-Padieu                                                                                                                                   |                |
| <b>Comparative analysis of the role of substrate specificity in biological control of <i>Botrytis elliptica</i> in lily and <i>B. cinerea</i> in cyclamen with <i>Ulocladium atrum</i></b>                         | <b>273–284</b> |
| G.J.T. Kessel, B.H. de Haas, C.H. Lombaers-van der Plas, J.E. van den Ende, M.G. Pennock-Vos, W. van der Werf, J. Köhl                                                                                             |                |
| <b>Genotypic diversity of the wheat leaf blotch pathogen <i>Mycosphaerella graminicola</i> (anamorph) <i>Septoria tritici</i> in Germany</b>                                                                       | <b>285–290</b> |
| F. Schnieder, G. Koch, C. Jung, J.-A. Verreet                                                                                                                                                                      |                |
| <b>Phenotypic and genotypic characterisation of Northern Ireland isolates of <i>Phytophthora infestans</i></b>                                                                                                     | <b>291–303</b> |
| D.J. Carlisle, L.R. Cooke, A.E. Brown                                                                                                                                                                              |                |
| <b>Diversity of <i>Phytophthora clandestina</i> isolated from subterranean clover in southern Australia: analysis of virulence and RAPD profiles</b>                                                               | <b>305–311</b> |
| A. Purwantara, A. Drenth, S.P. Flett, W. Guppy, P.J. Keane                                                                                                                                                         |                |
| <b>Detection of <i>Plasmodiophora brassicae</i> by PCR in naturally infested soils</b>                                                                                                                             | <b>313–321</b> |
| A.-C. Wallenhammar, O. Arwidsson                                                                                                                                                                                   |                |
| <b>A new cell wall located N-rich protein is strongly induced during the hypersensitive response in <i>Glycine max</i> L.</b>                                                                                      | <b>323–336</b> |
| A.A. Ludwig, R. Tenhaken                                                                                                                                                                                           |                |
| <b>Fitness of <i>Cercospora beticola</i> field isolates – resistant and – sensitive to demethylation inhibitor fungicides</b>                                                                                      | <b>337–347</b> |
| G.S. Karaoglaniadis, C.C. Thanassouloupoulos, P.M. Ioannidis                                                                                                                                                       |                |
| <b>The use of a GUS transformant of <i>Trichoderma harzianum</i>, strain T3a, to study metabolic activity in the spermosphere and rhizosphere related to biocontrol of <i>Pythium</i> damping-off and root rot</b> | <b>349–359</b> |
| H. Green, N. Heiberg, K. Lejbølle, D.F. Jensen                                                                                                                                                                     |                |
| <i>Short communication</i>                                                                                                                                                                                         |                |
| <b>Rice yellow mottle virus is seed-borne but not seed transmitted in rice seeds</b>                                                                                                                               | <b>361–364</b> |
| G. Konate, S. Sarra, O. Traore                                                                                                                                                                                     |                |
| <b>Announcement</b>                                                                                                                                                                                                | <b>365</b>     |

### Volume 107, No. 4 May 2001

|                                                                                                                                                             |                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Population dynamics of <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> in relation to the onset of Fusarium crown and root rot of tomato</b> | <b>367–375</b> |
| Y. Rekah, D. Shtienberg, J. Katan                                                                                                                           |                |

|                                                                                                                                                                                                                                                            |                |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>The effect of treatment with <i>Ulocladium atrum</i> on <i>Botrytis cinerea</i>-attack of geranium (<i>Pelargonium zonale</i>) stock plants and cuttings</b>                                                                                            | <b>377–386</b> |
| M. Gerlagh, J.J. Amsing, W.M.L. Molhoek, A.I. Bosker-van Zessen, C.H. Lombaers-van der Plas, J. Köhl                                                                                                                                                       |                |
| <b>Conventional PCR and real-time quantitative PCR detection of <i>Helminthosporium solani</i> in soil and on potato tubers</b>                                                                                                                            | <b>387–398</b> |
| D.W. Cullen, A.K. Lees, I.K. Toth, J.M. Duncan                                                                                                                                                                                                             |                |
| <b>Genetic structure of <i>Melampsora epitea</i> populations in Swedish <i>Salix viminalis</i> plantations</b>                                                                                                                                             | <b>399–409</b> |
| B. Samils, U. Lagercrantz, M. Lascoux, U. Gullberg                                                                                                                                                                                                         |                |
| <b>Characterization and detection of several filamentous viruses of cherry: Adaptation of an alternative cloning method (DOP-PCR), and modification of an RNA extraction protocol</b>                                                                      | <b>411–420</b> |
| M.E. Rott, W. Jelkmann                                                                                                                                                                                                                                     |                |
| <b>Differential control of head blight pathogens of wheat by fungicides and consequences for mycotoxin contamination of grain</b>                                                                                                                          | <b>421–431</b> |
| D.R. Simpson, G.E. Weston, J.A. Turner, P. Jennings, P. Nicholson                                                                                                                                                                                          |                |
| <b>Interaction of root-knot nematodes (RKN) and the bacterium <i>Agrobacterium tumefaciens</i> in roots of <i>Prunus cerasifera</i>: evidence of the protective effect of the <i>Ma</i> RKN resistance genes against expression of crown gall symptoms</b> | <b>433–441</b> |
| M.-J. Rubio-Cabetas, J.-C. Minot, R. Voisin, D. Esmenjaud                                                                                                                                                                                                  |                |
| <b>Comparative study of genetic diversity and pathogenicity among populations of <i>Verticillium dahliae</i> from cotton in Spain and Israel</b>                                                                                                           | <b>443–456</b> |
| N. Korolev, E. Pérez-Artés, J. Bejarano-Alcázar, D. Rodríguez-Jurado, J. Katan, T. Katan, R.M. Jiménez-Díaz                                                                                                                                                |                |
| <b>Cloning, sequencing and expression of a xylanase gene from the maize pathogen <i>Helminthosporium turcicum</i></b>                                                                                                                                      | <b>457–465</b> |
| Y. Degefu, L. Paulin, P.S. Lübeck                                                                                                                                                                                                                          |                |
| <i>Short communication</i>                                                                                                                                                                                                                                 |                |
| <b>Scytalidium wilt of citrus</b>                                                                                                                                                                                                                          | <b>467–470</b> |
| Y. Oren, A. Sadowsky, D. Gefen, Z. Solel, M. Kimchy                                                                                                                                                                                                        |                |
| <b>Book reviews</b>                                                                                                                                                                                                                                        | <b>471–472</b> |

## Volume 107, No. 5 June 2001

|                                                                                                                    |                |
|--------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Activation of defense responses to <i>Fusarium</i> infection in <i>Asparagus densiflorus</i></b>                | <b>473–483</b> |
| C. He, T. Hsiang, D.J. Wolyn                                                                                       |                |
| <b>Characterization of a new nepovirus infecting apricot in Southeastern France: apricot latent ringspot virus</b> | <b>485–494</b> |
| P. Gentit, R.-P. Delbos, T. Candresse, J. Dunez                                                                    |                |
| <b>Effect of grapevine training systems on development of powdery mildew</b>                                       | <b>495–501</b> |
| T. Zahavi, M. Reuveni, D. Scheglov, S. Lavee                                                                       |                |
| <b>The epidemiology of purple leaf blotch on leeks in Victoria, Australia</b>                                      | <b>503–510</b> |
| H. Suheri, T.V. Price                                                                                              |                |

|                                                                                                                                                           |                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Phenazines are involved in biocontrol of <i>Pythium myriotylum</i> on cocoyam by <i>Pseudomonas aeruginosa</i> PNA1</b>                                | <b>511–521</b> |
| J.T. Tambong, M. Höfte                                                                                                                                    |                |
| <b>Induction of systemic resistance in cucumber against several diseases by plant growth-promoting fungi: lignification and superoxide generation</b>     | <b>523–533</b> |
| N. Koike, M. Hyakumachi, K. Kageyama, S. Tsuyumu, N. Doke                                                                                                 |                |
| <b>Spread of <i>Phytophthora</i> root and crown rot in <i>Saintpaulia</i>, <i>Gerbera</i> and <i>Spathiphyllum</i> pot plants in ebb-and-flow-systems</b> | <b>535–542</b> |
| D.J. van der Gaag, A. Kerssies, C. Lanser                                                                                                                 |                |
| <b>Nutritional enhancement of biocontrol activity of <i>Candida sake</i> (CPA-1) against <i>Penicillium expansum</i> on apples and pears</b>              | <b>543–551</b> |
| C. Nunes, J. Usall, N. Teixidó, M. Miró, I. Viñas                                                                                                         |                |
| <i>Short communications</i>                                                                                                                               |                |
| <b>Production and germination of oospores of <i>Phytophthora infestans</i> (Mont.) de Bary in Morocco</b>                                                 | <b>553–556</b> |
| A. Hammi, A. Bennani, A. El Ismaili, Y. Msatef, M.N. Serrhini                                                                                             |                |
| <b>Involvement of iron and ferritin in the potato–<i>Phytophthora infestans</i> interaction</b>                                                           | <b>557–562</b> |
| C.G. Mata, L. Lamattina, R.O. Cassia                                                                                                                      |                |
| <b>Sequence analysis shows that a dwarfing disease on rice, wheat and maize in China is caused by <i>Rice black-streaked dwarf virus</i></b>              | <b>563–567</b> |
| H. Zhang, J. Chen, J. Lei, M.J. Adams                                                                                                                     |                |
| <b>Book review</b>                                                                                                                                        | <b>569</b>     |

## Volume 107, No. 6 July 2001

|                                                                                                                                                                 |                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Light and scanning electron microscopy studies on the infection of oriental lily leaves by <i>Botrytis elliptica</i></b>                                     | <b>571–581</b> |
| T.F. Hsieh, J.W. Huang, T. Hsiang                                                                                                                               |                |
| <b>Investigation of genomic variability of <i>Xanthomonas arboricola</i> pv. <i>juglandis</i> by AFLP analysis</b>                                              | <b>583–591</b> |
| S. Loreti, A. Gallelli, A. Belisario, E. Wajnberg, L. Corazza                                                                                                   |                |
| <b>Ribosomal DNA sequence comparisons of <i>Colletotrichum graminicola</i> from turfgrasses and other hosts</b>                                                 | <b>593–599</b> |
| T. Hsiang, P.H. Goodwin                                                                                                                                         |                |
| <b>Evaluation of natural chemical compounds against root-lesion and root-knot nematodes and side-effects on the infectivity of arbuscular mycorrhizal fungi</b> | <b>601–605</b> |
| C. Calvet, J. Pinochet, A. Camprubí, V. Estaún, R. Rodríguez-Kábana                                                                                             |                |
| <b>Susceptibility of clematis varieties and species to stem infection by <i>Phoma clematidina</i> as an indicator for resistance to wilt</b>                    | <b>607–614</b> |
| P. van de Graaf, T.M. O'Neill, J.M. Chartier-Hollis, M.E. Joseph                                                                                                |                |
| <b>Epidemiology of grey mould in annual waiting-bed production of strawberry</b>                                                                                | <b>615–624</b> |
| P. Boff, P. Kastelein, J. de Kraker, M. Gerlagh, J. Köhl                                                                                                        |                |
| <b>Incidence of <i>Soil-borne wheat mosaic virus</i> in mixtures of susceptible and resistant wheat cultivars</b>                                               | <b>625–631</b> |
| D. Hariri, M. Fouchard, H. Prud'homme                                                                                                                           |                |

|                                                                                                                                                                |                |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Characterisation of a tobamovirus from trailing petunias</b>                                                                                                | <b>633–638</b> |
| N.J. Spence, I. Sealy, P.R. Mills, G.D. Foster                                                                                                                 |                |
| <b>RAPD characterization of single females of the root-knot nematodes, <i>Meloidogyne</i> spp.</b>                                                             | <b>639–643</b> |
| O. Randig, F. Leroy, M. Bongiovanni, P. Castagnone-Sereno                                                                                                      |                |
| <b>Papaya ringspot virus resistance of transgenic Rainbow and SunUp is affected by gene dosage, plant development, and coat protein homology</b>               | <b>645–653</b> |
| P. Tennant, G. Fermin, M.M. Fitch, R.M. Manshardt, J.L. Slightom, D. Gonsalves                                                                                 |                |
| <i>Short communications</i>                                                                                                                                    |                |
| <b>The <i>Cladosporium fulvum</i> Bap1 gene: evidence for a novel class of Yap-related transcription factors with ankyrin repeats in phytopathogenic fungi</b> | <b>655–659</b> |
| H.-J. Bussink, A. Clark, R. Oliver                                                                                                                             |                |
| <b>Molecular identification of <i>Oat mosaic virus</i> as a <i>Bymovirus</i></b>                                                                               | <b>661–666</b> |
| W.A. Monger, G.R.G. Clover, G.D. Foster                                                                                                                        |                |

## Volume 107, No. 7 September 2001

### *Mini review*

|                                                                                                                                                                                         |                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Pine wilt disease caused by the pine wood nematode: the induced resistance of pine trees by the avirulent isolates of nematode</b>                                                   | <b>667–675</b> |
| H. Kosaka, T. Aikawa, N. Ogura, K. Tabata, T. Kiyohara                                                                                                                                  |                |
| <b>Incidence of viruses infecting <i>Allium</i> spp. in Greece</b>                                                                                                                      | <b>677–684</b> |
| C.I. Dovas, E. Hatziloukas, R. Salomon, E. Barg, Y. Shibolet, N.I. Katis                                                                                                                |                |
| <b>Improving control of green and blue molds of oranges by combining <i>Pantoea agglomerans</i> (CPA-2) and sodium bicarbonate</b>                                                      | <b>685–694</b> |
| N. Teixidó, J. Usall, L. Palou, A. Asensio, C. Nunes, I. Viñas                                                                                                                          |                |
| <b>Transmission characteristics of the European stone fruit yellows phytoplasma and its vector <i>Cacopsylla pruni</i></b>                                                              | <b>695–700</b> |
| L. Carraro, N. Loi, P. Ermacora                                                                                                                                                         |                |
| <b>Characterization of a new begomovirus from Egypt infecting hollyhock (<i>Althea rosea</i>)</b>                                                                                       | <b>701–711</b> |
| L. Bigarré, M. Chazly, M. Salah, M. Ibrahim, M. Padidam, M. Nicole, M. Peterschmitt, C. Fauquet, J.C. Thouvenel                                                                         |                |
| <b>Complementation of CMV subgroup IA strains in replicase-mediated resistant tobacco plants after co-inoculation with different cucumoviruses</b>                                      | <b>713–721</b> |
| K.-H. Hellwald, D. Glenewinkel, S. Hauber, S. Wittlinger                                                                                                                                |                |
| <b>Possible mechanisms influencing the dynamics of rhizoctonia disease of tulips</b>                                                                                                    | <b>723–738</b> |
| J.H.M. Schneider, C.G. Kocks, M.T. Schilder                                                                                                                                             |                |
| <b>Repetitive sequence-derived PCR profiling using the BOX-A1R primer for rapid identification of the plant pathogen <i>Clavibacter michiganensis</i> subspecies <i>sepedonicus</i></b> | <b>739–748</b> |
| N.C. Smith, J. Hennessy, D.E. Stead                                                                                                                                                     |                |
| <b>Field response to <i>Pratylenchus thornei</i> of a wheat line with the <i>CreAet</i> gene for resistance to <i>Heterodera avenae</i></b>                                             | <b>749–755</b> |
| G. Nombela, M.D. Romero                                                                                                                                                                 |                |

*Short communications*

- An efficient microtiter system to determine *Agrobacterium* biovar** 757–760  
J. Cubero, M.M. López
- Occurrence of *Pythium ultimum* var. *ultimum* in a greenhouse on Spitsbergen Island, Svalbard** 761–765  
M. Tojo, T. Hoshino, M.L. Herrero, S.S. Klemsdal, A.M. Tronsmo

**Volume 107, No. 8 October 2001**

- Infection of *Nicotiana* species by the anthracnose fungus, *Colletotrichum orbiculare*** 767–773  
S. Shen, P.H. Goodwin, T. Hsiang
- Aubian wheat mosaic virus, a new soil-borne wheat virus emerging in France** 775–785  
D. Hariri, H. Prud'homme, M. Fouchard, G. Boury, P. Signoret, H. Lapierre
- Effect of bactericides, phosphonates and nutrient amendments on blast of dormant flower buds of pear: a field evaluation for disease control** 787–794  
E. Montesinos, P. Vilardell
- Leaf wax layer may prevent appressorium differentiation but does not influence orientation of the leaf rust fungus *Puccinia hordei* on *Hordeum chilense* leaves** 795–803  
M.C. Vaz Patto, R.E. Niks
- Repeated applications of *Penicillium oxalicum* prolongs biocontrol of fusarium wilt of tomato plants** 805–811  
*Biological control of fusarium wilt in tomato plants*  
A. De Cal, P. Melgarejo
- Surface characteristics of necrotrophic secondary hyphae produced by the bean anthracnose fungus, *Colletotrichum lindemuthianum*** 813–819  
S.E. Perfect, J.R. Green, R.J. O'Connell
- Application of a putative fatty-acid binding protein to discriminate serologically the two European quarantine root-knot nematodes, *Meloidogyne chitwoodi* and *M. fallax*, from other *Meloidogyne* species** 821–832  
C. Tastet, F. Val, M. Lesage, L. Renault, L. Marché, M. Bossis, D. Mugniéry

*Short communications*

- Molecular evidence that a lily-infecting strain of *Tulip breaking virus* from Japan is a strain of *Lily mottle virus*** 833–837  
Y. Yamaji, X. Lu, S. Kagiwada, K. Oshima, S. Namba
- Influence of amylovoran production on virulence of *Erwinia amylovora* and a different amylovoran structure in *E. amylovora* isolates from *Rubus*** 839–844  
M. Maes, K. Orye, S. Bobev, B. Devreese, J. Van Beeumen, A. De Bruyn, R. Busson, P. Herdewijn, K. Morreel, E. Messens
- Existence of two serological subclusters of *Plum pox virus*, strain M** 845–848  
A. Myrta, D. Boscia, O. Potere, M. Kölber, M. Németh, B. Di Terlizzi, M. Cambra, V. Savino

**Volume 107, No. 9 November 2001**

|                                                                                                                                                                                                               |                |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <b>Vegetative growth of sorghum and <i>Striga hermonthica</i> in response to nitrogen and the degree of host root infection</b>                                                                               | <b>849–860</b> |
| W. Sinebo and D.S.H. Drennan                                                                                                                                                                                  |                |
| <b>Phytogard® and DL-<math>\beta</math>-amino butyric acid (BABA) induce resistance to downy mildew (<i>Bremia lactucae</i>) in lettuce (<i>Lactuca sativa</i> L)</b>                                         | <b>861–869</b> |
| E. Pajot, D. Le Corre and D. Silué                                                                                                                                                                            |                |
| <b>Genetic diversity in relation to sexual and asexual reproduction in populations of <i>Melampsora larici-epitea</i></b>                                                                                     | <b>871–881</b> |
| B. Samils, V. Stepien, U. Lagercrantz, M. Lascoux and U. Gullberg                                                                                                                                             |                |
| <b>The effect of soil moisture and cabbage amendment on the thermoinactivation of <i>Phytophthora nicotianae</i></b>                                                                                          | <b>883–894</b> |
| L. Coelho, D.J. Mitchell and D.O. Chellemi                                                                                                                                                                    |                |
| <b>Production of indole-3-acetic acid and tryptophol by <i>Pythium ultimum</i> and <i>Pythium</i> group F: Possible role in pathogenesis</b>                                                                  | <b>895–904</b> |
| P. Rey, S. Leucart, H. Désilets, R.R. Bélanger, J.P. Larue and Y. Tirilly                                                                                                                                     |                |
| <b>PCR-based assays to assess wheat varietal resistance to blotch (<i>Septoria tritici</i> and <i>Stagonospora nodorum</i>) and rust (<i>Puccinia striiformis</i> and <i>Puccinia recondita</i>) diseases</b> | <b>905–917</b> |
| B.A. Fraaije, D.J. Lovell, J.M. Coelho, S. Baldwin and D.W. Hollomon                                                                                                                                          |                |
| <b>Specific behaviour of French <i>Aphanomyces euteiches</i> Drechs. populations for virulence and aggressiveness on pea, related to isolates from Europe, America and New Zealand</b>                        | <b>919–929</b> |
| E. Wicker and F. Rouxel                                                                                                                                                                                       |                |
| <i>Short communication</i>                                                                                                                                                                                    |                |
| <b>Rapid identification of <i>Agrobacterium</i> species by staircase electrophoresis of low molecular weight RNA profiles</b>                                                                                 | <b>931–938</b> |
| E. Velázquez, J.L. Palomo, B. Lastra, P.F. Mateos, P. García-Benavides and E. Martínez-Molina                                                                                                                 |                |
| <b>Volume contents</b>                                                                                                                                                                                        | <b>939–946</b> |
| <b>Author index</b>                                                                                                                                                                                           | <b>947–949</b> |