

Expert Opinion

1. Introduction
2. Promise and perils of cell-mediated drug delivery
3. Neurodegenerative disorders and cell-based carriage of therapeutic nanoparticles
4. Cancer
5. Lung injury
6. Microbial infections
7. Epilepsy
8. Expert opinion

informa
healthcare

Cell-mediated drug delivery

Elena V Batrakova[†], Howard E Gendelman & Alexander V Kabanov

[†]University of Nebraska Medical Center, Center for Drug Delivery and Nanomedicine, Department of Pharmaceutical Sciences, Omaha, NE, USA

Introduction: Drug targeting to sites of tissue injury, tumor or infection with limited toxicity is the goal for successful pharmaceuticals. Immunocytes (including mononuclear phagocytes (dendritic cells, monocytes and macrophages), neutrophils and lymphocytes) are highly mobile; they can migrate across impermeable barriers and release their drug cargo at sites of infection or tissue injury. Thus, immune cells can be exploited as Trojan horses for drug delivery.

Areas covered: This paper reviews how immunocytes laden with drugs can cross the blood-brain or blood-tumor barriers to facilitate treatments for infectious diseases, injury, cancer, or inflammatory diseases. The promises and perils of cell-mediated drug delivery are reviewed, with examples of how immunocytes can be harnessed to improve therapeutic end points.

Expert opinion: Using cells as delivery vehicles enables targeted drug transport and prolonged circulation times, along with reductions in cell and tissue toxicities. Such systems for drug carriage and targeted release represent a new disease-combating strategy being applied to a spectrum of human disorders. The design of nanocarriers for cell-mediated drug delivery may differ from those used for conventional drug delivery systems; nevertheless, engaging different defense mechanisms in drug delivery may open new perspectives for the active delivery of drugs.

Keywords: cell carriers, drug delivery, immunocytes, nanoparticles, targeted drug transport

Expert Opin. Drug Deliv. (2011) 8(4):415-433

1. Introduction

The development of targeted drug delivery is among the most important goals of pharmaceutical research. Its realization can lead to improved therapeutic efficacy, reductions in drug dosing intervals and decreased toxicities. However, this is not a simple task as drug homing to pathologically relevant disease sites has only recently been investigated. Obstacles are substantial and include sustained time-based plasma concentrations and local blood flow.

2. Promise and perils of cell-mediated drug delivery

Using immunocytes, mononuclear phagocytes (monocytes, macrophages and dendritic cells) lymphocytes, and neutrophils and stem cells as drug carrier systems offers several advantages over common drug administration regimens. These include: targeted drug transport to disease sites; prolonged drug half-lives; time-controlled drug release; and diminished drug immunogenicity and cytotoxicity profiles. Immunocytes and stem cells show an intrinsic homing property, enabling them to migrate to sites of injury, inflammation and tumor. In addition, they can act as Trojan horses, carrying concealed drug cargoes while migrating across impermeable barriers (e.g., blood-brain or blood-tumor barriers) to sites of disease.

Despite such advantages, there is as yet limited success, for several reasons. First, drug loading in cell carriers is low. Second, there are limitations owing to the ability of immunocytes and stem cells efficiently to disintegrate and clear entrapped

Article highlights.

- Using cell-mediated drug delivery systems offers several advantages, including: targeted drug transport to disease sites; prolonged drug half-lives; time-controlled release of loaded drugs; and diminished drug immunogenicity and cytotoxicity profiles.
- The goals that need to be achieved for successful cell-mediated drug delivery include: high drug loading into cell carriers; efficient preservation of entrapped therapeutic agents against disintegration and clearance in the host cells; drug-triggered release at the site of action; efficient homing of cell carriers to a disease site; and safety of cell-based drug formulations for the whole organism.
- Two different approaches are utilized in cell-mediated drug delivery: genetically modified cell carriers producing therapeutically active molecules; and drug-loaded cell carriers used as 'Trojan horses' to deliver the drug to the disease site.
- Living cells for drug carriage and release represent a new disease-combating strategy that can be applied to a spectrum of human infectious, cancerous and degenerative disorders.

This box summarizes key points contained in the article.

therapeutic agents. Third, the loaded drug should not be prematurely released, but unloaded, in continuous action, on the cell's arrival to the site of action or disease. Fourth, the cell carriers should migrate to the disease site in substantial quantities. This should not be compromised during drug loading. Finally, all used formulations must be safe for both the cell carrier and the organism.

Many of these limitations can be addressed by incorporation of drugs into protective polymeric nanocarriers (liposomes and lipid nanoparticles [1-4], micelles [5-8], nanogels [9,10], nanospheres and nanocapsules [11,12], solid nanoparticles and nanosuspensions [13-16], block ionomer complexes [17,18] or nanofibers and nanotubes [19,20]) that preserve drugs inside sub-cellular organelles (Figure 1). The ideal drug carrier for cell-mediated delivery should have: optimal size, shape and surface characteristics; multivalent attachment; controlled drug release; and biocompatibility. All these characteristics are essential for translating cell-mediated drug delivery systems for human use.

2.1 Drug loading into cell carriers

Nanocarriers commonly have a core-shell structure. Their central part (such as aqueous pool of nanospheres, hydrophobic core of micelles, or polyelectrolyte core of nanogels) permits drug entrapment. It is surrounded with a polymer shell (or lipids in liposomes and lipid nanoparticles), which defines the nanocarrier dispersion stability, circulation time and cell interactions [1-8,10,11,13,14,17,18]. Both the core and shell are crucial for successful cell-mediated delivery. For example, the surface coating affects the ability of particles to internalize into the cells and therefore affects the loading of cell carriers.

In general, charged nanocarriers are rapidly taken up by mononuclear phagocytes (MP) and other immunocytes or stem cells compared with neutral nanoparticles [1,21-25]. The recognition occurs by receptors located on the cell plasma membranes, in particular mannose, complement and Fc receptors (MR, CR and FcR) [26-28]. In general, MR recognize mannans, as well as integrins (e.g., CD11b/CD18), CR interact with particles after nonspecific complement opsonization, and FcR recognize particles after specific antibody opsonization [28]. Furthermore, positively charged nanoparticles accumulate in MP to a greater extent than negatively charged particles. Thus, positively charged nanoparticles prepared by high-pressure homogenization with antiretroviral drugs indinavir (IDV), zidovudine (ZDV) and efavirenz (EFV) accumulate in MP about twofold more than the same size negatively charged carriers [14]. A similar effect was reported for nanoparticles ('nanozymes') comprising redox enzyme, catalase and variant synthetic polyelectrolyte block copolymers [29]. Loading capacity of nanozymes comprising positively charged mono-polymers (polyethyleneimine [PEI]- and poly-L-lysine [PL]-based) is greater than the nanozyme prepared with negatively charged block poly-L-glutamic acid (PGLU)-based copolymer. This probably occurs owing to the greater absorption and internalization of positively charged nanoparticles to negatively charged outside plasma membrane of cell carriers.

Electrostatically neutral and hydrophilic poly(ethylene glycol) (PEG) is perhaps the most common shell-forming polymer now used in injectable nanocarriers [30-32]. In particular, commercial PEGylated liposome-encapsulated Dox, Doxil, is approved for use in the treatment of recurrent ovarian cancer, AIDS-related Kaposi's sarcoma [33] and metastatic breast cancer [34]. However, PEG is not the only polymer that should be considered in the context of the cell-mediated delivery, as it can limit cellular uptake, as demonstrated for microspheres bearing high-density surface PEG chains that were resistant to phagocytosis [35]. Similar results were obtained with nanoparticles consisting of catalase and positively charged polymers based on PEI [29]. Thus, loading capacity of nanozymes without PEG corona in MP was significantly greater than those with PEG. To this end, modification of nanocarrier shell with specific vector moieties can provide for targeted cell delivery of nanocarriers [1,26,36-38]. In particular, modification of the PEG shell of a poly(amidoamine) nanoparticle (PAMAM) with streptavidin facilitates targeting to biotinylated T-cell markers such as anti-CD3 or peptide/major histocompatibility (MHC) complexes (Figure 2) [39].

Size and shape of nanocarriers are also of importance for cell uptake, although the phagosome may have different sizes depending on the size of the particles, which can range from as little as a few hundred nanometers [35,40,41] to several micrometers [42]. For example, murine bone marrow-derived macrophages accumulate IgG-opsonized latex beads > 20 µm in diameter [43]. Furthermore, particles with size ~ 1 µm were accumulated at a greater extent (2.56-fold) than smaller drug carriers at 500 µm [14].




Figure 1. Types of nanocarrier for drug delivery. A. Liposomes. B. Polymer and lipid nanoparticles. C. Nanospheres and nanocapsules. D. Nanosuspensions. E. Polymer micelles. F. Nanogels. G. Block ionomer complexes. H. Nanofibers and nanotubes.

Reproduced with permission from [110].

A recent study by Champion *et al.* reported striking effects of shape of particles on phagocytosis in alveolar rat macrophages [44]. Polystyrene-based particles of > 20 shapes including spheres, rectangles, rods, worms, oblate ellipses, elliptical disks and UFO-like particles were manufactured. The local particle shape at the point where it was attached to the cell played a crucial role in phagocytosis. For example, a macrophage attached to a sharper side of the ellipse would internalize the particle in a few minutes. By contrast, a macrophage attached to a dull side would not internalize the same ellipse for hours. Although particle size played a reduced role in the initiation of the phagocytosis, it could of course affect its completion, especially when the particle volume exceeded that of a cell.

2.2 Drug preservation inside cells

The stability of the drug loaded into cell carriers can depend, among other factors, on the intracellular trafficking of the nanocarriers. In general, drug-loaded nanocarriers need to avoid lysosomes to reduce drug disintegration inside cells [14]. In this regard, it is noteworthy that cationic and anionic nanoparticles show divergent fates inside MP [38]. For example, phagocytosis of cationic polyamine-coated nanoparticles leads to diminished phagosomal acidification when compared with anionic protein-coated particles. Such cationic nanoparticles protect the incorporated drug against lysosomal degradation [29,38]. Furthermore, loading of 'nanozymes' containing positively charged block copolymers (PEI-PEG and PL-PEG) protected the enzyme in macrophages. Increasing the amount




Figure 2. Schematic representation of streptavidin-PEG^{FITC}/PAMAM construction. **A.** The construction of streptavidin-PEG^{FITC}/PAMAM. **B.** Absorbance profiles of streptavidin-PEG^{FITC}/PAMAM and PEG^{FITC}/PAMAM. **C.** Reverse-phase HPLC of streptavidin-PEG^{FITC}/PAMAM and PEG^{FITC}/PAMAM.

Reproduced with permission from [39].

of block copolymer in the nanozyme formulation improved the stability of the enzyme. By contrast, catalase loaded in a polyion complex with a negatively charged block copolymer (PGLU-PEG) was degraded in macrophages to an even greater extent than catalase loaded alone. Protection of the enzymatic activity inside carrier cells may be the result, in part, of a 'proton sponge' effect of block copolymers [45] when an excess of amino groups on the surface of the nanoparticles buffers acidification of the cell's endocytic compartments. This serves to inhibit protease activity and decrease drug degradation [38]. In other words, a positively charged block copolymer prevents phagosome-lysosomal clearance functions and, as a result, enzyme degradation.

In extreme case, intracellular drug degradation can be avoided by attaching the drug to the surface of cell carriers. This 'back pack' approach would still provide targeted transport and increased blood circulation along with the preserved drug activity. Thus, attaching avidin-coated nanoparticles to the biotinylated plasma membrane can be achieved through the avidin-biotin complex [46]. Another possibility is red blood cells (RBCs) with attached drugs on their surface [47]. Specifically, glycoprotein A covalently conjugated to the surface of the RBCs may provide extended half-life, controlled volume of distribution and multivalent therapeutic interactions. However, general limitations of the 'back pack' approach may include decreased loading of cell carriers,

impeded drug release at the disease site, as well as increased immunogenicity and toxicity.

2.3 Drug release from cell carriers

The mechanism of nanoparticle unloading at the site of action remains an active area of investigation. Controlled release of drugs from the cell carriers modulates the rate of drug appearance, dose and duration of exposure at the diseased sites. To this end, utilizing cellular responses to various conditions could provide the desired triggered release. Obviously, targeting of cell carriers and their prolonged residence at the disease site should provide opportunities for drug unloading. In addition, mononuclear phagocytes, in particular, are known to produce and store various compounds in intracellular vesicles, and liberate them by means of exocytosis at the site of the disease. A similar mechanism can trigger release of drug-loaded nanocarriers when macrophages serve as drug delivery vehicles [48]. Furthermore, the drug release can also be triggered by the increase of intracellular concentration of Ca^{2+} [49]. Finally, mild hypothermia was shown to facilitate controlled release of drug-loaded liposomes from macrophages in anticancer therapy [50].

Overall the structure and composition of protective nanocontainers play a crucial role in the effectiveness of formulations for cell-based drug delivery systems. For example, recent studies indicate that the structure of the block copolymer used for catalase nanoformulation, nanozyme, affects its cytotoxicity, loading and release capacities, as well as preservation of catalase enzymatic activity inside cell carriers [29]. Thus, nanozymes containing a negatively charged block copolymer (PGLU-PEG/catalase) demonstrated low toxicity, high loading capacity and effective release from macrophages. However, the polymer provided limited protection of the enzyme against cell-associated protease degradation. By contrast, nanozymes based on positively charged block copolymers, especially the PLs, showed increased cytotoxicity and low loading and release rates, but were protective of the catalase. Increasing the amount of positively charged block copolymer in nanozyme led to protection of catalase enzymatic activity but substantially reduced loadings and release. Importantly, nanozymes with PEG corona show water stability, limited cytotoxicity and efficient catalase protection. Nonetheless, these formulations also demonstrate decreased loading capacity and release rates. In addition, nanozymes based on a mono-polymer (without PEG corona) have higher loading and release levels, but did not protect catalase inside the cell carriers. Taken together, the optimal nanozyme formulation was the one based on positively charged block copolymers (PEI-PEG/catalase and PL-PEG/catalase), which demonstrate the most efficient protection of catalase enzymatic activity along with relatively high loading and release rates with limited cytotoxicity (Figure 3).

2.4 Homing of drug-loaded cell carriers at disease sites

The numbers of cell carriers that can reach the disease site is especially crucial in the case of CNS disorders when

drug-loaded cells need to penetrate the blood-brain barrier (BBB) to mediate therapeutic effect. Many neurological diseases, such as Alzheimer's and Parkinson's diseases, Prion disease, meningitis, encephalitis and HIV-associated neurocognitive disorders (HAND), have in common an inflammatory component [51]. The process of inflammation is characterized by extensive MP recruitment. Notably, mononuclear phagocytes migrate towards the inflammation site by means of the processes known as diapedesis and chemotaxis [52]. Such cells efficiently cross the BBB owing to their margination and extravasation properties causing barrier breakdown as a consequence of brain inflammation [53-57]. Many reports in the literature indicate that blood-borne monocytes traffic primarily between adjacent endothelial cells, that is, paracellularly through the junctional complexes [58,59]. Therefore, these cells can be loaded with a required drug and administered intravenously to reach the brain. For example, in an experimental model of Parkinson's disease, considerable levels of catalase loaded into bone marrow-derived macrophages (BMM) were reported in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-intoxicated brains [48]. Approximately 2.1% of the injected catalase dose was delivered to the brain during MPTP-associated inflammation. Furthermore, several studies confirmed the migration of inflammatory-response cells, in particular bone marrow-derived mesenchymal stromal cells towards injury sites such as infarcted myocardium [60,61], spinal cord injury [62] and cerebral ischemia [63].

Neural stem cells (NSCs) were also suggested as drug delivery vehicles for gene therapy in the CNS [64-66]. Indeed, these cells are highly migratory and migrate to areas of brain pathology, including ischemic and neoplastic brain lesions that are commonly present in Alzheimer's disease, Parkinson's disease, brain cancer, stroke and multiple sclerosis. How the mobility of stem cells is directed is not understood, although NSCs express a wide variety of receptors that should enable them to respond to many chemotactic signals present in brain pathologies (Figure 4) [66]. In particular, activated microglia induced NSC migration to the brain. Delivery of various neurotropic factors is in the focus of cell-mediated strategies for neurological diseases treatment.

With regard to cell-mediated drug delivery of antineoplastic agents, mononuclear phagocytes are known to accumulate in large numbers in vascular, hypoxic sites in cancer tumors, for example, in breast and prostate carcinomas [67,68]. Hypoxia is widespread in malignant human tumors owing to their poorly organized vasculature. The cytokines released by tumor cells in response to hypoxia and other physiological stresses usually attract macrophages and monocytes, which should facilitate anticancer drug delivery using these cells as vehicles, avoiding indiscriminate drug distribution and decreasing severe toxicity.

Finally, targeting to disease sites can be improved by drug-incorporated magnetic nanoparticles loaded into the cell carriers and followed by application of local magnetic fields [37]. Thus, albino rats with brain inflammation (induced by




Figure 3. Optimization of nanocarriers' structure for catalase delivery in macrophages. Nanozymes consisting of catalase and a negatively charged block copolymer demonstrate low toxicity, high loading capacity and release from bone marrow-derived macrophages; however, they provide limited protection of the enzyme against protease degradation inside the cell carriers. Nanozymes based on positively charged block copolymers, especially the poly-L-lysines, showed increased cytotoxicity and low loading and release rates, but were highly protective of catalase. Furthermore, nanozymes with PEG corona show good stability in water, limited cytotoxicity and efficient protection of catalase, but decreased loading capacity and cell release. The optimal nanozyme formulation is one based on positively charged block copolymers (PEI-PEG/catalase and PL-PEG/catalase).

Reproduced with permission from [29].

intrastriatal microinjection of human recombinant IL-1 β) received intravenous injections of RGD-coated magnetic liposomes and non-magnetic liposomes as a control. RGD peptide (i.e., small peptide domain Arg-Gly-Asp) was used for selective binding by monocytes and neutrophils that express integrin receptors on their surface. Magnetic liposomes demonstrated about a 10-fold increase in brain levels compared with non-magnetic control carriers when a local magnetic field was applied. In addition, magnetic neutrophils prepared *in vitro* target lungs under magnetic guidance following intravenous injection [69]. Moreover, drug-loaded magnetic liposomes can be targeted for selective and preferential presentation to blood monocytes/neutrophils, which results in both drug and magnetic incorporation into these cells and can be guided to target tissue sites.

2.5 Safety of cell-mediated drug delivery systems

An obvious concern for inflammatory-response cell-based drug delivery relates to possible cytotoxic effects of cell carriers. Mononuclear phagocytes attracted to the site of pathology by cytokines release reactive oxygen species (ROS) that cause cell damage. Moreover, several therapeutic strategies for CNS neurodegenerative disorders are based on the prevention of

monocyte-macrophage infiltration [70]. Therefore, precluding cytotoxicities for cell-based drug formulations is a requirement for formulation developments. Furthermore, genetically modified and immortalized cell carriers show atypical characteristics, such as higher degree of multipotency, which may increase the probability of tumor formation. Nevertheless, studies so far report no cytotoxic effects after macrophage-mediated drug delivery in the brain of healthy C57BL/6 mice adoptively transferred with macrophages carrying nanoformulated catalase [71]. In addition, propagation and expansion of immortalized and genetically modified cell carriers should allow improved quality controls [66].

2.6 Cell-mediated drug delivery systems in clinical settings

Finally, developing such methods in a clinical setting may be successful when cell carriers are harvested from peripheral blood by apheresis, then loaded with particles and re-infused into the patient. An alternative approach may be harvesting stem cells from bone marrow, propagating them in culture to obtain monocytes, and then loading the cells with a drug and adoptively transferring them. This procedure will allow for expansion of the cell population, although this would




Figure 4. Using neural stem cells for drug delivery to the CNS. Neural stem cells are attracted by at least three physiological processes that are common to many brain pathologies: inflammation, reactive astrocytosis and angiogenesis. Pathology-induced CNS inflammation is mediated by activated microglia that release cytokines and chemokines, which, in turn, increases the inflammatory reaction. The brain lesion and subsequent inflammation trigger reactive astrocytosis. The lesion-induced angiogenesis and inflammation-activated endothelial cells enhance neural stem cell homing to brain pathology by secreting chemoattractant factors, and also offer an atypical, perivascular niche for support of immigrating neural stem cells. Reproduced with permission from [66].

require a more invasive procedure. To this end, immortalization of cell carriers can allow propagation of cells with definable properties so that clonal populations with particular traits can be established. Furthermore, drug loading into cell carriers can be achieved directly in the patient's peripheral blood; nano-formulated drugs would be injected and selectively taken up by circulating cell carriers. Targeting of such nanocarriers could be achieved by coating the nanocarrier surface with the receptor-specific moieties (e.g., folate, gelatin, fibronectin, A-protein, mannose, or RGD peptide), which is recognized by specific MP surface receptors. Overall, chronic diseases are more

suitable for this type of drug delivery than pathologies that require acute intervention. Based on such considerations, some successful investigations using living cells as drug delivery vehicles for therapeutics are now presented (Table 1).

3. Neurodegenerative disorders and cell-based carriage of therapeutic nanoparticles

3.1 Alzheimer's disease and Parkinson's disease

The progressive impairment of short-term memory and emotional disturbances that typify Alzheimer's disease results

Table 1. Examples of cell-mediated drug delivery systems.

Type of cell-carrier	Therapeutic agent	Route of administration	Ref.
<i>Alzheimer's disease</i>			
NSCs	Ciliary neurotrophic factor	Intracranial injection	[73]
NSCs	Nerve growth factor	Intracranial injection	[65]
NSCs	Brain-derived neurotrophic factor	Intracranial injection	[64,75]
NSCs	Choline acetyltransferase	Intracranial injection	[72,74]
<i>Parkinson's disease</i>			
Monocytes, macrophages	Catalase	Intravenous injection	[29,48,71]
Fetal dopamine neurons	Dopamine	Intracranial injection	[76]
NSCs	Glial cell line-derived neurotrophic factor	Intracranial injection	[77,80]
NSCs	Vascular endothelial growth factor	Intracranial injection	[78,79]
<i>HIV-1 infection and neuroAIDS</i>			
Monocytes, macrophages	Atazanavir, ritonavir, indinavir	Intravenous injection	[14,81-83]
<i>Cancer</i>			
T cells	Doxorubicin	<i>In vitro</i>	[94]
MSCs	TNF-related apoptosis-inducing ligand	<i>Ipsilateral</i>	[93]
MSCs	Interferon- β	Intravenous injection	[86,90,95]
MSCs	Oncolytic adenovirus	Intravenous injection	[91,92]
MSCs	Antineoplastic agent	<i>In vitro</i> , in three-dimensional collagen gels	[46]
Erythrocytes	5-Fluorouracil	Intravenous injection	[96]
Monocytes and macrophages	Au-containing nanoparticles	<i>In vitro</i> , in spheroids	[85,89]
Intraperitoneal macrophages	5-Fluorouracil	Intraperitoneal injection	[50]
Bacterially derived minicells	Small interfering RNA, cytotoxic drugs	Intravenous, intratumoral injections	[97,98]
Bacterially derived minicells	Doxorubicin	Intravenous, intratumoral injections	[98]
Macrophages	Imaging agents, quantum dots	Intravenous injection	[103]
Macrophages	<i>N</i> -(2-hydroxypropyl)methacrylamide copolymer-gadolinium chelates	Intravenous injection	[104]
Hybridoma cell lines 2A and OX86	Anti-CD137 and anti-OX40 mAb	Subcutaneous injection	[102]
<i>Inflammation</i>			
Macrophages	Betamethasone phosphate	<i>In situ</i>	[25]
Erythrocytes	Glycoprotein A		[25]
Sertoli cells	Anti-inflammatory agent, curcumin	Intravenous injection	[105]
<i>Microbial infections</i>			
Erythrocytes	Antifungal agent, amphotericin B	<i>In vitro</i>	[106]
Macrophages	Chloroquine	Intravenous injection	[107]
<i>Epilepsy</i>			
Human MSCs and human embryonic stem cells	Adenosine	Intracranial injection in cell-encapsulation device	[108,109]

MSCs: Mesenchymal stem cells; NSCs: Neural stem cells.

from synaptic dysfunction and neuronal death in the hippocampus and linked regions in the cerebral cortex and limbic system; because of the increasing numbers of affected people, there is an urgent need to develop strategies able to interfere with disease progression and protect neurons. In this respect,

delivery of neurotrophic factors (NTFs) including nerve growth factor (NGF) [65], brain-derived neurotrophic factor (BDNF) [64] and choline acetyltransferase (ChAT) [72] is of urgent need. To overcome limited drug transport across the BBB, direct brain implantation of cells engineered to produce

NTFs was developed [73]. In particular, ciliary neurotrophic factor delivered in cells implanted in the brain prevents A β oligomer-induced neuronal damage and neurobehavioral impairments in mouse models of Alzheimer's disease. In another study, NSCs overexpressing ChAT cDNA targeted acetylcholine deficits [74]. The ability of cell carriers, placed by cortical transplants, to improve cognition was tested after induced cholinergic lesions in rats. Significant improvements were recorded in water maze acquisition as well as in the retention and spatial probe trials [74]. A robust enhancement of hippocampal synaptic density, mediated by BDNF and delivered by NSCs, was seen [75]. NSCs ameliorated complex behavioral deficits associated with Alzheimer's disease pathology by BDNF. In these experiments, intracranial transplantation of the cell carriers was utilized.

Owing to specific nigrostriatal degeneration, Parkinson's disease was one of the first targets for cell therapy transplants. Fetal dopamine neurons were inoculated into the putamen, where the cells worked as dopamine pumps, as seen during systemic administration of L-DOPA [76]. It was suggested that the transplanted cells were gene therapy vehicles for dopamine delivery rather than replacement neurons [76]. Delivery of neurotrophic factors such as glial cell line-derived neurotrophic factor (GDNF) or vascular endothelial growth factor (VEGF) in neural stem cell transplants was tested in Parkinson's disease mouse models [77-79]. GDNF was delivered in bone marrow-derived macrophages to the affected brain [80]. The cells were transduced *ex vivo* with lentivirus expressing a *GDNF* gene driven by a synthetic macrophage-specific promoter and then transplanted into recipient mice. Eight weeks after transplantation, the mice were injected with MPTP for 7 days to induce nigrostriatal neurodegeneration. Macrophage GDNF treatments dramatically ameliorated MPTP-induced degeneration of tyrosine hydroxylase (TH)-positive neurons seen in the substantia nigra and TH(+) terminals in the striatum. This resulted in axon regeneration and reversed hypoactivity in the open field test.

The development of new CNS drug delivery using macrophage carriers for delivery of the antioxidant enzyme catalase in MPTP mice was reported by the authors' laboratories [48,71]. In this system, nanoformulated catalase (nanozyme) was obtained by coupling the enzyme to a cationic block copolymer PEI-PEG, leading to a polyion complex micelle. BMM carried significant amounts of catalase then slowly released the active enzyme over several days [48]. The enzyme released on stimulation of nanozyme-loaded cell carriers decomposed microglial hydrogen peroxide produced on nitrated alpha-synuclein (N- α -syn) or TNF- α -induced activation *in vitro*. Subsequent studies examined the relationships between the composition of catalase nanozyme, their physicochemical characteristics (morphology, size and Zeta potential), and cell loadings, release and enzymatic activities for macrophage carriage [29].

A significant amount of catalase was detected in brains of mice after transfer of BMM loaded with nanoformulated catalase following MPTP intoxication. It was demonstrated that such nanozyme-loaded BMM injected into MPTP-intoxicated mice reduce neuroinflammation and attenuate nigrostriatal degeneration [71]. In particular, MPTP intoxication decreased the number of TH-positive nigral dopaminergic neurons (32% survival) compared with saline-treated controls (Figure 5). By contrast, the number of surviving dopaminergic neurons in MPTP-intoxicated mice treated with nanozyme-loaded BMM was greater than the total number of neurons in MPTP-treated mice (62.4% survival). Furthermore, treatment with nanozyme alone (without cell carriers) also produced some neuroprotection effect, although with fewer neurons (41.3% survival) compared with the mice treated with cell/nanozyme formulation. Finally, treatment with empty monocytes did not preserve neurons in MPTP-intoxicated animals (31.1% survival). This signified a neuroprotective effect of nanozyme-loaded monocytes in MPTP-induced neurodegeneration.

The possible means of BMM-mediated therapeutic effects of the nanozymes have been suggested (Figure 6): i) nanozyme-loaded BMM cross the BBB and release catalase at the site of inflammation (particularly in the substantia nigra pars compacta [SNpc]); ii) a 'depot' was established such that nanozyme is slowly released from BMM to the bloodstream and bypasses the BBB independently of cell carriers; and iii) catalase nanozyme released from BMM in the liver and spleen suppresses peripheral leukocyte activation, which results in significant protection of SNpc neurons against MPTP-induced neurodegeneration [71]. Overall, as few therapeutic modalities exist that affect progression of Parkinson's disease and are aimed at neuroregenerative therapies, the cell-mediated delivery of catalase and GDNF may represent efficacious strategies that attenuate neuroinflammation and provide neuroprotection for disease.

3.2 HIV-1 infections and neuroAIDS

NeuroAIDS is a clinical disorder that encompasses neurologic disorders seen primarily as a consequence of damage to the CNS by HIV. The clinical syndromes include cognitive, motor and behavioral disorders of varying severity. Such syndromes affect 30 – 40% of adults and children with AIDS and, despite the advent of potent combination antiretroviral therapy (cART), incidence rates remain constant although disease severity is less. Mononuclear phagocyte carriage has been utilized as a delivery system for antiretroviral therapy of HIV in mice [14,57,81-83]. To protect the drug against degradation, as well as the cell carriers against the drug, it was incorporated into nanosized drug crystals (termed nano-antiretroviral therapy, or nanoART, Figure 7). NanoART was made by high-pressure homogenization of crystalline drug with various poloxamer and/or phospholipid surfactants, or by wet-milling homogenization, which




Figure 5. Neuroprotective effect of nanozyme-loaded bone marrow-derived macrophages in an MPTP model of Parkinson's disease. MPTP-intoxicated C57Bl/6 mice (18 mg/kg) were intravenously injected with saline (second bar), nanozyme alone (third bar), macrophages loaded with nanozyme (5×10^6 cells/(mouse 100 μ l)) (fourth bar), or empty macrophages (fifth bar). **A.** Healthy non-intoxicated animals were used as control group (first bar). Seven days after treatment the animals were killed; brain slices were stained for tyrosine hydroxylase-positive nigral dopaminergic neurons. **B.** Results from $N = 5$ animals per group demonstrate a significant loss of nigrostriatal neurons in MPTP-treated mice, which is prevented by adoptive transfer of macrophages loaded with nanozyme. No significant neuroprotective effect was detected after treatment with nanozyme alone, or empty macrophages. Values are means \pm s.e.m. and $p < 0.05$ compared with (a) saline, (b) MPTP and (c) MPTP + macrophage/nanozyme.

Reproduced with permission from [71].

BMM: Bone marrow-derived macrophages; MPTP: 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine.




Figure 6. A pictorial scheme for cell-based nanoformulated drug delivery. Three possible ways of BMM-mediated therapeutic effects of catalase nanozyme in Parkinson's disease mouse model: pathway I, BMM loaded with nanozyme cross the BBB and release catalase in the substantia nigra pars compacta; pathway II, nanozyme is released from BMM to the bloodstream and bypasses the BBB independently of cell carriers; and pathway III, catalase nanozyme released from BMM in the liver and spleen suppresses peripheral leukocyte activation, which results in significant protection of nigrostriatal neurons against MPTP-induced neurodegeneration.

Reproduced with permission from [71].

BBB: Blood-brain barrier; BMM: Bone marrow-derived macrophages; MPTP: 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine.

produced nanoformulations with high loading capacities (> 80%) and relatively small particle sizes (200 – 350 nm). Scanning electron microscopy analysis revealed particles of various sizes and morphologies, which were polygonal, rod, cuboidal, or spherical in shape with smooth surfaces.

Simultaneous uptake of nanoformulated cART into the cell carriers was shown [83]. It was also demonstrated that size, charge, coating and shape of the nanoparticles are crucial for efficient MP uptake, drug release and cell migration [14]. Coating of nanoART greatly affected drug accumulation in cell carriers. Positively charged nanoART preparations are taken up better than negatively charged ones. To determine whether any of the wet-milled nanoformulations could induce long-term antiretroviral effects, macrophages were pretreated with individual nanoART preparations and then challenged with HIV-1_{ADA} up to 15 days after drug treatment. Nanoparticles loaded with atazanavir, ritonavir and indinavir were released from macrophages and demonstrated > 85, 80 and 40% inhibition, respectively, of progeny virion production and HIV-1p24 antigen to challenge day 15. All formulations

of EFV showed almost complete suppression of viral infection [83]. These results supported the continued development of macrophage-mediated nanoART carriage for treating HIV-1 disease. In addition, delivery of antisense oligodeoxynucleotides, ribozymes, incorporated into liposomes to HIV-1-infected cells was reported in human monocyte-derived macrophages [84]. Thus, in addition to nanoART a ribozyme complementary to HIV-1 5'-long terminal repeat delivered in pH-sensitive liposomes inhibited virus production by 90%, whereas the free ribozyme caused limited viral inhibitions.

4. Cancer

Cell-mediated delivery is also a potential therapeutic and diagnostic strategy for cancer [46,50,85-93]. The advantage herein is that cytotoxic activity of the cell carriers can be used in tandem with site-specific delivery of antineoplastic agents. In addition, antineoplastic agents are safely packaged into cell carriers to reduce secondary cytotoxicities. However,




Figure 7. NanoART. A. NanoART preparations of antiretroviral agents were prepared by high-pressure homogenization using an AvestinC-5 homogenizer. B. Scanning electron microscopy demonstrated different nanoparticle morphology. C. Uptake of drug-loaded nanoART (red) in monocyte-derived macrophages (green). D. Cells treated with nanoART showed complete or near complete suppression of HIV1 p24 antigen production. E. Biodistribution of nanoART in mice following intravenous injections. F. Patient treatment with nanoART in clinical settings.

Reproduced with permission from [111].

adverse effects of the cytotoxic drugs on the cell carriers themselves need to be considered. For example, when nanoparticles coated with cytotoxic antibiotic doxorubicin were loaded into T cells, the release of drug inside the cell carrier caused cell damage [94]. As a result > 60% T cells loaded with the nanoparticles were dead over 15 h after the loading.

To overcome these limitations, mesenchymal stem cells (MSCs) were genetically engineered to produce antitumor proteins [86,90-93,95]. MSCs possess a set of several unique properties, which make them ideally suited for both cellular therapies in regenerative medicine and as vehicles for gene and drug delivery to treat cancer. These properties include: i) relative ease of isolation; ii) the ability to differentiate into a wide variety of functional cell types of both mesenchymal

and non-mesenchymal origin; iii) the ability to be expanded extensively in culture without loss of differentiative capacity; iv) the hypoimmunogenicity and even ability to produce immunosuppression on transplantation; v) the pronounced anti-inflammatory properties; and vi) the ability to home to damaged tissues, tumors and metastases following *in vivo* administration. Thus, MSCs of human origin were genetically modified to express TNF-related apoptosis-inducing ligand (TRAIL) at the glioma tumor site. First, the retention of tumor tropic ability of hMSC S-TRAIL cells by *in vitro* and *in vivo* migration assays was clearly demonstrated [93]. Next, for the *in vivo* assessment of therapeutic efficacy, hMSCs were injected ipsilateral to an established intracranial glioma tumor in a mouse xenograft model. Genetically engineered

hMSC S-TRAIL cells were effective at inhibiting intracranial U87 glioma tumor growth (81.6%) *in vivo* and resulted in significantly longer animal survival. Immunohistochemical studies demonstrated a significant, eightfold greater tumor cell apoptosis in the hMSC S-TRAIL-treated group compared with controls. Overall, the study demonstrated the therapeutic efficacy of hMSC S-TRAIL cells and confirmed that hMSCs can serve as a powerful cell-based delivery vehicle for the site-specific release of therapeutic proteins. Furthermore, reduction of tumor growth by genetically modified hMSCs expressing interferon- β was reported in models of metastatic breast cancer, melanoma [86,90] and gliomas [95]. The hMSC were also used as carriers for conditionally replicating adenoviruses in a breast cancer metastasis model [91] and in a model of intracranial malignant glioma [92].

Attachment of nanoparticles with cytotoxic agent to the cell surface allowed cell carriers to remain intact. Thus, NeutrAvidin-coated nanoparticles were anchored on biotinylated plasma membrane of hMSC, where they remained attached for up to 2 days [46]. The hMSC with such nanoparticulate patches retained their inherent tumorigenic properties in a tumor model with a three-dimensional extracellular matrix. These results provide a new strategy actively to deliver nanostructures and therapeutics to tumors utilizing stem cells as carriers.

Besides MSCs, erythrocytes, mononuclear phagocytes and bacterially derived minicells were also evaluated as carriers for delivery of antineoplastic agents. In particular, erythrocytes were used as carriers for anticancer agent 5-fluorouracil (5-FU) to treat malignant ascites [96]. Intravenous injections of the cell carriers loaded with 5-FU resulted in significant regression of the quantity of malignant ascites and an increase of survival time in mice. Notably, mononuclear phagocytes can serve as 'Trojan horses' for nanoparticle transport into tumor regions [85,89]. Thus, infiltration of monocytes and active delivery of Au-containing nanoparticles to the center of tumor spheroids (which contain hypoxic centers) were reported [89]. Following the delivery the tumor spheroids were destroyed by irradiation with near-infrared light, which actuated the delivered Au-containing nanoparticles. In another study, macrophages acted as a cellular vehicle for 5-FU encapsulated in oligomannose-coated liposomes (OMLs) [50]. The successful decrease of tumor growth by co-administration of OML-encased 5-FU and OML-encased magnetic nanoparticles followed by treatment with an alternating magnetic field was reported in mice. These OMLs were injected into the peritoneal cavity, and then gradually accumulated in the omentum and the other lymphoid tissues within 24 h. Treatment of macrophages at 39°C for 30 min led to the release of 5-FU from the macrophages, suggesting that controlled release from macrophages could be achieved by mild hyperthermia. The encased magnetic nanoparticles, which are known to convert electromagnetic energy to heat in the OMLs, allowed *in vivo* hyperthermia to be achieved. Finally, mononuclear phagocytes were utilized for delivery of therapeutic DNA constructs into tumors [85].

Bacterially derived minicells with diameters of 400 nm were suggested for targeted delivery of different chemotherapeutics and small interfering RNA (siRNA) duplexes that cause stabilization and regression of tumor size in human cancer xenografts in mice [97,98]. The cell carriers were specifically targeted to tumor cell-surface proteins with biospecific antibodies (BsAb). About 80% efficiency of packing minicells with siRNA was reported. Thus, minicells carrying at least 12,000 siRNA molecules and 100 copies of short hairpin RNA (shRNA)-encoding plasmid can specifically and sequentially deliver to tumor xenografts these therapeutics and compromise drug resistance by knocking down a drug-resistant protein. This cell-mediated drug delivery system enables one to use several thousand-fold less drug than is needed for conventional systemic administration of cancer therapies. Evaluation of uptake and intracellular kinetics of minicell-delivered siRNA indicates that after endocytosis the minicells traverse the well-established early and late endosomal pathways, terminating into acidified organelles, the lysosomes, where minicells are degraded and release their cargo [98]. In addition, several methods of packaging in minicells of a range of anticancer chemotherapeutic drugs, such as doxorubicin, paclitaxel, irinotecan, 5-FLU, cisplatin, carboplatin monastrol and vinblastine, despite their disparate structure, charge, hydrophobicity and solubility, were developed [99-101].

Antibody-producing hybridoma cells were used for delivery of anti-CD137 and anti-OX40 mAb [102]. Microcapsules containing viable cells that secrete antibody were implanted by injection into the subcutaneous tissue of Balb/c mice bearing CT26 colon carcinomas. CD137 and OX40 are members of the TNF receptor family that provide costimulatory effect on T lymphocytes. The treatment resulted in complete tumor eradication in an elevated fraction of cases and strong tumor-specific cytotoxic T-lymphocyte responses with either anti-CD137- or anti-OX40-producing hybridomas.

Besides cytotoxic agents, delivery of imaging agents is of importance for cancer therapy. Mononuclear phagocyte-mediated delivery of quantum dots to brain tumors was explored for this purpose [103]. Thus, the surgical management of brain tumors requires the precise localization of tumor tissues within normal brain parenchyma in order to achieve accurate diagnostic biopsy and complete surgical resection. To this end, quantum dots are optical semiconductor nanocrystals that show stable, bright fluorescence. It was demonstrated that the intravenous injection of quantum dots is accompanied by reticuloendothelial system and macrophage sequestration. Macrophages infiltrate brain tumors and phagocytize intravenously injected quantum dots, optically labeling the tumors.

Another class of imaging agents, *N*-(2-hydroxypropyl)methacrylamide (HPMA) copolymer-gadolinium (Gd) chelates, was used for enhanced MRI of macrophage-mediated malignancies [104]. To increase the nanoparticles uptake in macrophages, copolymers were vectorized with mannoseamine, which was shown to increase mannose receptor-mediated uptake in

human MP. It was suggested that intravenously administered Gd-nanoparticles will be accumulated in macrophages and then transported to the pathological sites (e.g., solid tumor). Overall, the tumor recruitment of inflammatory-response cells may be exploited for cell-mediated delivery of antineoplastic and imaging agents.

5. Lung injury

Drug delivery to small airways, terminal bronchioles and alveoli is complicated owing to the methodological limitations in targeting the deep lung with high-efficiency drug distribution to the site of pathology. To overcome these limitations, chitosan nanoparticles with entrapped curcumin (an anti-inflammatory compound) were loaded into testis-derived Sertoli cells and then injected intravenously into mice with bronchial or alveolar injuries [105]. By 24 h post-injection, most of the curcumin load (~ 90%) delivered into the injected Sertoli cells was present and distributed throughout the lungs, including the perivascular sac area in the lower lungs. These results identify a new and efficient method for targeted delivery of drugs for treatment of bronchitis and alveolitis. In addition, lungs are suitable for both local and systemic drug delivery, therefore this route may be used broadly for other diseases.

6. Microbial infections

The efficiency of antifungal agents has been diminished by their severe side effects and poor pharmacokinetics. To this end, erythrocytes were suggested as delivery vehicles for antifungal agent amphotericin B (AmB). To avoid toxic effects and achieve efficient drug loading, AmB was encapsulated in nanosuspension (AmB-NS) by high-pressure homogenization [106]. AmB-NS was loaded into RBCs by using hypotonic hemolysis, leading to intracellular AmB amounts of 3.81 ± 0.47 pg/RBC and an entrapment efficacy of 15 – 18%. Hence, RBCs served as primary carriers. Then, the uptake of RBC loaded with AmB-NS by leukocytes was studied by flow cytometry. More than 98% of the phagocyte population (granulocytes and peripheral monocytes) accumulated AmB-NS after 4 h of incubation, and then showed a slow AmB release over 10 days without any alteration in cell viability. This results in an immediate, permanent inhibition of intra- and extracellular fungal activity. AmB-NS-RBC-leukocyte-mediated delivery of AmB was efficient in amounts 1000 times lower than the toxic dose.

Another antifungal agent, chloroquine, loaded into phosphatidylserine-containing negatively charged liposomes was used in MP-mediated delivery against *Cryptococcus* infection in the mouse brain [107]. Administration of chloroquine-loaded liposomes accumulated inside macrophage phagolysosomes resulted in a remarkable reduction in fungal load in the brain even in low doses compared with free drug in high doses, thus increasing antifungal activity of

macrophages. Therefore, this drug delivery method is effective for the transport of water-insoluble substances such as AmB, and this warrants consideration for further testing.

7. Epilepsy

A relatively straightforward method of engineering adenosine-releasing cells was developed for the treatment of epilepsy [108]. Lack of adenosine, a modulator of neuronal activity with anticonvulsant and neuroprotective properties, was found to contribute to ictogenesis. Therefore, focal reconstitution of adenosine within an epileptogenic brain region constitutes a rational therapeutic approach, whereas systemic augmentation of adenosine is precluded by side effects. To this end, human mesenchymal stem cells and human embryonic stem cells were embedded in a cell-encapsulation device to release adenosine and reduce acute injury and seizures, as well as chronic seizures [108]. Human embryonic stem cells have a high proliferative capacity and can be subjected to specific cellular differentiation pathways. Human embryonic stem cells differentiated *in vitro* into neuroepithelial cells and grafted into the mouse brain displayed intrahippocampal location and neuronal morphology. Overall, this therapeutic approach demonstrated antiepileptic and neuroprotective properties when grafted into the mouse hippocampus. The therapeutic potential of this approach suggests the feasibility of engineering autologous adenosine-releasing stem cells derived from a patient [109].

8. Expert opinion

The concept of using cells as drug delivery vehicles is under development. Challenges include sufficient loading capacity and triggered drug release, preservation of therapeutic agents against degradation inside the cells, protection of cell carriers from drug cytotoxic effects, and efficient homing of cells to the disease site. In addition, harvesting of cells in sufficient quantities or their expansion is also important issue.

Two different approaches are utilized in cell-mediated drug delivery. First, cell carriers (predominantly mononuclear phagocytes) are genetically modified to produce therapeutically active molecules. Second, host cells are loaded with a drug usually incorporated into protective container, and then used as 'Trojan horses' to deliver the drug to the disease site (Figure 8). Most cell-based delivery systems suggest unloading the therapeutic agent at the disease site. However, a few use another approach, when the targeted cells accumulate in the carrier along with the drug (minicells with anticancer agents and red blood cells with antifungal agents).

Regarding the first approach, the ideal cell delivery vehicle would be stable in tissue culture and capable of sustained, preferably regulated expression of therapeutic molecules. The cells should have appropriate and predictable differentiation pattern and survive for a long time *in vivo* after transplantation. Obviously, for both approaches the cell carriers should

Disease	Cell-based drug delivery systems	
	Genetically modified cell carriers	Drug-loaded cell carriers
PD, AD	NSCs or BMM producing neurotrophic factors	BMM loaded with redox enzymes
HIV	Monocytes and BMM with antiretroviral agents	
Cancer	<ul style="list-style-type: none"> • MSCs producing antitumor proteins • Antibody-producing hybridoma cells 	<ul style="list-style-type: none"> • T cells, RBC, and BMM loaded with chemotherapeutics • BMM loaded with Au- or magnetic nanoparticles • MSCs with nanoparticles attached to the surface • Mini cells loaded with chemotherapeutics or siRNA • BMM loaded with imaging agents
Lung injury	Sertoli cells loaded with anti-inflammatory agents	
Microbial infections	RBC loaded with antifungal agents	
Epilepsy	MSCs producing adenosine	

Figure 8. Cell-based drug delivery systems.

AD: Alzheimer's disease; BMM: Bone marrow-derived macrophages; MSCs: Mesenchymal stem cells; NSCs: Neural stem cells; PD: Parkinson's disease; RBC: Red blood cell; siRNA: Small interfering RNA.

demonstrate responsiveness to the chemotactic signals produced by the type of pathology that they are used to treat. Regarding the second approach, loading cell carriers with drug in the bloodstream by injecting nanoparticulate drug targeted to these cells is the most attractive strategy. To this end, it is important to note that these cells are moving targets. Nanosystems that bind to these targets are potentially a powerful approach for imaging their movement and delivering a therapeutic drug dose to the disease site. Clearly, a multifunctional molecular scaffold that is aimed at delivery of drug specifically to the cell carrier but in addition could be engineered for imaging purposes would offer substantial benefits over current approaches.

Such a strategy is opposite to the common approach when drug-loaded nanoparticles are designed to minimize their entrapment in the reticuloendothelial system and avoid drug decomposition and clearance. Consequently, the design of nanocarriers for cell-mediated drug delivery (size, shape, surface charge, etc.) may differ

from those used for conventional drug delivery systems. Nevertheless, engaging different defense mechanisms into drug delivery may open a new perspective of active delivery of drugs for the treatment of various devastating diseases.

Declaration of interest

This work was supported by the United States National Institutes of Health grants 1R01 NS057748 (to EV Batrakova), 2R01 NS034239, 2R37 NS36126, P01 NS31492, P20RR 15635, P01 MH64570, 5 P01 DA026146, 1P01 DA028555, P01 NS43985 (to HE Gendelman), RO1 NS051334, 2RO1 CA89225, 1 R01 CA116591, UO1 CA151806, 1P20 RR021937, the United States Department of Defense grants W81XWH-09-1-0386 and DoD USAMRMC 06108004, and the Government of Russian Federation grants 02.740.11.523 and 11.G4.31.0004 (to AV Kabanov).

Bibliography

Papers of special note have been highlighted as either of interest (●) or of considerable interest (●●) to readers.

1. Fujiwara M, Baldeschwieler JD, Grubbs RH. Receptor-mediated endocytosis of poly(acrylic acid)-conjugated liposomes by macrophages. *Biochim Biophys Acta* 1996;1278(1):59-67
2. Torchilin VP. Drug targeting. *Eur J Pharm Sci* 2000;11(Suppl 2):S81-91
3. Mora M, Sagrista ML, Trombetta D, et al. Design and characterization of liposomes containing long-chain N-acylPEs for brain delivery: penetration of liposomes incorporating GM1 into the rat brain. *Pharm Res* 2002;19(10):1430-8
4. Aoki H, Kakinuma K, Morita K, et al. Therapeutic efficacy of targeting chemotherapy using local hyperthermia and thermosensitive liposome: evaluation of drug distribution in a rat glioma model. *Int J Hyperthermia* 2004;20(6):595-605
5. Kabanov AV, Chekhonin VP, Alakhov V, et al. The neuroleptic activity of haloperidol increases after its solubilization in surfactant micelles. Micelles as microcontainers for drug targeting. *FEBS Lett* 1989;258(2):343-5
6. Kabanov AV, Vinogradov SV, Suzdaltseva YG, et al. Water-soluble block polycations as carriers for oligonucleotide delivery. *Bioconjug Chem* 1995;6(6):639-43
7. Kabanov A, Alakhov V. Pluronic block copolymers in drug delivery: from micellar nanocontainers to biological response modifiers. *Crit Rev Ther Drug Carrier Syst* 2002;19(1):1-72
8. Kwon GS. Polymeric micelles for delivery of poorly water-soluble compounds. *Crit Rev Ther Drug Carrier Syst* 2003;20(5):357-403
9. Vinogradov S, Batrakova E, Kabanov A. Poly(ethylene glycol)-polyethyleneimine NanoGel (TM) particles: novel drug delivery systems for antisense oligonucleotides. *Colloids Surf B Biointerfaces* 1999;16(1-4):291-304
10. Vinogradov SV, Batrakova EV, Kabanov AV. Nanogels for oligonucleotide delivery to the brain. *Bioconjug Chem* 2004;15(1):50-60
11. Gref R, Minamitake Y, Peracchia M, et al. Biodegradable long-circulating polymeric nanospheres. *Science* 1994;263(5153):1600-3
12. Hyuk IS, Jeong U, Xia Y. Polymer hollow particles with controllable holes in their surfaces. *Nat Mater* 2005;4(9):671-5
13. Calvo P, Gouritin B, Chacun H, et al. Long-circulating PEGylated polycyanoacrylate nanoparticles as new drug carrier for brain delivery. *Pharm Res* 2001;18(8):1157-66
14. Nowacek AS, Miller RL, McMillan J, et al. NanoART synthesis, characterization, uptake, release and toxicology for human monocyte-macrophage drug delivery. *Nanomomed* 2009;4(8):903-17
- **This study describes characterization of nanoformulated antiretroviral drugs for HIV-1 infection and its application for human cell-mediated drug delivery.**
15. Muller RH, Jacobs C, Kayser O. Nanosuspensions as particulate drug formulations in therapy. Rationale for development and what we can expect for the future. *Adv Drug Deliv Rev* 2001;47(1):3-19
16. Friedrich I, Reichl S, Muller-Goymann CC. Drug release and permeation studies of nanosuspensions based on solidified reverse micellar solutions (SRMS). *Int J Pharm* 2005;305(1-2):167-75
17. Harada A, Kataoka K. Chain length recognition: core-shell supramolecular assembly from oppositely charged block copolymers. *Science* 1999;283(5398):65-7
18. Jaturanpinyo M, Harada A, Yuan X, et al. Preparation of bionanoreactor based on core-shell structured polyion complex micelles entrapping trypsin in the core cross-linked with glutaraldehyde. *Bioconjug Chem* 2004;15(2):344-8
19. Bull SR, Guler MO, Bras RE, et al. Self-assembled peptide amphiphile nanofibers conjugated to MRI contrast agents. *Nano Lett* 2005;5(1):1-4
20. Guler MO, Pokorski JK, Appella DH, et al. Enhanced oligonucleotide binding to self-assembled nanofibers. *Bioconjug Chem* 2005;16(3):501-3
21. Juliano R, Stamp D. The effect of particle size and charge on the clearance rates of liposomes and liposome encapsulated drugs. *Biochem Biophys Res Commun* 1975;63:651-8
22. Lee KD, Hong K, Papahadjopoulos D. Recognition of liposomes by cells: in vitro binding and endocytosis mediated by specific lipid headgroups and surface charge density. *Biochim Biophys Acta* 1992;1103(2):185-97
23. Nishikawa K, Arai H, Inoue K. Scavenger receptor-mediated uptake and metabolism of lipid vesicles containing acidic phospholipids by mouse peritoneal macrophages. *J Biol Chem* 1990;265(9):5226-31
24. Miller CR, Bondurant B, McLean SD, et al. Liposome-cell interactions in vitro: effect of liposome surface charge on the binding and endocytosis of conventional and sterically stabilized liposomes. *Biochemistry* 1998;37(37):12875-83
25. Ishihara T, Izumo N, Higaki M, et al. Role of zinc in formulation of PLGA/PLA nanoparticles encapsulating betamethasone phosphate and its release profile. *J Control Release* 2005;105(1-2):68-76
26. Tempone AG, Perez D, Rath S, et al. Targeting Leishmania (L.) chagasi amastigotes through macrophage scavenger receptors: the use of drugs entrapped in liposomes containing phosphatidylserine. *J Antimicrob Chemother* 2004;54(1):60-8
27. Krieger M. Molecular flypaper and atherosclerosis: structure of the macrophage scavenger receptor. *Trends Biochem Sci* 1992;17(4):141-6
28. Aderem A, Underhill DM. Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol* 1999;17:593-623
29. Zhao Y, Haney MJ, Klyachko NL, et al. Polyelectrolyte complex optimization for macrophage delivery of redox enzyme nanoparticles. *Nanomedicine (Lond)* 2011;6(1):25-42
30. Harada A, Kataoka K. Pronounced activity of enzymes through the incorporation into the core of polyion complex micelles made from charged

- block copolymers. *J Control Release* 2001;72(1-3):85-91
31. Harada A, Kataoka K. Switching by pulse electric field of the elevated enzymatic reaction in the core of polyion complex micelles. *J Am Chem Soc* 2003;125(50):15306-7
 32. Knop K, Hoogenboom R, Fischer D, et al. Poly(ethylene glycol) in drug delivery: pros and cons as well as potential alternatives. *Angew Chem Int Ed Engl* 2010;49(36):6288-308
 33. Gabizon A, Shmeeda H, Barenholz Y. Pharmacokinetics of pegylated liposomal Doxorubicin: review of animal and human studies. *Clin Pharmacokinet* 2003;42(5):419-36
 34. Papaldo P, Fabi A, Ferretti G, et al. A phase II study on metastatic breast cancer patients treated with weekly vinorelbine with or without trastuzumab according to HER2 expression: changing the natural history of HER2-positive disease. *Ann Oncol* 2006;17(4):630-6
 35. Gbadamosi JK, Hunter AC, Moghimi SM. PEGylation of microspheres generates a heterogeneous population of particles with differential surface characteristics and biological performance. *FEBS Lett* 2002;532(3):338-44
 36. Daleke DL, Hong K, Papahadjopoulos D. Endocytosis of liposomes by macrophages: binding, acidification and leakage of liposomes monitored by a new fluorescence assay. *Biochim Biophys Acta* 1990;1024(2):352-66
 37. Jain S, Mishra V, Singh P, et al. RGD-anchored magnetic liposomes for monocytes/neutrophils-mediated brain targeting. *Int J Pharm* 2003;261(1-2):43-55
 38. Thiele L, Merkle HP, Walter E. Phagocytosis and phagosomal fate of surface-modified microparticles in dendritic cells and macrophages. *Pharm Res* 2003;20(2):221-8
 39. Fahmy TM, Schneck JP, Saltzman WM. A nanoscopic multivalent antigen-presenting carrier for sensitive detection and drug delivery to T cells. *Nanomedicine* 2007;3(1):75-85
 40. Tabata Y, Ikada Y. Effect of the size and surface charge of polymer microspheres on their phagocytosis by macrophage. *Biomaterials* 1988;9(4):356-62
 41. Mizushima Y, Hamano T, Yokoyama K. Tissue distribution and anti-inflammatory activity of corticosteroids incorporated in lipid emulsion. *Ann Rheum Dis* 1982;41(3):263-7
 42. Hillaireau H, Couvreur P. Nanocarriers' entry into the cell: relevance to drug delivery. *Cell Mol Life Sci* 2009;66(17):2873-96
 43. Cannon GJ, Swanson JA. The macrophage capacity for phagocytosis. *J Cell Sci* 1992;101(Pt 4):907-13
 44. Champion JA, Katare YK, Mitragotri S. Making polymeric micro- and nanoparticles of complex shapes. *Proc Natl Acad Sci USA* 2007;104(29):11901-4
 45. Behr J-P. The proton sponge: a trick to enter cells the viruses did not exploit. *CHIMIA. Int J Chem* 1997;51(1):34-6
 46. Cheng H, Kastrup CJ, Ramanathan R, et al. Nanoparticulate cellular patches for cell-mediated tumor tropic delivery. *ACS Nano* 2010;4(2):625-31
 47. Krantz A. Red cell-mediated therapy: opportunities and challenges. *Blood Cells Mol Dis* 1997;23(1):58-68
 48. Batrakova EV, Li S, Reynolds AD, et al. A macrophage-nanozyme delivery system for Parkinson's disease. *Bioconjug Chem* 2007;18(5):1498-506
 - **This paper reported the development and characterization of a cell-mediated delivery system of antioxidant enzyme catalase. Loading, release and protection of enzymatic activity of catalase in bone marrow-derived monocytes are evaluated in an *in vitro* model of Parkinson's disease.**
 49. Sollner T, Bennett MK, Whiteheart SW, et al. A protein assembly-disassembly pathway *in vitro* that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell* 1993;75(3):409-18
 50. Ikehara Y, Niwa T, Biao L, et al. A carbohydrate recognition-based drug delivery and controlled release system using intraperitoneal macrophages as a cellular vehicle. *Cancer Res* 2006;66(17):8740-8
 51. Perry VH, Bell MD, Brown HC, et al. Inflammation in the nervous system. *Curr Opin Neurobiol* 1995;5(5):636-41
 52. Kuby J. *Immunology*. Freeman, WH. and Co., New York; 1994
 53. Anthony DC, Bolton SJ, Fearn S, et al. Age-related effects of interleukin-1 beta on polymorphonuclear neutrophil-dependent increases in blood-brain barrier permeability in rats. *Brain* 1997;120(Pt 3):435-44
 54. Anthony DC, Blond D, Dempster R, et al. Chemokine targets in acute brain injury and disease. *Prog Brain Res* 2001;132:507-24
 55. Blamire AM, Anthony DC, Rajagopalan B, et al. Interleukin-1beta-induced changes in blood-brain barrier permeability, apparent diffusion coefficient, and cerebral blood volume in the rat brain: a magnetic resonance study. *J Neurosci* 2000;20(21):8153-9
 56. Persidsky Y, Ghorpade A, Rasmussen J, et al. Microglial and astrocyte chemokines regulate monocyte migration through the blood-brain barrier in human immunodeficiency virus-1 encephalitis. *Am J Pathol* 1999;155(5):1599-611
 57. Dou H, Destache CJ, Morehead JR, et al. Development of a macrophage-based nanoparticle platform for antiretroviral drug delivery. *Blood* 2006;108(8):2827-35
 58. Pawlowski NA, Kaplan G, Abraham E, et al. The selective binding and transmigration of monocytes through the junctional complexes of human endothelium. *J Exp Med* 1988;168(5):1865-82
 59. Lossinsky AS, Shivers RR. Structural pathways for macromolecular and cellular transport across the blood-brain barrier during inflammatory conditions [review]. *Histol Histopathol* 2004;19(2):535-64
 60. Orlic D, Kajstura J, Chimenti S, et al. Bone marrow cells regenerate infarcted myocardium. *Nature* 2001;410(6829):701-5
 61. Orlic D, Kajstura J, Chimenti S, et al. Mobilized bone marrow cells repair the infarcted heart, improving function and survival. *Proc Natl Acad Sci USA* 2001;98(18):10344-9
 62. Hofstetter CP, Schwarz EJ, Hess D, et al. Marrow stromal cells form guiding

- strands in the injured spinal cord and promote recovery. *Proc Natl Acad Sci USA* 2002;99(4):2199-204
63. Mahmood A, Lu D, Lu M, et al. Treatment of traumatic brain injury in adult rats with intravenous administration of human bone marrow stromal cells. *Neurosurgery* 2003;53(3):697-702, discussion-3
64. Martinez-Serrano A, Hantzopoulos PA, Bjorklund A. Ex vivo gene transfer of brain-derived neurotrophic factor to the intact rat forebrain: neurotrophic effects on cholinergic neurons. *Eur J Neurosci* 1996;8(4):727-35
65. Martinez-Serrano A, Bjorklund A. Ex vivo nerve growth factor gene transfer to the basal forebrain in presymptomatic middle-aged rats prevents the development of cholinergic neuron atrophy and cognitive impairment during aging. *Proc Natl Acad Sci USA* 1998;95(4):1858-63
66. Muller FJ, Snyder EY, Loring JF. Gene therapy: can neural stem cells deliver? *Nat Rev Neurosci* 2006;7(1):75-84
67. Leek RD, Lewis CE, Whitehouse R, et al. Association of macrophage infiltration with angiogenesis and prognosis in invasive breast carcinoma. *Cancer Res* 1996;56(20):4625-9
68. Lewis JS, Landers RJ, Underwood JC, et al. Expression of vascular endothelial growth factor by macrophages is up-regulated in poorly vascularized areas of breast carcinomas. *J Pathol* 2000;192(2):150-8
69. Ranney DF, Huffaker HH. Magnetic microspheres for the targeted controlled release of drugs and diagnostic agents. *Ann NY Acad Sci* 1987;507:104-19
70. Hendriks JJ, Teunissen CE, de Vries HE, et al. Macrophages and neurodegeneration. *Brain Res Brain Res Rev* 2005;48(2):185-95
71. Brynskikh AM, Zhao Y, Mosley RL, et al. Macrophage delivery of therapeutic nanozymes in a murine model of Parkinson's disease. *Nanomedicine (Lond)* 2010;5(3):379-96
- **This study supports the feasibility of cell-mediated drug delivery to the brain by examining: the nanozyme loading capacity for cell carriers; the effect of nanozymes on cell viability and function; and the neuroprotective activities of BMM-carried nanozyme against MPTP intoxication in an *in vivo* model of Parkinson's disease.**
72. Low WC, Lewis PR, Bunch ST, et al. Function recovery following neural transplantation of embryonic septal nuclei in adult rats with septohippocampal lesions. *Nature* 1982;300(5889):260-2
73. Garcia P, Youssef I, Utvik JK, et al. Ciliary neurotrophic factor cell-based delivery prevents synaptic impairment and improves memory in mouse models of Alzheimer's disease. *J Neurosci* 2010;30(22):7516-27
74. Pizzo DP, Coufal NG, Lortie MJ, et al. Regulatable acetylcholine-producing fibroblasts enhance cognitive performance. *Mol Ther* 2006;13(1):175-82
75. Blurton-Jones M, Kitazawa M, Martinez-Coria H, et al. Neural stem cells improve cognition via BDNF in a transgenic model of Alzheimer disease. *Proc Natl Acad Sci USA* 2009;106(32):13594-9
76. Zurn AD, Tseng J, Aebischer P. Treatment of Parkinson's disease. Symptomatic cell therapies: cells as biological minipumps. *Eur Neurol* 1996;36(6):405-8
77. Akerud P, Canals JM, Snyder EY, et al. Neuroprotection through delivery of glial cell line-derived neurotrophic factor by neural stem cells in a mouse model of Parkinson's disease. *J Neurosci* 2001;21(20):8108-18
78. Casper D, Engstrom SJ, Mirchandani GR, et al. Enhanced vascularization and survival of neural transplants with ex vivo angiogenic gene transfer. *Cell Transplant* 2002;11(4):331-49
79. Yasuhara T, Shingo T, Muraoka K, et al. Neurorescue effects of VEGF on a rat model of Parkinson's disease. *Brain Res* 2005;1053(1-2):10-8
80. Biju K, Zhou Q, Li G, et al. Macrophage-mediated GDNF delivery protects against dopaminergic neurodegeneration: a therapeutic strategy for Parkinson's disease. *Mol Ther* 2010;18(8):1536-44
81. Dou H, Morehead JR, Destache C, et al. Laboratory investigations for the morphologic, pharmacokinetic, and anti-retroviral properties of indinavir nanoparticles in human monocyte-derived macrophages. *Virology* 2007;358(1):148-58
82. Dou H, Grotetas CB, McMillan JM, et al. Macrophage delivery of nanoformulated antiretroviral drug to the brain in a murine model of neuroAIDS. *J Immunol* 2009;183(1):661-9
83. Nowacek AS, McMillan J, Miller R, et al. Nanoformulated antiretroviral drug combinations extend drug release and antiretroviral responses in HIV-1-infected macrophages: implications for neuroAIDS therapeutics. *J Neuroimmune Pharmacol* 2010;5(4):592-601
84. Duzgunes N, Pretzer E, Simoes S, et al. Liposome-mediated delivery of antiviral agents to human immunodeficiency virus-infected cells. *Mol Membr Biol* 1999;16(1):111-18
85. Chokri M, Lopez M, Oleron C, et al. Production of human macrophages with potent antitumor properties (MAK) by culture of monocytes in the presence of GM-CSF and 1,25-dihydroxy vitamin D3. *Anticancer Res* 1992;12(6B):2257-60
86. Studeny M, Marini FC, Champlin RE, et al. Bone marrow-derived mesenchymal stem cells as vehicles for interferon-beta delivery into tumors. *Cancer Res* 2002;62(13):3603-8
87. Stagg J, Lejeune L, Paquin A, et al. Marrow stromal cells for interleukin-2 delivery in cancer immunotherapy. *Hum Gene Ther* 2004;15(6):597-608
88. Nakamura K, Ito Y, Kawano Y, et al. Antitumor effect of genetically engineered mesenchymal stem cells in a rat glioma model. *Gene Ther* 2004;11(14):1155-64
89. Choi MR, Stanton-Maxey KJ, Stanley JK, et al. A cellular Trojan horse for delivery of therapeutic nanoparticles into tumors. *Nano Lett* 2007;7(12):3759-65
90. Studeny M, Marini FC, Dembinski JL, et al. Mesenchymal stem cells: potential precursors for tumor stroma and targeted-delivery vehicles for anticancer agents. *J Natl Cancer Inst* 2004;96(21):1593-603
91. Stoff-Khalili MA, Rivera AA, Mathis JM, et al. Mesenchymal stem cells as a vehicle for targeted delivery of CRAds to lung

- metastases of breast carcinoma. *Breast Cancer Res Treat* 2007;105(2):157-67
92. Sonabend AM, Ulasov IV, Tyler MA, et al. Mesenchymal stem cells effectively deliver an oncolytic adenovirus to intracranial glioma. *Stem Cells* 2008;26(3):831-41
93. Menon LG, Kelly K, Yang HW, et al. Human bone marrow-derived mesenchymal stromal cells expressing S-TRAIL as a cellular delivery vehicle for human glioma therapy. *Stem Cells* 2009;27(9):2320-30
94. Steinfeld U, Pauli C, Kaltz N, et al. T lymphocytes as potential therapeutic drug carrier for cancer treatment. *Int J Pharm* 2006;311(1-2):229-36
95. Nakamizo A, Marini F, Amano T, et al. Human bone marrow-derived mesenchymal stem cells in the treatment of gliomas. *Cancer Res* 2005;65(8):3307-18
96. Wang GP, Guan YS, Jin XR, et al. Development of novel 5-fluorouracil carrier erythrocyte with pharmacokinetics and potent antitumor activity in mice bearing malignant ascites. *J Gastroenterol Hepatol* 2010;25(5):985-90
97. MacDiarmid JA, Mugridge NB, Weiss JC, et al. Bacterially derived 400 nm particles for encapsulation and cancer cell targeting of chemotherapeutics. *Cancer Cell* 2007;11(5):431-45
98. MacDiarmid JA, Amaro-Mugridge NB, Madrid-Weiss J, et al. Sequential treatment of drug-resistant tumors with targeted minicells containing siRNA or a cytotoxic drug. *Nat Biotechnol* 2009;27(7):643-51
99. Nikaido H. Multidrug efflux pumps of gram-negative bacteria. *J Bacteriol* 1996;178(20):5853-9
100. Nikaido H. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev* 2003;67(4):593-656
101. Poole K. Outer membranes and efflux: the path to multidrug resistance in Gram-negative bacteria. *Curr Pharm Biotechnol* 2002;3(2):77-98
102. Dubrot J, Portero A, Orive G, et al. Delivery of immunostimulatory monoclonal antibodies by encapsulated hybridoma cells. *Cancer Immunol Immunother* 2010;59(11):1621-31
103. Popescu MA, Toms SA. In vivo optical imaging using quantum dots for the management of brain tumors. *Expert Rev Mol Diagn* 2006;6(6):879-90
104. Zarabi B, Nan A, Zhuo J, et al. Macrophage targeted N-(2-hydroxypropyl)methacrylamide conjugates for magnetic resonance imaging. *Mol Pharm* 2006;3(5):550-7
105. Kumar A, El-Badri N, Glaum M, et al. Initial observations of cell mediated drug delivery to the deep lung. *Cell Transplant* 5 Nov 2010 [Epub ahead of print]
106. Staedtke V, Brahler M, Muller A, et al. In vitro inhibition of fungal activity by macrophage-mediated sequestration and release of encapsulated amphotericin B nanosuspension in red blood cells. *Small* 2010;6(1):96-103
107. Khan MA, Jabeen R, Nasti TH, et al. Enhanced anticryptococcal activity of chloroquine in phosphatidylserine-containing liposomes in a murine model. *J Antimicrob Chemother* 2005;55(2):223-8
108. Boison D. Engineered adenosine-releasing cells for epilepsy therapy: human mesenchymal stem cells and human embryonic stem cells. *Neurotherapeutics* 2009;6(2):278-83
109. Boison D. Adenosine augmentation therapies (AATs) for epilepsy: prospect of cell and gene therapies. *Epilepsy Res* 2009;85(2-3):131-41
110. Kabanov AV, Batrakova EV. Polymer nanomaterials. In: Gendelman HE, Ikezu T, editors. *Neuroimmune pharmacology*. Springer, Omaha; 2008. p. 691-707
111. Nowacek A, Gendelman HE. NanoART, neuroAIDS and CNS drug delivery. *Nanomedicine* 2009;4(5):557-74

Affiliation

Elena V Batrakova^{†1},
Howard E Gendelman^{1,2,3} &
Alexander V Kabanov^{1,3,4}
[†]Author for correspondence
¹University of Nebraska Medical Center,
Center for Drug Delivery and Nanomedicine,
Department of Pharmaceutical Sciences,
Omaha, NE 68198-5830, USA
Tel: +1 402 559 9364; Fax: +1 402 559 9365;
E-mail: ebatrako@unmc.edu
²University of Nebraska Medical Center,
Center for Neurodegenerative Disorders,
Omaha, NE 68198-5830, USA
³University of Nebraska Medical Center,
Department of Pharmacology and
Experimental Neuroscience,
Omaha, NE 68198-5830, USA
⁴MV Lomonosov Moscow State University,
Department of Chemistry,
Moscow, 119991 Russia