

## PHOTOSENSITIZATION AND QUENCHING OF SINGLET OXYGEN BY PIGMENTS AND LIPIDS OF PHOTORECEPTOR CELLS OF THE RETINA

A. A. KRASNOVSKY, jr and V. E. KAGAN

*Department of Biology, Moscow State University, Moscow 117234, USSR*

Received 2 October 1979

### 1. Introduction

Studies of  $^1\text{O}_2$  generation and quenching in solutions of pigments and lipids isolated from photoreceptor cells are important for understanding the molecular mechanisms of photodamage of the retina [1–5]. The first works in this field were published by Delmelle ([4] and references therein), who attempted to demonstrate the ability of retinal to produce  $^1\text{O}_2$ , using indirect 'chemical' methods of  $^1\text{O}_2$  detection. Here we have employed a more accurate method, i.e., registration of oxygen luminescence observed in air-saturated solutions of various pigments ([5,6] and references therein). With the aid of this emission the following questions were studied: is retinal able to generate  $^1\text{O}_2$  and what is the efficiency of  $^1\text{O}_2$  interaction with retinal and photoreceptor lipids?

### 2. Materials and methods

The luminescence was measured in an apparatus supplied with a photomultiplier and a mechanical phosphoroscope as in [6].  $\text{CCl}_4$  of spectral grade was used as a solvent. The retinals were synthesized in the M. V. Lomonosov Institute of Fine Chemical Technology. Egg lecithin, phosphatidylcholine and phosphatidylethanolamine from frog retina were isolated according to [7] and purified by thin-layer chromatography.

### 3. Results and discussion

#### 3.1. Generation of $^1\text{O}_2$ by retinal

Illumination of air-saturated solutions of 11-*cis*, 13-*cis* and all-*trans* retinals results in a delayed

luminescence in the region of  $^1\text{O}_2$  emission from  $^1\Delta_g$  state (1270 nm). The excitation spectra coincide with the absorption spectra of the pigments (fig.1.). The lifetime of the luminescence ( $\tau$ ) is equal to  $28 \pm 3$  ms, which corresponds to  $\tau$   $^1\text{O}_2$  in  $\text{CCl}_4$  [6]. The  $^1\text{O}_2$  quenchers,  $\beta$ -carotene,  $\alpha$ -tocopherol, 1,3-diphenylisobenzofuran (DPIBP) and 1,4-diazabicyclo(2,2,2)-octane (DABCO) quench the luminescence. The rate constants of quenching ( $k_q$ ) coincide with those for  $^1\text{O}_2$  determined by other methods (table 1).

The data obtained suggest that the luminescence results from emission of  $^1\text{O}_2$  formed due to energy transfer from the photoexcited molecules of retinals. The quantum yield of  $^1\text{O}_2$  generation ( $\gamma_g$ ) was evaluated by comparing the intensity of  $^1\text{O}_2$  emission in  $\text{CCl}_4$  solutions of retinals and porphyrins (protoporphyrin and tetraphenylporphyrin,  $\gamma_g = 0.8$  [6]). The obtained value of  $\gamma_g$  for the retinals is equal to  $0.55 \pm 0.1$ , which exceeds the value reported by Delmelle for retinal in ethanol (0.09) [4].

#### 3.2. Quenching of $^1\text{O}_2$ luminescence

This was studied in solutions of protoporphyrin ( $1.5 \times 10^{-7}$  M) upon illumination by red light at 627 nm. The retinals, retinyl acetate and lipids were found to quench  $^1\text{O}_2$  in accordance with the Stern–Volmer equation (fig.1). The values of  $k_q$  calculated from this equation suggest that these compounds are relatively weak quenchers of  $^1\text{O}_2$  (table 2). Amongst the retinals the all-*trans* isomer has the highest and 13-*cis* has the lowest activity. The quenching activity of the phospholipids studied was sufficiently lower and did not practically depend on their molecular structure.

In order to elucidate the mechanism of quenching




Fig.1. (A) Comparison of the absorption spectrum (1) with the excitation spectrum (2) of the  $^1\text{O}_2$  luminescence in solutions of 11-cis retinal in  $\text{CCl}_4$ . (B) Quenching of  $^1\text{O}_2$  luminescence by all-trans (1), 11-cis (2) and 13-cis (3) retinals and ROS phosphatidylcholine (4).  $L$ ,  $L_0$ , intensities of the luminescence in the presence and in the absence of the quencher;  $I$ , intensity of the exciting light; OD, absorbance.

the reaction between  $^1\text{O}_2$  and all-trans retinal was studied in more detail [5]. It was shown that  $^1\text{O}_2$  quenching results in a decrease of the absorption bands corresponding to the original pigment and an appearance of new bands at 325 and 340 nm. This

Table 1  
Values of  $k_q$  ( $\text{M}^{-1} \text{s}^{-1}$ ) for quenching of  $^1\text{O}_2$  and the luminescence sensitized by all-trans retinal

Quencher	Luminescence (own data)	$^1\text{O}_2$ (literature data)
DABCO	$6.6 \times 10^6$	$7.3 \times 10^6$ [8]
DPIBP	$8.0 \times 10^8$	$7.0 \times 10^8$ [9]
$\beta$ -Carotene	$8.0 \times 10^9$	$1.1 \times 10^{10}$ [10]
$\alpha$ -Tocopherol	$10^8$	$1.7 \times 10^8$ [11]

process was inhibited by  $\beta$ -carotene, the inhibition rate constant being equal to that for  $^1\text{O}_2$  quenching. The dependence of the reciprocal value of the quantum yield of bleaching ( $1/\psi_{\text{ox}}$ ) on the reverse concentration of retinal ( $1/C$ ) appeared to be linear (fig.2). The intersection of the straight line with the ordinate ( $1/C = 0$ ) corresponds apparently to the maximal value of  $\psi_{\text{ox}}$ , i.e.,  $1/\psi_{\text{ox}}^{\text{max}}$ . Such dependence is known to be characteristic of the photosensitization process involving  $^1\text{O}_2$ . In this case  $1/\psi_{\text{ox}}^{\text{max}} = k_q/\gamma_g \times k_{\text{ox}}$ , where  $k_{\text{ox}}$  is the rate constant of retinal oxidation by  $^1\text{O}_2$  (see also [5,6] and references therein). Hence, the value of  $k_{\text{ox}}$  can be determined from the following equation:

$$k_{\text{ox}} = \frac{\psi_{\text{ox}}^{\text{max}} \times k_q}{\gamma_g}$$

Table 2  
Values of  $k_q$  ( $\text{M}^{-1} \text{s}^{-1}$ ) for retinals and lipids (errors  $\pm 10\%$ )

Quencher	$k_q$	Quencher	$k_q$
All-trans retinal	$3.7 \times 10^6$	Egg lecithin	$5 \times 10^4$
13-cis retinal	$3.6 \times 10^5$	ROS phosphatidylcholine	$10^5$
11-cis retinal	$10^6$	ROS phosphatidylethanolamine	$10^5$
Retinyl acetate	$7 \times 10^5$		




Fig. 2. Protoporphyrin-sensitized photooxidation of all-*trans* retinal in  $CCl_4$ . (A) Absorption spectra of retinal before (1) and after (2) photooxidation. (B) Dependence of quantum yield of photooxidation on retinal concentration. Porphyrin was  $1.5 \times 10^{-7}$  M, the wavelength of the actinic light is 545 nm.

The values of  $k_q$  and  $\gamma_g$  are given above. The value of  $\psi_{ox}^{max}$  was determined by comparing the efficiency of protoporphyrin-sensitized photobleaching of retinal and naphthacene under the same illumination conditions. The absolute value of  $\psi_{ox}^{max}$  for naphthacene was established in [5]. In our experimental conditions it was 0.8. For retinal we have obtained  $\psi_{ox}^{max} = 0.16$ . Thus,  $k_{ox} = 0.2 \times k_q = 7.5 \times 10^5 M^{-1} s^{-1}$ . It may be concluded therefore that retinals are mainly physical quenchers of  $^1O_2$ ; however, the relative efficiency of the chemical quenching is rather high. We have not attempted to establish the role of chemical quenching in general quenching effect of lipids. However, a comparison with the data in [12] demonstrates that the obtained values of  $k_q$  exceed those of  $k_{ox}$  for methyl esters of oleic, linoleic and linolenic acids by  $\leq 3$ -fold.

Thus, these data support the assumption that free retinal which is known to be always present in the photoreceptor cells might be an effective generator of  $^1O_2$  in the retina. The results of experiments on luminescence quenching show that  $^1O_2$  can attack at least three components of the photoreceptor membranes, i.e., phospholipids, retinal and  $\alpha$ -tocopherol. The molar ratio of these components in the membranes is equal to 1000: <<10:1 [13], assuming that the amount of free retinal in the membrane is much

lower than that of rhodopsin. Taking into account the values of  $k_q$ , we have to assume that the proportion between the rates of  $^1O_2$  interaction with the above components corresponds to 10: <<1:10. This means that  $^1O_2$  interacts mainly with  $\alpha$ -tocopherol and phospholipids, whereas the destruction rate of the retinal is extremely low. Hence  $\alpha$ -tocopherol and phospholipids play the role of a protector of the retinal in the photoreceptor membrane. In this connection it is noteworthy that illumination of high intensity results in accumulation of lipid peroxides in the retinal photoreceptors, the effect being enhanced in vitamin E-deficient animals [2,3].

### Acknowledgements

We are grateful to Drs B. I. Mitsker and N. A. Sokolova for supply of the retinals and to Dr V. P. Karagodin for his valuable assistance in isolating and purifying of the lipids.

### References

- [1] Noell, W. K., Walker, V. S., Kang, B. S. and Berman, S. (1966) *Invest. Ophthalmol.* 5, 450–471.
- [2] Kagan, V. E., Shvedova, A. A., Novikov, K. N. and Kozlov, Yu. P. (1973) *Biochim. Biophys. Acta* 330, 76–79.
- [3] Kagan, V. E., Barybina, G. V. and Novikov, K. N. (1977) *Bull. Exptl. Biol. Med. (Russian)* 4, 411–413.
- [4] Delmelle, M. (1978) *Photochem. Photobiol.* 27, 731–734.
- [5] Krasnovsky, A. A., jr and Kagan, V. E. (1978) *Dokl. Acad. Nauk SSSR (Russian)* 242, 229–232.
- [6] Krasnovsky, A. A., jr (1979) *Photochem. Photobiol.* 29, 29–36.
- [7] Folch, J., Lees, M., Stanley, A. H. S., Meath, S. A. and Le Bason, F. N. (1951) *J. Biol. Chem.* 191, 883–891.
- [8] Young, R. N. and Martin, R. L. (1972) *J. Am. Chem. Soc.* 94, 5183–5185.
- [9] Stevens, B., Perez, S. R. and Ors, J. A. (1974) *J. Am. Chem. Soc.* 96, 6846–6850.
- [10] Farmilo, A. and Wilkinson, F. (1973) *Photochem. Photobiol.* 18, 447–450.
- [11] Stevens, B., Small, R. D., jr and Perez, S. K. (1974) *Photochem. Photobiol.* 20, 505–512.
- [12] Matsushita, S., Terao, J. and Yamauchi, R. (1978) in: *Tocopherol, oxygen and biomembranes* (de Duve, C. and Hayaishi, O. eds) pp. 23–39, Elsevier/North-Holland, Amsterdam, New York.
- [13] Daemen, F. J. M. (1973) *Biochim. Biophys. Acta* 300, 255–288.