

ADAPTATION TO SALT OF THE PHOTOSYNTHETIC APPARATUS IN CYANOBACTERIA

Elisha TEL-OR

Institute for Desert Research, Ben-Gurion University of the Negev, Sede Boqer, Israel

Received 6 November 1979

Revised version received 19 December 1979

1. Introduction

Cyanobacteria grow under a wide range of environmental conditions. Their distribution in environments of highly diversified osmotica is well documented [1]. Hence, the photosynthetic apparatus must adapt to the different degrees of salinity in the habitat. This is achieved by means of either osmoregulation or on the level of the organism [2] or modification of the enzyme structure [3].

Here we correlated the salt tolerance of the whole organism with the salt tolerance of activities of the photosynthetic apparatus and NADPH-cytochrome *c* reductase in 4 cyanobacteria characterized by different degrees of salt tolerance: marine *Spirulina* tolerates up to 3-times sea water [4]; *Calothrix scopulorum*, sea water [5]; *Spirulina platensis*, brackish water [6]; *Nostoc muscorum* 7119, fresh water.

We show that the soluble components of the final steps of photosynthetic electron transport constitute a salt-sensitive site and that salt tolerance may be acquired during growth in saline water.

2. Materials and methods

2.1. Organisms and growth conditions

Spirulina platensis (Cambridge culture collection no. 1475/4a) was grown in Zarouk's medium [8] in a turbidostat culture at 35°C. The culture was stirred by an air stream containing 1.5% CO₂ at 35°C and illuminated by cool white light at 1250 $\mu\text{W}\cdot\text{cm}^{-2}$. Marine *Spirulina* obtained from Dr A. Abielovich, was grown in batch cultures in sea water containing

(g/l): NaNO₃, 2.5; NaHCO₃, 2; FeSO₄, 0.01; Na₂·EDTA, 0.08; Na₂HPO₄, 0.5; and 1 ml of A₅ and B₆ microelements solution [4]. The cultures were grown on a rotary shaker at 30°C and illuminated by cool white light at 250 $\mu\text{W}\cdot\text{cm}^{-2}$. *Nostoc muscorum* 7119 [9] and *Calothrix scopulorum* (Cambridge culture collection no. 1410/5) were grown in the medium of [10] in batch cultures on a rotary shaker at 30°C and illuminated at 250 $\mu\text{W}\cdot\text{cm}^{-2}$.

2.2. Cell-free preparations

Algae were harvested at late logarithmic phase, washed and resuspended in 25 mM *N*-2-hydroxyethyl-piperazine-*N*-2-ethane-sulphonate (Hepes) (pH 7.5) disrupted by sonication and centrifuged (27 000 $\times g$ for 30 min). The supernatant was assayed for reduction of cytochrome *c* and the pellet, resuspended in Hepes buffer, was used for the assay of the photo-reduction of methylviologen mediated by photosystem 1.

Protein was determined according to [11], and chlorophyll extracted in 80% acetone was determined as in [12].

2.3. Assays

Oxygen evolution was measured in cells, which had been washed and resuspended in fresh medium at 30°C with a Clark-type electrode (YSI 4004, Yellow Springs Instr., OH) connected to a recorder. Cells were illuminated at $I=3.5 \text{ mW}\cdot\text{cm}^{-2}$. Methylviologen photoreduction was measured as O₂ consumption with an oxygen electrode. Reduction of cytochrome *c* (horse heart, Sigma) was measured at 550 nm in a Gilford 250 spectrophotometer.

3. Results and discussion

3.1. Effect of salt on photosynthetic electron transport

The response to salt of photosynthetic electron transport *in vivo* was tested by following O_2 evolution in intact filaments of 4 different cyanobacteria in their growth medium (fig.1). The photosynthetic activity of marine *Spirulina* tolerated high salt concentrations with no inhibition up to 2 M NaCl, while *Calothrix* was moderately sensitive and *Nostoc* and *S. platensis* were much more sensitive to NaCl. The photoreduction of methylviologen, mediated by photosystem 1 and supported by reduced dichlorophenolindophenol, was tolerant to NaCl when tested in membrane preparations of *Nostoc*, *Calothrix* and *S. platensis* as shown in fig.2. These results suggest that the membrane-bound components which participate in the photosystem 1 reaction of electron transport are insensitive to NaCl.




Fig.1. Effect of salt on O_2 evolution. Cultures grown as in section 2 were harvested at late logarithmic phase of growth. Cells containing 3–10 μg chl were resuspended in growth medium except for marine *Spirulina* which was resuspended in artificial sea water without NaCl. 100% activity: 82, 206, 138 and 271 $\mu\text{mol } O_2$ evolved $\text{mg} \cdot \text{chl}^{-1} \cdot \text{h}^{-1}$ for marine *Spirulina*, *Calothrix*, *Nostoc* and *Spirulina platensis*, respectively.




Fig.2. Effect of salt on the activity of photosystem 1. Reaction mixture (4 ml) contained 50 mM Hepes (pH 7.5), 5 mM sodium ascorbate, 50 μM dichlorophenolindophenol, 50 μM methylviologen, 2.5 mM sodium azide and membranes containing 20–50 μg chl. 100% activity: 730, 520 and 380 $\mu\text{mol } O_2$ consumed $\cdot \text{mg} \cdot \text{chl}^{-1} \cdot \text{h}^{-1}$ for *Spirulina platensis*, *Nostoc* 7119 and *Calothrix scopulorum*, respectively.




Fig.3. Effect of salt on NADPH–cytochrome *c* reduction. Activity was measured in a 1 ml reaction mixture containing 25 mM Hepes (pH 7.5), 5 mM MgCl_2 , 33 μM NADPH, 33 μM cytochrome *c* and sonicated supernatant cells containing 1–2 mg protein. 100% activity: 3.6, 10.2, 5.0 and 5.5 μmol cytochrome *c* reduced $\cdot \text{mg} \cdot \text{protein}^{-1} \cdot \text{min}^{-1}$ for marine *Spirulina*, *Calothrix scopulorum*, *Nostoc* 7119 and *Spirulina platensis*, respectively.

3.2. The effect of salt on the activity of NADPH-cytochrome *c* reductase

The effect of salt on the reduction of cytochrome *c* by NADPH was tested in the crude soluble fraction of ruptured cells of the 4 different cyanobacteria. As shown in fig.3, only the preparation derived from the marine *Spirulina* exhibited tolerance to NaCl. The reduction of cytochrome *c* involves the formation of a complex between ferredoxin and ferredoxin-NADP reductase [13], which was found to be sensitive to high ionic strength in higher plants [14]. The difference in salt sensitivity between marine *Spirulina* on the one hand and *Nostoc*, *Calothrix* and *S. platensis* on the other, suggests that the enzyme couple in the marine *Spirulina* has become modified in a way which allows the formation of a stable active complex at high ionic strength. The soluble components participating in the final steps of electron transport are therefore a salt-sensitive site.

3.3. Acquired tolerance to salt

We have tested the possibility that cyanobacteria may acquire salt tolerance during their growth in media containing salt. *S. platensis* was cultured in growth media containing either 0.1 M or 0.2 M NaCl.




Fig.4. Effect of salt on O_2 evolution in cultures of *S. platensis* previously grown at different salt concentrations. Cells grown at 0, 0.1 and 0.2 M NaCl in turbidostat cultures at identical conditions were harvested, washed and resuspended in salt-free medium. 100% activity: 469, 450 and 413 $\mu\text{mol } O_2$ evolved $\cdot \text{mg chl}^{-1} \cdot \text{h}^{-1}$ for cultures containing 0, 0.1 and 0.2 M NaCl, respectively.




Fig.5. Effect of salt on NADPH-cytochrome *c* reduction in extracts from cultures previously grown at different salt concentrations. Cells were harvested, washed, disrupted in 25 mM Hepes and the supernatant was assayed as in fig.3. 100% activity: (a) 5.5 and 7.6 μmol cytochrome *c* reduced $\cdot \text{mg protein}^{-1} \cdot \text{min}^{-1}$ for *Spirulina* grown with 0 and 0.2 M NaCl, respectively. (b) 5 and 8.5 μmol cytochrome *c* reduced $\cdot \text{mg protein}^{-1} \cdot \text{min}^{-1}$ for *Nostoc* grown with 0 and 0.2 M NaCl, respectively.

The photosynthetic activity of these cultures exhibited increased tolerance to salt (fig.4). Some acquired tolerance to salt was also found in the activity of NADPH-cytochrome *c* reductase (fig.5a).

Similar trends were also found in cultures of *Nostoc* grown in batch cultures containing either 0.2 M or 0.4 M NaCl. The photosynthetic activity was found to tolerate higher salt concentrations than the control cultures, and likewise the reduction of cytochrome *c* exhibited adaptation to salt (fig.5b). These results suggest that the enzyme couple ferredoxin and ferredoxin-NADP reductase undergoes a specific modification, which confers on it an increased tolerance to increased ionic strength. The reports that ferredoxin-NADP reductase is composed of a mixture of isozymes [13] and the evidence for two ferredoxins in *Nostoc* [15] may provide a possible explanation for this modification during the growth in salt-containing medium; in this medium, the cells may preferably synthesize the isozyme which is more resistant to the high ionic strength.

Acknowledgement

I am grateful to Mr Amos Goldstein for his assistance with the *Spirulina platensis* experiments.

References

- [1] Fogg, G. E., Stewart, W. D. P., Fay, P. and Walsby, A. E. (1973) *The Blue-Green Algae*, Academic Press, London.
- [2] Hellebust, J. A. (1976) *Ann. Rev. Plant Physiol.* 27, 485–505.
- [3] Lanyi, J. K. (1974) *Bact. Rev.* 40, 803–846.
- [4] Abeliovich, A. (1980) submitted.
- [5] Jones, K. and Stewart, W. D. P. (1969) *J. Mar. Biol. Ass. (UK)* 49, 475–488.
- [6] Richmond, A. E. and Vonshak, A. (1978) *Arch. Hydrobiol. Beih.* 11, 280–284.
- [7] Shin, M. (1971) *Methods Enzymol.* 23, 440–447.
- [8] Goldstein, A. (1979) MSc Dissert., Ben-Gurion University of the Negev, Beer-Sheva, (in Hebrew).
- [9] Tel-Or, E., Luijk, L. M. and Packer, L. (1977) *FEBS Lett.* 78, 49–52.
- [10] Allen, M. B. and Arnon, D. I. (1955) *Plant Physiol.* 30, 366–372.
- [11] Lowry, O. H., Rosenbrough, N. J., Farr, A. L. and Randall, B. J. (1951) *J. Biol. Chem.* 193, 265–275.
- [12] Mackinney, G. (1941) *J. Biol. Chem.* 140, 315–322.
- [13] Forti, G. (1977) *Encyclopedia of Plant Physiol.* (Trebst, A. and Avron, M. eds.) vol. 5, pp. 222–226, Springer-Verlag, Berlin.
- [14] Nelson, N. and Neumann, J. (1969) *J. Biol. Chem.* 244, 1926–1931.
- [15] Dutton, J. E. and Rogers, L. J. (1978) *Biochim. Biophys. Acta* 537, 501–506.