

## MAMMALIAN LIPOATE ACETYLTRANSFERASE: MOLECULAR WEIGHT DETERMINATION BY GEL FILTRATION IN THE PRESENCE OF GUANIDINIUM CHLORIDE

Georg-B. KRESZE, Brigitte DIETL and Hedwig RONFT

*Institut für Biochemie der Universität München Karlstrasse 23, D-8000 München 2, FRG*

Received 6 February 1980

### 1. Introduction

Lipoate acetyltransferase (E2) forms the core of the pyruvate dehydrogenase multienzyme complex (reviewed [1,2]). The subunit  $M_r$  of mammalian E2 has been a matter of controversy. Using SDS–polyacrylamide gel electrophoresis  $M_r$  74 000–76 000 has been found for the bovine as well as porcine kidney and heart enzyme [3–6]. The same value was reported for pig heart E2 in a study employing gel filtration and sedimentation equilibrium analysis in the presence of 6 M guanidine–HCl [4]. However, by this latter method, the subunit  $M_r$  of bovine kidney and heart E2 was determined to be ~52 000 [3].

Knowledge of the subunit  $M_r$  of E2 is indispensable to elucidate the molecular structure of the pyruvate dehydrogenase complex. In fact, due to the uncertainty about the E2 subunit, contradictory models have been proposed for the polypeptide chain stoichiometry of the mammalian E2 core as well as the enzyme complex [3–5]. We therefore saw fit to redetermine the subunit  $M_r$  of bovine kidney E2 by gel filtration using Sephacryl S-300 as the gel matrix.

### 2. Experimental

#### 2.1. Materials

Pyruvate dehydrogenase complex was purified from bovine kidney and E2 isolated from the complex

as in [6]. Its papain fragment was obtained as in [7]. Sephacryl S-300 and blue dextran-2000 were from Pharmacia, Uppsala. Guanidine–HCl was from Merck, Darmstadt, or Sigma, St Louis, and was purified according to [8]. Cytochrome *c*, aldolase, lactate dehydrogenase, and phosphorylase *a* were from Boehringer, Mannheim, chymotrypsinogen and DNP-Ala from Serva, Heidelberg, and bovine serum albumin from Behringwerke, Marburg. Iodo[1-<sup>14</sup>C]acetamide (57 Ci/mol) and [<sup>3</sup>H]NSP (43 Ci/mmol) were from the Radiochemical Centre, Amersham.

#### 2.2. Methods

The calibrating proteins as well as E2 were radio-labelled either with iodo[1-<sup>14</sup>C]acetamide according to [8] or were treated first with [<sup>3</sup>H]NSP at pH 8.5 as in [9] and then with unlabelled iodoacetamide in the presence of 6 M guanidine–HCl [8].

Gel filtration was performed according to [10] but using Sephacryl S-300 superfine instead of S-200 (gel bed dimensions, 1.6 × 83 cm). The elution buffer was 6 M guanidine–HCl (pH 5.0) (flowrate 4.8 ml/h). Fractions of ~0.88 ml (11 min) were collected. To improve reproducibility, the elution positions were determined by weight rather than by volume [11]. Blue dextran and DNP-Ala were used as the markers for the void and total volume, respectively, and were localized by measuring  $A_{630}$  and  $A_{360}$ . To determine the elution positions of the labelled proteins, samples of each fraction were counted for radioactivity, except for cytochrome *c* which was monitored at 410 nm. To obtain a calibration curve,  $K_d$  values were calculated according to [10] and plotted according to [12] or [13].

SDS–polyacrylamide gel electrophoresis was per-

**Abbreviations:** E2, lipoate acetyltransferase (EC 2.3.1.12);  $M_r$ , molecular weight; SDS, sodium dodecylsulfate; DNP-Ala, dinitrophenyl-alanine; [<sup>3</sup>H]NSP, *N*-succinimidyl-[2,3-<sup>3</sup>H]-propionate

formed as in [6]. Localization of radioactivity in the Coomassie blue-stained gels was accomplished according to [14].

### 3. Results

#### 3.1. Gel filtration

In our experiments, Sephacryl S-300 turned out to be a very suitable gel matrix for  $M_r$  determination of proteins in the presence of 6 M guanidine-HCl. Good linearity was obtained in the range from at least 11 000–93 000  $M_r$  when the data were plotted either according to [12] ( $r^2 = 0.998$ , fig.1) or [13] ( $r^2 = 0.989$ , not shown).

Bovine kidney E2 was eluted in a position corresponding to  $M_r$  51 100  $\pm$  1100 (or 53 600  $\pm$  1200 when evaluated according to [13]). This value is in excellent agreement with the result of sedimentation equilibrium analysis [3] but disagrees with the app.  $M_r$  in SDS gel electrophoresis.




Fig.1. Plot of  $\log 100 K_d$  vs  $N^{2/3}$  [12] for the calibrating proteins as well as (I) E2, (II) the fragment formed from E2 during incubation with 6 M urea, and (III) the papain fragment of E2.  $N$  is the number of amino acid residues in the polypeptide chain. The following values were used for  $N$  (in parentheses, no.  $K_d$  determinations): phosphorylase  $a$ , 797 (3); bovine serum albumin (BSA), 579 (3); aldolase, 358 (3); lactate dehydrogenase, 315 (1); chymotrypsinogen, 245 (8); cytochrome  $c$ , 104 (12).  $K_d$  values found for the test proteins were: E2,  $0.171 \pm 0.004$  (7); urea fragment,  $0.271 \pm 0.006$  (5); papain fragment,  $0.288 \pm 0.008$  (3).

The incorporation of radioactivity from [ $^3$ H]NSP into E2 was low ( $\sim 3 \times 10^7$  cpm/mg, corresponding to  $\sim 0.05$  mol/mol E2). We therefore tried to increase the amount of label by adding 6 M urea to the incubation mixture of E2 and [ $^3$ H]NSP. However, in this case, only part of the label was eluted at 52 000  $M_r$  whereas  $\sim 50\%$  of the radioactivity gave a peak corresponding to  $M_r$  30 400  $\pm$  900 (II in fig.1) which was quite insignificant (if at all detectable) in E2 samples labelled in the absence of urea. It appears that isolated E2 contains small amounts of a firmly bound protease which is activated in the presence of urea. The size of the fragment detected in these experiments is very similar to that of the principal stable fragment obtained from E2 after treatment with a lysosomal protease ( $M_r$  29 200  $\pm$  600 in SDS gel electrophoresis) [6].

We have reported [7] that after limited proteolysis of E2 with papain, the principal fragment detectable in polyacrylamide gels after staining with Coomassie blue has app.  $M_r$   $\sim 26$  000. When this fragment was analyzed by gel filtration, it was found to have  $M_r$  28 000  $\pm$  1200 (III in fig.1). This suggests that the papain fragment, unlike uncleaved E2, migrates normally in SDS gel electrophoresis and thus, may lack the region of the E2 chain responsible for its anomalous behaviour.

#### 3.2. SDS-polyacrylamide gel electrophoresis

The following experiment was done to exclude the possibility that E2, when eluted from the column in a position corresponding to  $M_r$   $\sim 52$  000 was itself degraded proteolytically. The labelled material of the E2 peak was pooled, dialyzed extensively against water to remove the guanidine-HCl, freeze-dried and dissolved in the sample buffer for SDS gel electrophoresis. Unlabelled pyruvate dehydrogenase complex was added as a carrier and molecular weight standard. After the run, the gel was stained and then sliced. The pattern of gel staining and the distribution of radioactivity among the slices are shown in fig.2. Nearly all of the radioactivity was found coinciding with the E2 band whose position corresponds to app.  $M_r$  74 000. Thus, E2 was not degraded by proteases. Since the same sample had shown  $M_r$  52 000 in gel filtration, the anomalous behaviour of E2 is clearly established.

### 4. Discussion

A number of proteins have been reported to display




Fig.2. SDS-polyacrylamide gel electrophoresis of E2 after elution from the column. The stained bands represent, from top to bottom: E2 ( $M_r$  in SDS gel electrophoresis, 74 000); lipoamide dehydrogenase (E3) ( $M_r$  56 000); pyruvate dehydrogenase (E1),  $\alpha$  subunit ( $M_r$  41 000) and  $\beta$  subunit ( $M_r$  36 000). The gel was sliced as indicated on the left side of the scale and the radioactivity contained in the slices was counted. Total radioactivity (sum of all slices) was 5828 cpm.

anomalous migration in SDS gel electrophoresis (e.g. [15–17]). In contrast, anomalous behaviour of proteins in gel filtration with 6 M guanidine-HCl has not yet been reported [18]. Since, furthermore,  $M_r$  ~52 000 has now been found for the subunit of mammalian E2 by two independent methods, this value must be considered correct rather than  $M_r$  74 000 resulting from SDS gel electrophoresis.

Two models have been proposed for the structure of the mammalian E2 core:

- (i) A 60 subunit pentagonal dodecahedron with  $M_r$  52 000 subunits [3];
- (ii) A 24 subunit aggregate with subunits of  $M_r$  74 000 [4,5].

The results reported here argue against the latter proposal. Furthermore, only a 60-subunit structure is in agreement with the symmetry of the E2 core as seen in electron micrographs [19,20].

$M_r$  of the undissociated core has been determined to be  $\sim 3.1 \times 10^6$  ([3], bovine kidney and heart enzymes) or  $1.8 \times 10^6$  ([4], pig heart enzyme). Since species differences between the bovine and porcine

enzymes are very unlikely to cause such discrepancies, an obvious explanation suggested by our results would be that, in the latter study, E2 was fragmented by an endogenous protease to give a core consisting of  $M_r$  60 000 ~ 30 000 subunits.

The reasons for the anomalous migration of kidney E2 in SDS gel electrophoresis may be similar to those discussed for *Escherichia coli* E2 [21]. This question is under study.

### Acknowledgement

This work was supported by the Deutsche Forschungsgemeinschaft.

### References

- [1] Reed, L. J. (1974) *Acc. Chem. Res.* 7, 40–46.
- [2] Koike, M. and Koike, K. (1976) *Adv. Biophys.* 9, 187–227.
- [3] Barrera, C. R., Namihira, G., Hamilton, L., Munk, P., Eley, M. H., Linn, T. C. and Reed, L. J. (1972) *Arch. Biochem. Biophys.* 148, 343–358.
- [4] Hamada, M., Otsuka, K.-I., Tanaka, N., Ogasahara, K., Koike, K., Hiraoka, T. and Koike, M. (1975) *J. Biochem.* 78, 187–197.
- [5] Sugden, P. H. and Randle, P. J. (1978) *Biochem. J.* 173, 659–668.
- [6] Kresze, G.-B. and Steber, L. (1979) *Eur. J. Biochem.* 95, 569–578.
- [7] Kresze, G.-B., Ronft, H. and Dietl, B. (1980) *Eur. J. Biochem.* in press.
- [8] Fohlman, J., Eaker, D., Karlsson, E. and Thesleff, S. (1976) *Eur. J. Biochem.* 68, 457–469.
- [9] Bolton, A. E. and Hunter, W. M. (1973) *Biochem. J.* 133, 529–539.
- [10] Belew, M., Fohlman, J. and Janson, J.-C. (1978) *FEBS Lett.* 91, 302–304.
- [11] Fish, W. W., Mann, K. G. and Tanford, C. (1969) *J. Biol. Chem.* 244, 4989–4994.
- [12] Rydén, L. (1971) *FEBS Lett.* 18, 321–325.
- [13] Porath, J. (1963) *Pure Appl. Chem.* 6, 233–244.
- [14] Bates, D. L., Harrison, R. A. and Perham, R. N. (1975) *FEBS Lett.* 60, 427–430.
- [15] Mitchell, E. D., Riquetti, P., Loring, R. H. and Carraway, K. L. (1973) *Biochim. Biophys. Acta* 295, 314–322.
- [16] Weber, K. and Osborn, M. (1975) in: *The Proteins* (Neurath, H. and Hill, R. L. eds) 3rd edn, vol. 1, pp. 179–223, Academic Press, New York.
- [17] Lehtovaara, P. (1978) *Biochem. J.* 169, 251–253.
- [18] Mann, K. G. and Fish, W. W. (1972) *Methods Enzymol.* 26, 28–42.
- [19] Reed, L. J. and Oliver, R. M. (1968) *Brookhaven Symp. Biol.* 21, 397–412.
- [20] Junger, E. and Reinauer, H. (1972) *Biochim. Biophys. Acta* 250, 478–490.
- [21] Bleile, D. M., Munk, P., Oliver, R. M. and Reed, L. J. (1979) *Proc. Natl. Acad. Sci. USA* 76, 4385–4389.