

5'-NUCLEOTIDASE ACTIVITY OF TWO POPULATIONS OF MOUSE THYMOCYTES SEPARATED BY PEANUT AGGLUTININ AGGLUTINATION

Jacques DORNAND, Jean-Claude BONNAFOUS and Jean-Claude MANI

*Laboratoire de Biochimie des Membranes, ER CNRS no. 228, Ecole Nationale Supérieure de Chimie,
8 rue de l'Ecole Normale, F-34075 Montpellier, France*

Received 18 July 1980

1. Introduction

The adenosine-generating enzyme 5'-nucleotidase is localized on the external surface of the plasma membrane of many cells [1], including lymphocytes [2,3]. This enzyme is a concanavalin A (con A) receptor [4] and is inhibited by the lectin binding [3,5]. Although little is known about its function, 5'-nucleotidase plays a critical role in lymphocyte adenosine metabolism; it is involved in adenosine uptake from 5'-AMP [6,7] and regulates intracellular cyclic AMP level [8] through membrane adenosine receptor sites [9]. As adenosine appears to exert a negative control on lymphocyte proliferation [10,11] the distribution of 5'-nucleotidase activity among lymphocyte subpopulations is a very important problem. Non-uniform distribution of 5'-nucleotidase among lymphoid tissues has been described [12–17] but these results are sometimes contradictory; moreover completely different distributions are reported for different animal species.

However low 5'-nucleotidase levels are unambiguously found in lymphocytes from patients with chronic lymphocytic leukemia (CLL) ([17–20] and ourselves) or hypogammaglobulinemia [21,22] and in cord blood lymphocytes [22,23]. As CLL cells are thought to be arrested at an early stage of development [24] and as cord blood cells might also be immature [25], it was hypothesized that the 5'-nucleotidase deficiency in hypogammaglobulinemic patients may reflect a stage of maturation arrest in T and B lymphocytes [23].

As others [13,14,16] we found that 5'-nucleotidase activity in thymocytes is 6–10-fold lower than in splenocytes. We checked the hypothesis that this low activity may reflect the fact that the major thymo-

cyte subpopulation (85–95%) which is hydrocortisone-sensitive and immature displays very low 5'-nucleotidase activity.

2. Materials and methods

Thymocytes were obtained from 4–7-week-old male mice (C57/BL6, DBA2, C3H/eb, CB-20 or Swiss) killed by cervical luxation. Thymuses were removed, minced in Hank's salt and passed through a fine stainless steel mesh to give a single cell suspension ($1\text{--}1.5 \times 10^8$ cells/thymus); when necessary erythrocytes were selectively killed by 0.14 M ammonium chloride. Cell viability was 95–98%. Hydrocortisone-resistant thymocytes were obtained by the same procedure 48 h after intraperitoneal injection of 2.5 mg hydrocortisone acetate. Splenocytes were prepared as in [7].

Mature and immature thymocyte populations were isolated by agglutination with peanut agglutinin (PNA) as in [26]. Thymocytes, 2×10^8 in 0.25 ml PBS, were incubated with 0.25 mg PNA in 0.25 ml PBS for 10 min at 25°C; the mixture was then layered on 8 ml FCS 30% in PBS. After 30 min sedimentation ($1 \times g$) agglutinated (immature) cells (bottom layer) were removed from non-agglutinated (mature) cells (top layer), dissociated into single cells by 10 min incubation at 37°C with 5 ml 0.2 M D-galactose in PBS, washed twice with 5 ml 0.2 M D-galactose and twice with PBS. Unagglutinated cells (5–10% of total thymocytes) received the same treatment. Viability of both populations was 95%.

5'-Nucleotidase activity was determined using 5'-[^{32}P]AMP as substrate. Cells ($2.5\text{--}5 \times 10^6$) were incubated 30 min at 37°C in 250 μl 199 Hepes

medium with 0.2 mM 5'-[32 P]AMP. The reaction was stopped with 150 μ l 0.1 N HCl and 32 P_i was determined as in [3]. The contribution of non specific phosphatases to 5'-AMP hydrolysis was negligible [3] and increases in P_i up to 0.1 M had no effect on the enzyme activity [1,8]. 5'-Nucleotidase activity was expressed as nmol P_i · h⁻¹ · mg protein⁻¹. The amount of proteins corresponding to 2.5–5 × 10⁶ cells was determined by Lowry's method.

Thymidine uptakes induced by PHA Difco (7 μ l/ml) or con A Pharmacia (5 μ g/ml) were determined as in [11]. Cells 250 μ l (3 × 10⁶/ml) were cultured in RPMI medium supplemented with 15% fetal calf serum for 60 h at 37°C under an air–CO₂ (95:5) atmosphere. For the last 5 h the cells were pulsed with [3 H]thymidine and harvested as in [11].

3. Results

The separation of mouse thymocytes into two subpopulations (mature and immature) was done by the method in [26] for mouse thymocytes and recently applied to human thymocytes [27]. This method is based on the observation that immature thymocytes bind PNA while mature cells do not. Agglutinated cells (80–90% of total thymocytes) were dissociated into single cells by incubation with

D-galactose which bind specifically PNA; this population was unambiguously characterized as immature and hydrocortisone-sensitive, while non-agglutinated cells were found identical to mature hydrocortisone-resistant thymocytes [26,27]. Table 1 confirms these characterizations; it reports the stimulation of unseparated thymocytes and thymocyte subpopulations by PHA or con A. PNA⁺ (agglutinated) lymphocytes were not stimulated by PHA, while PNA⁻ lymphocytes and thymocytes from hydrocortisone-treated mice were highly stimulated.

Table 2 shows the 5'-nucleotidase activity of unseparated, PNA⁺ and PNA⁻ thymocytes. Ten experiments were performed with Swiss mouse thymocytes and one with each of the other mouse strains. The percentage of PNA⁻ thymocytes was 5–15%; the 5'-nucleotidase activity of this subpopulation was 3–5-fold higher than that of unseparated thymocytes. PNA⁺ thymocytes accounted for 80–90% of total thymocytes and displayed a 5'-nucleotidase activity 1.5–2-fold lower than unseparated cells. 5'-Nucleotidase activity of PNA⁻ thymocytes was thus 5–11-fold higher than that of PNA⁺ cells. PNA had no effect on pure 5'-nucleotidase from pig lymph node lymphocytes [5] (not shown). Thymocyte 5'-nucleotidase was neither affected by incubation with 0.2 M D-galactose for 10–60 min, nor by preincubation with PNA (≤ 50 μ g/

Table 1
Mitogenic response of mouse thymocytes to Con A and PHA

Mitogen	Unseparated thymocytes	PNA ⁺ thymocytes	PNA ⁻ thymocytes	Hydrocortisone-resistant thymocytes
None	600 ± 200	520 ± 50	900 ± 300	650 ± 60
Con A	93 000 ± 7000	23 000 ± 3000	210 000 ± 15 000	80 000 ± 10 000
PHA	1500 ± 300	600 ± 200	80 000 ± 6000	63 000 ± 4000

[3 H]Thymidine incorporation (cpm) in mouse thymocyte populations, 60 h after stimulation by PHA or con A

Table 2
5'-Nucleotidase activities (nmol P_i · h⁻¹ · mg protein⁻¹) of thymocytes (unseparated, PNA⁺, PNA⁻) from different mouse strains

	C3H/eb	CB20	C57BL6	DBA2	Swiss
Unseparated thymocytes	30	40		30	15–35
PNA ⁺ thymocytes	18	25	11.2	20	8.3–25
PNA ⁻ thymocytes	130	130	125	150	90–210

Table 3
Effect of hydrocortisone in vivo on the 5'-nucleotidase activity of mouse thymocytes

	Expt 1	Expt 2	Expt 3
Thymocytes from hydrocortisone-treated mice	194	144	162
Unseparated thymocytes from untreated mice	30	35	30
PNA ⁺ thymocytes from untreated mice	20	17.5	18
PNA ⁻ thymocytes from untreated mice	110	133	130

5'-Nucleotidase activities (nmol $P_i \cdot h^{-1} \cdot mg$ protein⁻¹) of thymocytes from hydrocortisone-treated Swiss mice and of control thymocytes (unseparated, PNA⁺, PNA⁻) from untreated Swiss mice

2.5×10^6 cells), which ruled out the possibility that the low 5'-nucleotidase level of PNA⁺ thymocytes was due to D-galactose treatment or to inhibition by PNA incompletely removed from the cell membrane.

To study mature thymocytes, we used hydrocortisone treatment to deplete the immature cell population [26]. Table 3 represents the 5'-nucleotidase activity of Swiss mouse thymocytes, two days after an intraperitoneal injection of 2.5 mg hydrocortisone acetate. Hydrocortisone-resistant cells were pooled from 10 mice ($\sim 15 \times 10^6$ cells/thymus); they did not agglutinate with PNA. Their 5'-nucleotidase activity was compared with that of thymocytes from untreated mice (unseparated, PNA⁺, PNA⁻). 5'-Nucleotidase activity of hydrocortisone-resistant cells was 5- and 8–10-fold higher than that of unseparated and PNA⁻ control thymocytes, respectively. It appeared that PNA⁻ control thymocytes and hydrocortisone-resistant thymocytes, which have been claimed to be identical (mature) population [26] had similar 5'-nucleotidase activities.

5'-Nucleotidase activity of splenic lymphocytes from Swiss mice was determined in 10 different expt; its mean value was 320 ± 70 nmol $P_i \cdot h^{-1} \cdot mg^{-1}$. As no difference between T and B splenocyte 5'-nucleotidase activity was evidenced [14], it appeared that PNA⁻ or hydrocortisone-resistant thymocytes had lower 5'-nucleotidase activity than T splenocytes.

4. Discussion

Using the general technique in [28] we separated

mouse thymocytes into PNA⁻ thymocytes identified as mature cells (medullary) and PNA⁺ thymocytes which are immature (cortical). As it was shown [26,27] that PNA agglutination and subsequent dispersion with galactose yielded fully functional cells, and as these treatments had no effect on 5'-nucleotidase activity, we can state that the low 5'-nucleotidase activity of mouse thymocytes results from the presence of 90–95% immature cells. In mouse thymus immature cells can be selectively eliminated by hydrocortisone treatment, the remaining mature cells had the same high 5'-nucleotidase activity than PNA⁻ control thymocytes. These repeatedly obtained results confirmed that mature thymocyte 5'-nucleotidase activity is 10-fold higher than that of immature thymocytes. No difference for 5'-nucleotidase activity were found in thymocytes from hydrocortisone-treated rats or untreated control rats (410 ± 105 and 370 ± 110 nmol $P_i \cdot h^{-1} \cdot mg^{-1}$, respectively) [15]; however it must be emphasized that the 5'-nucleotidase value reported for control thymocytes was abnormally high as compared to rat splenocyte 5'-nucleotidase (890 nmol $P_i \cdot h^{-1} \cdot mg^{-1}$), as generally the thymocyte activity is 6–10-fold lower ([13,14] and ourselves) or even more in rabbit thymocytes [16].

5'-Nucleotidase activity in PNA⁻ mature thymocytes appeared lower than in T splenocytes, if we assume that T and B splenocytes display the same mean activity [14]. This result can be explained if thymus maturation generates several subclasses of T-cells [29] with different 5'-nucleotidase activity and if PNA⁻ thymocyte subpopulation is heterogeneous [30]. On the other hand a small subpopulation of mature lymphocytes bearing PNA receptors is mostly found in the thymus [29] and could account for the 5'-nucleotidase activity of PNA⁺ thymocytes.

5'-Nucleotidase activity appears to be restricted to hydrocortisone-resistant thymocytes. It was recently shown that the steroid-resistant intrathymic pool is a resident subpopulation which is not exported to the periphery [31]. We do not know if 5'-nucleotidase increase in peripheral T cells occurs after migration from the thymus to secondary lymphoid tissues or before.

Recent reports on the low 5'-nucleotidase activity in lymphocytes from patients with CLL [17–20] or hypogammaglobulinemia [21,22] increase the interest of studying the function and distribution of this enzyme among various lymphocyte populations.

T and B cells from cord blood have a lower activity than adult lymphocytes [22,23]. Malignant transformed cells display also very low activity [32]. Most of these cells are thought to be arrested at an early stage of maturation [24,25]. As we found a low 5'-nucleotidase activity in immature thymocytes, the absence of 5'-nucleotidase activity might be considered as a marker of a selective maturation arrest. This was confirmed by the absence of detectable 5'-nucleotidase in new-born cat liver cells and in cells from regenerating liver [33].

Acknowledgements

This work received financial support from the Centre National de la Recherche Scientifique and the Fondation pour la Recherche Médicale Française. The authors acknowledge the skillful technical assistance of R. Larguier and N. Bernad.

References

- [1] De Pierre, J. W. and Karnovsky, M. L. (1974) *J. Biol. Chem.* 249, 7111–7120.
- [2] Misra, D. N., Gill, T. J. iii and Estes, L. W. (1974) *Biochim. Biophys. Acta* 352, 455–461.
- [3] Dornand, J., Reminiac, C. and Mani, J. C. (1977) *Biochimie* 59, 425–432.
- [4] Dornand, J., Bonnafous, J. C. and Mani, J. C. (1978) *Biochem. Biophys. Res. Commun.* 82, 685–692.
- [5] Dornand, J., Bonnafous, J. C. and Mani, J. C. (1978) *Eur. J. Biochem.* 87, 459–465.
- [6] Fleit, H., Conklyn, M., Stebbins, R. D. and Silber, R. (1975) *J. Biol. Chem.* 250, 8889–8892.
- [7] Dornand, J., Bonnafous, J. C., Gavach, C. and Mani, J. C. (1979) *Biochimie* 61, 973–977.
- [8] Dornand, J., Bonnafous, J. C. and Mani, J. C. (1980) *FEBS Lett.* 110, 30–34.
- [9] Bonnafous, J. C., Dornand, J. and Mani, J. C. (1979) *FEBS Lett.* 107, 95–99.
- [10] Hovi, T., Smyth, J. F., Allison, A. C. and Williams, S. C. (1976) *Clin. Exp. Immunol.* 23, 395–403.
- [11] Bonnafous, J. C., Dornand, J. and Mani, J. C. (1979) *Biochim. Biophys. Acta* 587, 180–191.
- [12] Allan, D. and Crumpton, M. J. (1970) *Biochem. J.* 120, 133–143.
- [13] Misra, D. N., Ladoulis, C. T., Estes, L. W., Gill, T. J. iii (1975) *Biochemistry* 14, 3014–3024.
- [14] Uusitalo, R. J. and Karnovsky, M. J. (1977) *J. Histochem. Cytochem.* 25, 87–96.
- [15] Barton, R. W. and Goldschneider, I. (1979) *J. Immunol.* 121, 2329–2334.
- [16] Gregory, S. H. and Kern, M. (1979) *J. Immunol.* 123, 1078–1082.
- [17] Rowe, M., De Gast, G. C., Platts-Mills, T. A. E., Asherson, G. L., Webster, A. D. B. and Johnson, S. M. (1979) *Clin. Exp. Immunol.* 96, 97–101.
- [18] Quagliata, F., Faig, D., Conklyn, M. and Silber, R. (1974) *Cancer Res.* 34, 3197–3202.
- [19] Kramers, M. T. C., Catovsky, D., Foa, R., Cherchi, M. and Galton, D. A. B. (1976) *Biomedicine* 25, 363–365.
- [20] Petitou, M., Tuy, F., Rosenfeld, C., Mishal, Z., Paintrand, M., Jasmin, C., Mathé, G. and Inbar, M. (1978) *Proc. Natl. Acad. Sci. USA* 75, 2306–2310.
- [21] Johnson, S. M., North, M. E., Asherson, G. L., Allsop, J., Watts, R. W. and Webster, A. D. B. (1977) *Lancet* i, 168–170.
- [22] Rowe, M., De Gast, G. C., Platts-Mills, T. A. E., Asherson, G. C., Webster, A. D. B. and Johnson, S. M. (1980) *Clin. Exp. Immunol.* 39, 337–343.
- [23] Kramers, M. T. C., Catovsky, D., Cherchi, M. and Galton, D. A. B. (1977) *Leukemia Res.* 1, 279–281.
- [24] Salmon, S. E. and Seligmann, M. (1974) *Lancet* ii, 1230–1233.
- [25] Hayward, A. R. and Lawton, A. R. (1977) *J. Immunol.* 119, 1213–1217.
- [26] Reisner, Y., Israeli-Linker, M. and Sharon, N. (1976) *Cell. Immunol.* 25, 129–134.
- [27] Reisner, Y., Biniaminov, E. R., Sharon, N. and Ramot, B. (1979) *Proc. Natl. Acad. Sci. USA* 76, 447–451.
- [28] Sharon, N. (1979) in: *Affinity Chromatography and Molecular Interactions* (Egly, J. M. ed) pp. 197–206, INSERM edn, Paris.
- [29] London, J. and Horton, M. A. (1980) *J. Immunol.* 124, 1803–1807.
- [30] London, J., Petit, M. and Horton, M. A. (1979) in: *Protides of Biological Fluids, Proc. 27th Colloq. Brussels* (Peters, H. ed) Pergamon, Oxford.
- [31] Stutman, O. (1978) *Immunol. Rev.* 42, 138–184.
- [32] Raz, A., Collard, J. G. and Inbar, M. (1978) *Cancer Res.* 38, 1258–1262.
- [33] Koizumi, K., Ito, Y., Kojima, K. and Fujii, T. (1976) *Biochem. J.* 79, 739–748.