

Glycine and asparagine tRNA sequences from the archaeobacterium, *Methanobacterium thermoautotrophicum*

Xiang-Rong Gu*, K. Nicoghossian and R.J. Cedergren⁺

Département de Biochimie, Université de Montréal, Montréal, Québec H3C 3J7, Canada

Received 7 August 1984

Two tRNA sequences from *Methanobacterium thermoautotrophicum* are reported. Both tRNA^{Gly}_{GCC} and tRNA^{Asn}_{NUU}, the first tRNA sequences from methanogens, were determined by partial hydrolyses (both chemical and enzymatic) and analyzed by gel electrophoresis. The two tRNAs contain the unusual T-loop modifications, Cm and m¹I, which are present in other archaeobacterial tRNAs. Finally the presence of an unknown modification in the D-loop has been inferred by a large jump in the sequence ladder. These tRNAs are approximately equidistant from eubacterial or eukaryotic tRNAs.

tRNA sequence Evolution Methanogen

1. INTRODUCTION

The kingdom of archaeobacteria is composed of a diverse set of organisms including thermoacidophiles, halophiles and methanogens [1,2]. Curiously, halophiles have enjoyed an extensive popularity even though they represent only a relatively small niche within the much larger grouping of methanogens. In our desire to use tRNA sequences as a probe of evolutionary relationships, we have turned our attention to the methanogens [3,4]. To date no tRNA sequences from methanogens have been reported, although a tRNA gene sequence has been published as part of a study of ribosomal RNA genes [5]. The two tRNA sequences determined here, tRNA^{Asn}_{NUU} and tRNA^{Gly}_{GCC} from *Methanobacterium thermoautotrophicum* are compared with other tRNA structures.

2. METHODS

2.1. Purification of the tRNAs

Cells of *M. thermoautotrophicum* were generously supplied by Dr R.S. Wolfe. Crude tRNA was isolated by phenol extraction of cells and then purified by gel electrophoresis [3]: 20 A₂₆₀ units were applied to a 10% polyacrylamide-7 M urea gel which was subjected to electrophoresis at 600 V for 48 h at 4°C in TBE buffer (50 mM Tris-borate and 5 mM EDTA). The tRNA was resolved into 13 main bands, which were isolated and further purified by electrophoresis in a 20% gel at room temperature (1000 V). Pure tRNA contained in the bands was extracted from the gel in a 0.5 M NH₄ acetate, 1 mM MgCl₂, 0.1 mM EDTA and 0.1% SDS solution overnight at room temperature, precipitated twice with ethanol and stored dry at -20°C.

2.2. Preparation and sequencing of ³²P-labelled tRNA

Labelling of tRNAs was carried out as in [6] and [7]. End-labelled tRNAs were electrophoresed for 15 h at 1500 V in thin (0.08 cm) 15% polyacrylamide-7 M urea gels. Under these conditions one

* Present address: Institute of Biochemistry, 320 Yoyang Road, Shanghai, China

⁺ To whom correspondence should be addressed

main band for each of the different tRNAs (purified as above) was observed on the gel. tRNA sequencing was carried out by enzymatic hydrolysis of the 3'- or 5'-end labelled tRNA with RNases T₁, U₂, A, and Phy-M [7]. A variation of the technique in [8] was also used to identify and position modified nucleotides. In this latter technique, 1 μ g tRNA was partially hydrolyzed in 5 μ l deionized formamide at 100°C for 3–4 min. The resulting fragments were labelled with 100–200 μ Ci [γ -³²P]-ATP (3000 Ci/mmol) using 5 units polynucleotide kinase and separated on 15 and 20% polyacrylamide-7 M urea gels at room temperature in TBE buffer. The individual bands were excised from the gel and extracted with 300 μ l of 300 mM NaCl, 0.1% SDS. To the spin filtrate, 2.5 vols of ethanol were added to precipitate the RNA. Digestion of labelled RNA fragments in 5 μ g carrier tRNA was carried out with 0.5 units RNase T₂, 1 unit RNase T₁ and 0.5 μ g RNase A in 100 mM Na acetate (pH 4.5) overnight at 37°C. The 5'-labelled nucleoside 3',5'-diphosphates were identified by thin-layer chromatography on PEI-cellulose sheets developed in 0.55 M ammonium sulfate at 4°C or in 1.75 M ammonium formate (pH 3.5) at room temperature.

3. RESULTS

3.1. Primary sequence

The primary sequence was elucidated using rapid gel sequencing techniques. In most cases, fragments of the full molecule were essential to the final assignment of sequence. The variable loop region is particularly recalcitrant to structure determination, yet this region is readily available using a labelled fragment. Autoradiograms of typical sequencing gels are shown in figs 1,2. Fig.1 is the enzymatic degradation of the 5'-end labelled tRNA^{Gly} whereas fig.2 is the enzymatic sequencing gel of a 5'-end labelled fragment of tRNA^{Gly} covering A₉–G₁₉. This D-loop region was particularly difficult to determine because of an unknown modified nucleotide at position 15. Our preliminary sequence of this region did not take into account the inordinate jump in the sequence ladder, indicating that the adjacent phosphodiester bond is not cleaved during formamide hydrolysis. Our feeling now is that this position is occupied by a modified G because of its slight susceptibility to




Fig.1. Enzymatic degradation of the 5'-end labelled tRNA^{Gly}. L is formamide ladder. T₁, U₂, A and PhyM represent the various enzymatic digestion tracks.

T₁ RNase and is perhaps the same modification which is present in some other archaebacterial tRNAs [9].

Fig.3,4 show the primary structure of tRNA^{Gly} and tRNA^{Ala} as deduced from the above results.




Fig.2. Enzymatic degradation of the 5'-end labelled fragment of tRNA^{Gly} covering the sequence A-9-G-19.




Fig.3. Cloverleaf folding of tRNA^{Gly}. Numbering is according to the Cold Spring Harbor convention.




Fig.4. Cloverleaf folding of tRNA^{Asn}. Numbering is according to the Cold Spring Harbor convention.




Fig. 5. Two-dimensional thin-layer chromatography of position N₃₄ of tRNA^{Asn}. First dimension is isobutyric acid: 0.5 M NH₄OH, 5:3 (v/v) and second dimension is isopropanol: conc. HCl: H₂O, 20:15:15 (v/v).

3.2. Modified nucleosides

Separate chromatographic analyses of the nucleoside diphosphates in 0.55 M ammonium sulfate and 1.75 M ammonium formate as well as the two-dimensional chromatography of the 5'-nucleoside monophosphates [10] indicated the presence of: ψ_{54} , ψ_{55} , Cm₅₆ and m¹I₅₇ [2] in these tRNAs and the presence of t⁶A₃₇ in tRNA^{Asn}. The dinucleotide triphosphate pCmpm¹Ip has the following R_f values: the R_{pAp} in 0.55 M ammonium sulfate is 1.95 and the R_{pUp} in 1.75 M ammonium formate (pH 3.5) is 1.56. N-34 in tRNA^{Asn} is an unknown modification of G that could not be identified with the available standards: its position in a two-dimensional system on cellulose thin-layer plates is seen in fig. 5.

4. DISCUSSION

Methanogens along with halobacteria, thermoacidophiles and some thermophiles are classified together in the kingdom of archaeobacteria. We report here the first tRNA sequences from the methanogens, although a tRNA^{Ala} gene sequence has been reported [5].

Among present data, only in the case of alanine, are representatives of both halobacteria and methanogens available for comparative analyses. The number of structural differences between these tRNAs (78% identical) could be a result of the phylogenetic divergence, or the fact that the two tRNAs have different anticodons and therefore may not have a recent ancestor [4].

The sequences reported here have many features found in other archaeobacterial tRNAs. The *M. thermoautotrophicum* sequences however, exhibit rather long (10–11 bases) D-loops. The consensus sequence N ψ C_mm¹IAAU is present in the T-loop, but rather than m¹ ψ , as in the halobacterial sequences, N is ψ in the methanogen sequences. The high G-C content (Gly and Asn 65%) is characteristically archaeobacterial [1], and moreover inherent to thermophiles like *M. thermoautotrophicum* [11]. Finally, these tRNAs, as other archaeobacterial tRNAs, have no dihydrouridine which may be because of the instability of this nucleotide under the growth conditions of these organisms. The D-loop is complicated however by an unknown nucleotide which defies hydrolysis in formamide or chemical degradation.

Comparison of the methanogen tRNAs with either eubacterial or eukaryotic tRNAs of the same family shows both methanogen sequences to be too distant to enable any conclusion as to their phylogeny (average number of differences was 31 after conversion of modified nucleotides to parents [4]). In the case of tRNA^{Asn} and particularly tRNA^{Gly}, the D-loop is longer than either the eubacterial or the eukaryotic sequence. The length of the variable loop is different only in the case of the eukaryotic glycine tRNA. More precise comparisons will be possible as new sequences become available.

ACKNOWLEDGEMENTS

The authors sincerely thank Dr R.S. Wolfe for supplying them with frozen cells and the Natural Science and Engineering Research Council of Canada for financial support.

REFERENCES

- [1] Balch, W.E., Fox, G.E., Magnum, L.J., Woese, C.R. and Wolfe, R.S. (1979) Microbiol. Rev. 43, 260–296.

- [2] Woese, C.R. and Fox, G.E. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5088–5090.
- [3] Gu, X.-R., Nicoghossian, K., Cedergren, R.J. and Wong, J.T.-F. (1983) *Nucleic Acids Res.* 16, 5433–5443.
- [4] Cedergren, R.J., LaRue, B., Sankoff, D. and Grosjean, H. (1981) *CRC Crit. Rev. Biochem.* 11, 35–104.
- [5] Jarsch, M. and Böck, A. (1983) *Nucleic Acids Res.* 11, 7537–7545.
- [6] Peattie, D.E. (1979) *Proc. Natl. Acad. Sci. USA* 76, 1760–1764.
- [7] Donis-Keller, H., Maxam, A.M. and Gilbert, W. (1977) *Nucleic Acids Res.* 4, 2527–2535.
- [8] Gupta, R. and Randerath, K. (1979) *Nucleic Acids Res.* 6, 3443–3458.
- [9] Kilpatrick, M.W. and Walker, R.T. (1981) *Nucleic Acids Res.* 9, 4387–4390.
- [10] *Methods Enzymol.* 54, 20–70, and Sprinzl, M., Gruter, F. and Gauss, D.H. (1979) *Nucleic Acids Res.* 6, r1.
- [11] Kagawa, Y., Nojira, H., Nukiwa, H., Ishizuka, M., Nakajima, T., Yasuhara, T., Tanaka, T. and Oshima, T. (1984) *J. Biol. Chem.* 259, 2956–2960.