

Import inhibition of poly(His) containing chloroplast precursor proteins by Ni²⁺ ions

R. Rothen, M. Thiess, P. Schumann, A. Boschetti*

Institut für Biochemie, Universität Bern, Freiestr. 3, CH-3012 Bern, Switzerland

Received 9 July 1996; revised version received 28 July 1996

Abstract The precursor of the small subunit of ribulose-1,5-bisphosphate carboxylase (pSS) and a modified pSS containing a C-terminal hexahistidyl tail (pSS(His)₆) were imported into isolated *Chlamydomonas* chloroplasts with comparable efficiency. In the presence of Ni²⁺ ions the import of pSS(His)₆ was inhibited and the precursor bound to the envelope remained protease sensitive, while import of pSS was not affected. Addition of an excess of L-histidine suppressed the inhibition demonstrating that the hexahistidyl-Ni²⁺ complex was responsible for import inhibition. Inhibition could be observed between about 0.5 and 10 mM Ni²⁺, depending on the total protein content in the assay. Import incompetent Ni²⁺-precursor complexes can be used to study early events in chloroplast protein import.

© 1997 Federation of European Biochemical Societies.

Key words: Chloroplast protein import; Ni²⁺ inhibition; *Chlamydomonas reinhardtii*

1. Introduction

Most proteins of chloroplasts as well as of mitochondria are imported into the organelle from the cytoplasm where they are synthesized as precursor proteins containing an N-terminal transit peptide [1]. Proteins of the envelope membrane are involved in import as receptors and components of the translocation apparatus [2,3]. Our knowledge on the chloroplast import apparatus is based mainly on the biochemical analysis of so-called translocation complexes. As import is a rather fast process such analysis will be most successful when importation of in vitro synthesized, labeled precursor proteins into isolated intact chloroplasts is arrested at an intermediate stage, thereby accumulating translocation complexes. Such an accumulation might be induced by precursor proteins bearing a C-terminal domain that can be transformed into a rigid or bulky structure which is sterically hindered from passing the membrane pore. Import of precursors into mitochondria is blocked, e.g. by binding of antibodies to the C-terminal domain [4], or by binding of the substrate analogue methotrexate to a DHFR domain attached to the C-terminus of the precursor protein. This inhibition is probably due to hindered unfolding of the import protein, i.e. stabilization of a rigid conformation [5,6].

Similar attempts have been made for the analysis of components of the import apparatus in chloroplast envelopes,

however, with contradictory results. On the one hand, the import of a transit peptide-DHFR fusion protein is not blocked by the presence of methotrexate in higher plants, pointing to a strong unfolding activity associated with chloroplast envelope membranes [7,8]. Also, the protease sensitivity of a chimeric precursor protein containing the ricin A chain, when bound to chloroplasts, is indicative of an unfolding activity of the chloroplast envelope [9]. On the other hand, binding of specific antibodies to the DHFR moiety or to the protein A moiety of a precursor-protein A fusion protein arrested import [10,11]. Also consistent with an unfolding/refolding model is the inhibition by glyphosate of the import of the precursor for 5-enolpyruvylshikimate-3-phosphate synthase in the presence of its substrate [12].

Not much attention has so far been paid to the protein import into chloroplasts of *Chlamydomonas reinhardtii*, although this unicellular green alga is widely used as a model organism to investigate various cell biological problems of green plants [13]. This neglect is probably due to the tedious isolation procedure for chloroplasts and to the fact that precursor proteins from higher plants are not imported or not correctly processed by *Chlamydomonas* chloroplasts, and vice versa [14].

A very common affinity tag introduced into fusion proteins is the hexa-histidyl sequence simplifying the isolation of proteins via its high affinity to nickel ions bound to a chelating resin [15]. This rather small sequence, however, was never used for formation of a rigid structure to arrest translocation. Here, we present evidence that the import into chloroplasts of *Chlamydomonas* of a fusion precursor protein containing a hexahistidyl tail at the C-terminus is hindered by the addition of nickel ions.

2. Materials and methods

2.1. Construction of plasmid

The plasmid pSSpSP64 containing the cDNA of the gene *rbcS2*, which codes for one of the two forms of pSS present in *C. reinhardtii* was a generous gift of Dr. M.L. Mishkind. The two pSS in *Chlamydomonas* differ only in a few amino acids in the mature protein but not in the transit sequence [16].

The 0.8 kb *Pst*I fragment of plasmid pSSpSP64 was cloned into the *Pst*I site of the commercial expression vector pQE-3 (Qiagen GmbH, Hilden, Germany). The recombinant plasmid was cut with *Eco*47III and *Eco*RI and the 0.8 kb fragment was isolated from an agarose gel. The plasmid pQE-12 (Qiagen GmbH, Hilden, Germany) which contained the hexahistidyl-coding sequence was cut with *Bgl*II, blunted with Klenow enzyme and cut with *Eco*RI. The 3.4 kb fragment was ligated with the 0.8 kb fragment of the *Eco*47III-*Eco*RI restriction. The 0.8 kb *Pvu*II-*Pst*I fragment of the resulting plasmid was ligated with the 2.9 kb *Pvu*II-*Pst*I fragment of plasmid pSSpSP64, finally leading to plasmid pSSHspSP64. The in vitro transcript obtained from this plasmid served to synthesize in vitro precursor proteins of which the last amino acid (val) was replaced by the hexahistidyl tail

*Corresponding author: Fax: (41) (31) 631 48 87.

Abbreviations: DHFR, dihydrofolate reductase; (His)₆, hexahistidyl peptide; hsp70, 70 kDa heat shock protein; pSS, precursor of the small subunit of ribulose-1,5-bisphosphate carboxylase; pSS(His)₆, pSS with C-terminal hexahistidyl tail

[17]. All operations were essentially performed according to Maniatis et al. [18].

2.2. Preparation of precursor proteins

The in vitro transcription of the genes was performed in 50 µl assays as described [19]. For the in vitro translation a wheat germ extract was prepared according to Mishkind et al. [20]. To synthesize radiolabeled precursor proteins each 150 µl of translation mixture contained 30 µl wheat germ extract, 2 mM ATP, 0.4 mM GTP, 25 µM non-radioactive amino acids (all but methionine), 60 µCi [³⁵S]methionine, 2 mM dithiothreitol, 40 µM spermine, 24 mM HEPES buffer, pH 7.5, 1 mM Mg acetate, 60 mM K acetate and 5 µl mRNA (about 2 µg). The translation was carried out over 40 min at 30°C.

2.3. Preparation of chloroplasts

Chloroplasts were isolated from synchronized cultures of *C. reinhardtii* cw-15 (stock CC-277 from the Chlamydomonas Genetics Center, Duke University, Durham, NC, USA) as described previously [21].

2.4. Import assay

20 µl of chloroplasts ($2\text{--}4 \times 10^8/\text{ml}$) and 10 µl translation mixture containing the radiolabeled precursor proteins were incubated in a final volume of 100 µl import buffer (250 mM sorbitol, 35 mM HEPES buffer, pH 7.8) with an excess of 4 mM ATP for 30 min at 25°C under white light. The chloroplasts were reisolated by centrifugation through 20% Percoll in import buffer and carefully washed with import buffer. Blockage of import was achieved by addition of 1 mM NiSO₄ to the import assay. The reisolated chloroplasts were solubilized and the proteins analyzed by SDS-polyacrylamide gel electrophoresis [24]. For quantitative determination of the radioactivity in the protein bands, the dried gel was scanned by use of a Phosphor-Imager (Molecular Dynamics).

3. Results and discussion

Starting with the cloned gene of the precursor of the small subunit of ribulose-1,5-bisphosphate carboxylase (pSS), an expression vector was constructed coding for a fusion protein composed of pSS with the last amino acid replaced by a C-terminal hexahistidyl tail (pSS(His)₆). The genes for the native protein pSS and for the fusion protein pSS(His)₆ were transcribed in vitro and the precursor proteins synthesized and labeled with [³⁵S]methionine in a wheat germ translation system. Driven by light, the labeled precursors were imported into isolated chloroplasts of *C. reinhardtii* and the proteins electrophoretically analyzed. Fig. 1 demonstrates that the fu-




Fig. 1. Import of in vitro synthesized precursor protein into isolated chloroplasts. Fluorogram of a 10–20% SDS-polyacrylamide gel with electrophoretically separated incubation mixtures. Each lane was loaded with an equal amount of chlorophyll *a+b*. Isolated chloroplasts were incubated with in vitro synthesized, radiolabeled pSS (lanes 1–3) or pSS(His)₆ (lanes 4–6). After incubation the chloroplasts were centrifuged through Percoll and washed. Lanes: 1,4, incubation in the dark without addition of ATP; 2,5, incubation in the light with externally added ATP; 3,6, chloroplasts were treated with thermolysin after incubation. p, precursor protein; m, mature protein.




Fig. 2. Inhibition of pSS(His)₆ import by Ni²⁺ ions. Fluorogram of a 10–20% polyacrylamide gel with electrophoretically separated incubation mixtures. Isolated chloroplasts were incubated in the light and in the presence of ATP with in vitro synthesized, radiolabeled pSS(His)₆ (lanes 1–3) or pSS (lane 4). Additions: 1 mM Ni²⁺ to the samples of lanes 2–4 prior to incubation. Lane 3: after half-time of incubation, i.e. 15 min, 5 mM L-histidine was added. Chloroplasts were reisolated by centrifugation through Percoll and washed. Each lane was loaded with an equal amount of chlorophyll *a+b*. p, precursor protein; m, mature protein.

sion protein pSS(His)₆ was processed in isolated chloroplasts to about the same degree as the non-modified pSS. With the very active chloroplast preparation of Fig. 1, the radioactivity of the mature proteins SS(His)₆ and SS was 83% (lane 2) and 92% (lane 5), respectively (100% = mature and precursor protein). With chloroplasts of average activity 55 and 57% (Fig. 2) or 72 and 63% (Fig. 3) of the precursors were processed. To prove that the processed proteins were internalized, the chloroplasts were treated with thermolysin after the import reaction (Fig. 1). This protease is known to degrade and remove the external precursor proteins, but not the mature processed proteins protected within the chloroplast. In spite of the poly(His)₆ tail the in vitro synthesized chimeric protein was imported and processed with high efficiency by isolated chloroplasts.

However, an important difference between the two precursor proteins with and without poly(His)₆ tail became obvious when the import reaction was performed in the presence of nickel ions. At a concentration of 1 mM Ni²⁺ the import of pSS(His)₆ was inhibited while the import of pSS was not affected (Fig. 2, lanes 2,4). The precursor protein pSS(His)₆ blocked by Ni²⁺ and adsorbed at the envelope surface was completely thermolysin sensitive (data not shown). If indeed metal ions are responsible for the import inhibition, addition of Ni²⁺-binding agents, such as EDTA or histidine, to the reaction mixture should have an antagonistic effect. While

Table 1

Binding constants related to complexes of the binding domain in chimeric import proteins and ligands

Ligand	Binding domain	Binding constant	Reference
Ni ²⁺	L-His	$4.6 \times 10^8 \text{ M}^{-1} (K_1)$	[25]
		$3.3 \times 10^{15} \text{ M}^{-1} (\beta_2)$	[25]
Ni ²⁺	-(His) ₆	$1 \times 10^{13} \text{ M}^{-1}$	[26]
Ni ²⁺	Histidylhistidin	$3.1 \times 10^{10} \text{ M}^{-1} (\beta_2)$	[27]
Ni ²⁺	EDTA ³⁻	$3.6 \times 10^{11} \text{ M}^{-1}$	[27]
Ni ²⁺	Imidazole	$1 \times 10^3 \text{ M}^{-1} (K_1)$	[25]
		$5 \times 10^{10} \text{ M}^{-1} (\beta_5)$	[25]
Ni ²⁺	ATP	$1 \times 10^5 \text{ M}^{-1}$	[27]
Methotrexate	DHFR	$3.2 \times 10^{10} \text{ M}^{-1}$	[27]

$K_1 = (Me \cdot L)/(Me)(L)$; $\beta_n = (Me \cdot L_n)/(Me)(L)^n$ and applies to the equation $Me + nL = Me \cdot L_n$, where *Me* = metal ion, *L* = ligand.




Fig. 3. Effect of increasing Ni^{2+} concentrations on the chloroplast import of pSS and pSS(His)₆. Composite picture of three fluorograms of a 10–20% polyacrylamide gel with electrophoretically separated incubation mixtures. Each lane was loaded with an equal amount of chlorophyll *a+b*. Isolated chloroplasts were incubated in the presence of the indicated Ni^{2+} concentrations with in vitro synthesized, radiolabeled pSS (lanes a) or pSS(His)₆ (lanes b). p, precursor protein; m, mature protein.

addition of 5 mM L-histidine could indeed revert the import inhibition (Fig. 2, lane 3) the addition of 5 mM EDTA had only a slight effect (not shown). The import reactions per se of pSS and of pSS(His)₆ were not impaired, either by EDTA or by L-histidine. Therefore, this inhibition by Ni^{2+} is clearly different from the previously described effect of the oxidizing CuCl_2 on protein import [22].

The import inhibition of pSS(His)₆ by nickel ions could only be observed over a narrow concentration range between about 0.5 and 10 mM Ni^{2+} . The limits could not be determined precisely. As proteins generally have an inherent binding capacity for heavy metals, and as in different experiments the amount of protein in the assays varied depending on the quality of the chloroplast preparation and of the in vitro translation system, the Ni^{2+} concentration available for complex formation with pSS(His)₆ probably varied to a considerable degree from experiment to experiment.

Nevertheless, Fig. 3 shows the processing and import activity of intact chloroplasts for pSS and pSS(His)₆ as a function of low Ni^{2+} concentrations in the assay. Between 0.5 and 1 mM Ni^{2+} the import of pSS(His)₆ became increasingly inhibited as shown by the disappearance of the distinct band of mature SS(His)₆. In contrast, the import of pSS was much less affected by Ni^{2+} , demonstrating that the chloroplasts were still active in the presence of even 3 mM Ni^{2+} . The quantitative evaluation of this experiment by PhosphorImager is shown in Fig. 4. Even under Ni^{2+} inhibition a background level of mature SS(His)₆ was formed. It is not yet clear whether this reflects incomplete import inhibition or more likely the processing of some precursor by a stromal peptidase released from broken chloroplasts during incubation. In experiments with additional thermolysin treatment of the assays this background level was less pronounced (data not shown).

To explain the inhibition of protein import by formation of complexes between the import protein and a ligand, the idea was proposed – as a working hypothesis – that an import protein with a rigid conformation will mechanically not be able to pass through the import pore [5]. The envelope, however, equipped with a membrane-bound chaperonin of the *hsp70* class [23,24] contains an unfolding capacity. Weak conformational forces in an import protein will be compensated when the precursor protein binds to the receptor, hereby facilitating the translocation of the protein [7]. Yet, a rigid conformation stabilized by strong forces cannot be compensated. Such a protein remains adsorbed or sticks firmly as an intermediate import complex in the envelope. Table 1 shows that the affinity of Ni^{2+} is highest to an excess of monomeric L-histidine and decreases from poly(His)₆ to EDTA. This could

explain why, in contrast to L-histidine, EDTA was not able to suppress import inhibition. Furthermore, the binding constant of methotrexate to DHFR is lower than that of Ni^{2+} to (His)₆. Eventually the unfolding capacity of the chloroplast envelope membrane [8,9] is able to discriminate between the two binding constants, thereby explaining why the import of (His)₆-containing precursor protein is inhibited by Ni^{2+} , while DHFR-fusion proteins are still imported in the presence of methotrexate. To investigate this point further, we are currently producing a precursor-fusion protein with two C-terminal additions: a DHFR sequence followed by a hexahistidyl peptide.

This construct will allow us to test concomitantly the blocking capacities of methotrexate and Ni^{2+} ions in *C. reinhardtii*.

Acknowledgements: We thank Dr. M. Mishkind for the kind gift of the cloned pSS. This work was supported partly by the Swiss National Foundation of Scientific Research (grant No. 3100-36 459.92).




Fig. 4. Quantitative estimation of the inhibition of pSS(His)₆ import into isolated chloroplasts at increasing Ni^{2+} concentration. The amount of radioactivity in the mature and precursor protein band of Fig. 3 was measured using a PhosphorImager. The amount of mature protein is represented in percent of the sum of mature and precursor protein.

References

- [1] De Boer, A.D. and Weisbeek, P.J. (1991) *Biochim. Biophys. Acta* 1071, 221–253.
- [2] Pfanner, N. (1992) in: *Plant Gene Research – Cell Organelles* (Herrmann, R.G., ed.) pp. 371–400, Springer Verlag, Berlin.
- [3] Perry, S.E. and Keegstra, K. (1994) *Plant Cell* 6, 93–105.
- [4] Schwaiger, M., Herzog, V. and Neupert, W. (1987) *J. Cell Biol.* 105, 235–246.
- [5] Eilers, M. and Schatz, G. (1986) *Nature* 322, 228–232.
- [6] Rassow, J., Guiard, B., Wienhues, U., Herzog, V., Hartl, F.U. and Neupert, W. (1989) *J. Cell Biol.* 109, 1421–1428.
- [7] America, T., Hageman, J., Guéra, A., Rook, F., Archer, K., Keegstra, K. and Weisbeek, P. (1994) *Plant Mol. Biol.* 24, 283–294.
- [8] Endo, T., Kawakami, M., Goto, A., America, T., Weisbeek, P. and Nakai, M. (1994) *Eur. J. Biochem.* 225, 403–409.
- [9] Walker, D., Chaddock, A.M., Chaddock, J.A., Roberts, L.M., Lord, J.M. and Robinson, C. (1996) *J. Biol. Chem.* 271, 4082–4085.
- [10] Schnell, D.J. and Blobel, G. (1993) *J. Cell Biol.* 120, 103–115.
- [11] Wu, C., Seibert, F.S. and Ko, K. (1994) *J. Biol. Chem.* 269, 32264–32271.
- [12] Della Cioppa, G. and Kishore, G.M. (1988) *EMBO J.* 7, 1299–1305.
- [13] Rochaix, J.-D. (1995) *Annu. Rev. Genet.* 29, 209–230.
- [14] Su, Q.X. and Boschetti, A. (1994) *Eur. J. Biochem.* 217, 1039–1047.
- [15] Stüber, D., Gentz, R., Döbeli, H., Bannwart, W. and Hochuli, E. (1988) *Biotechnology* 1988, 1321–1325.
- [16] Goldschmidt-Clermont, M. and Rahire, M. (1986) *J. Mol. Biol.* 191, 421–432.
- [17] Su, Q.X. and Boschetti, A. (1993) *Eur. J. Biochem.* 217, 1039–1047.
- [18] Maniatis, T., Fritsch, E.F. and Sambrook J. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- [19] Su, Q.X., Niklaus, A., Rothen, R. and Boschetti, A. (1992) *FEBS Lett.* 300, 157–161.
- [20] Mishkind, M.L., Greer, K.L. and Schmidt, G.W. (1987) *Methods Enzymol.* 148, 274–294.
- [21] Mendiola-Morgenthaler, L., Leu, S. and Boschetti, A. (1985) *Plant Sci.* 38, 33–39.
- [22] Seedorf, M. and Soll, J. (1995) *FEBS Lett.* 367, 19–22.
- [23] Waegemann, K. and Soll, J. (1991) *Plant J.* 1, 149–158.
- [24] Ko, K., Bornemisza, O., Kourtz, L., Ko, Z.W., Plaxton, W.C. and Cashmore, A.R. (1992) *J. Biol. Chem.* 267, 2986–2993.
- [25] Martel, A.E. and Smith, R.M. (1982) *Critical Stability Constants*, vol. 5, 1st Suppl., Plenum Press, New York.
- [26] Crowe, J. and Henco, K. (1992) *The QIAexpressionist*, DIAGEN GmbH, Hilden, Germany.
- [27] Martel, A.E. (1971) *Stability Constants of Metal-ion Complexes*, Suppl. 1, part II, The Chemical Society, London.