

# Sequence of subunit *c* of the Na<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub> ATPase of *Acetobacterium woodii*: proposal for determinants of Na<sup>+</sup> specificity as revealed by sequence comparisons

Stefan Rahlfs, Volker Müller\*

Institut für Mikrobiologie der Universität, Grisebachstr. 8, D-37077 Göttingen, Germany

Received 23 December 1996

**Abstract** A 3.2 kb *EcoRI* fragment carrying genes for Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase was cloned from chromosomal DNA of *Acetobacterium woodii*. DNA sequence analysis revealed the presence of an open reading frame which was identified by data base searches and comparison with the experimentally derived N-terminal amino acid sequence to code for subunit *c* of Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase. A comparison of the primary sequences of the two well established Na<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub> ATPases from *Acetobacterium woodii* and *Propionigenium modestum* with H<sup>+</sup>-translocating enzymes indicates the length of the C-terminus as well as specific residues located in the cytoplasmic membrane to be important for Na<sup>+</sup> transport.

© 1997 Federation of European Biochemical Societies.

**Key words:** ATPase; Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub>; Subunit *c*; DNA sequence; *Acetobacterium woodii*

## 1. Introduction

F<sub>1</sub>F<sub>0</sub> ATPases are reversible enzymes which couple ATP synthesis in the F<sub>1</sub> domain with the flow of ions through the F<sub>0</sub> domain [1,2]. Although the enzymes have been known for decades, the mechanism of ion transport as well as the coupling of ATP synthesis with ion flow is far from being settled. An alternative approach to elucidate their function is to perform a comparative study with enzymes exhibiting an altered ion specificity. The anaerobic bacteria *Propionigenium modestum* and *Acetobacterium woodii* are the only well established examples of organisms having ATPases which, as revealed by inhibitor studies, subunit composition, primary sequences and electron microscopy, are typical F<sub>1</sub>F<sub>0</sub> enzymes but use Na<sup>+</sup> instead of H<sup>+</sup> as coupling ion [3–5]. The F<sub>0</sub> complex was shown to determine the ion specificity of the enzyme [6] and the ‘active carboxylate’ of subunit *c* was shown biochemically to contribute to Na<sup>+</sup> liganding [7,8]. As a first step towards a better understanding of Na<sup>+</sup> transport through the F<sub>1</sub>F<sub>0</sub> ATPase from *A. woodii* we cloned and sequenced the gene encoding subunit *c*. This sequence is now the second sequence available for subunit *c* from Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPases. A comparison of these two sequences to those from H<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPases revealed determinants probably responsible for Na<sup>+</sup> specificity.

## 2. Materials and methods

### 2.1. Organism and plasmids

*A. woodii* (DSM 1030) was obtained from the Deutsche Sammlung für Mikro-organismen und Zellkulturen, Braunschweig, Germany, and grown under strictly anaerobic conditions on carbonate-buffered medium supplied with 0.4% glycine [9]. *E. coli* DH5α (*supE44 ΔlacU169(Φ80lacZΔM15) hsdR17 recA1 endA1 gyrA96 thi1 relA1* [10]) was grown on LB at 37°C. Plasmids used were pSE420 [11], pHSG 398 and 399 [12], and pBluescript II SK and KS (Stratagene, San Diego, CA, USA).

### 2.2. Molecular procedures

Chromosomal DNA of *A. woodii* was isolated by a modified Mar-mur preparation [13]. Cells were grown to late logarithmic growth phase, pelleted by centrifugation and lysed by alkaline SDS solution. The protein was precipitated with 6 M guanidinium thiocyanate and isobutanol, and the DNA was bound to glassmilk in the presence of NaJ (90.8 g NaJ, 1.5 g Na<sub>2</sub>SO<sub>3</sub>, H<sub>2</sub>O ad 100 ml). After washing with ethanol (50%), NaCl (0.1 M), Tris-HCl (0.02 M, pH 7.5) and EDTA (1 mM), the DNA was eluted with TE buffer. The DNA was then restricted, size fractionated by gradient centrifugation and cloned into pSE420. All procedures used were standard techniques [10]. Oligonucleotides were synthesized on a Gene Assembler Plus apparatus as recommended by the manufacturer (Pharmacia LKB). DNA sequence was determined by the chain termination method of Sanger and analyzed on a VAX computer using the Wisconsin genetics computer group sequence analysis software package, version 8.1 (University of Wisconsin Biotechnology Center, Madison, WI, USA).

## 3. Results and discussion

A homologous probe covering the DNA region upstream of the *EcoRI* site in *atpA* [5] was amplified by PCR with oligonucleotides derived from the known DNA sequence in *atpA* and from the N-terminal amino acid sequence of the 19 kDa subunit. Southern blot analysis of genomic DNA restricted with *EcoRI* revealed a 3.2 kb fragment that hybridized with the homologous probe. This fragment was cloned in pSK<sup>+</sup>, designated pSR4 and partially sequenced.

At the 5′ end of pSR4 an open reading frame of 270 bp was found. This ORF was identified by data base searches as *atpE* which encodes subunit *c* of F<sub>1</sub>F<sub>0</sub> ATPases. The identity of the gene was verified by comparison of the deduced amino acid sequence which matched the experimentally derived N-terminal sequence of subunit *c* of the Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase from *A. woodii* [4]. *atpE* starts with an ATG codon and encodes a strongly hydrophobic protein with a molecular mass of 8.96 kDa (Fig. 1). 60, 47, and 23% of its residues are identical with subunit *c* from the Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase from *P. modestum*, the H<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase from *Synechocystis* 6803 and the H<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPase from *Escherichia coli*, respectively.

A comparison of the deduced amino acid sequences with H<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub> ATPases revealed the following un-

\*Corresponding author. Fax: (49) (551) 393793.  
E-mail: wmueller2@gwdg.de

```

1
GGAGGAAACATAATTTATGGAGGTTTATGATTTTATTAAGCATGTTGAGCCATTGGAGC
  M E G L D F I K A C S A I G A
  ? E (I) L D F (I) K

61
TGGGATTGCGATGATCGCTGGGCTGGGACCTGGGATTTGGTCAGGGTTTTCGCCCGCTAA
  G I A M I A G V G P G I G Q G F A A G K

121
AGGCCCGAAGCCGTTGGCCGCCAGCCGGAAGCGCAAGTGATATTATCAGAACCATGCT
  G A E A V G R Q P E A Q S D I I R T M L

181
TTTGGGTGCAGCAGTTGCTGAAACCACTGGTATTTATGGTTTGATTGTCGCAITGATTTT
  L G A A V A E T T G I Y G L I V A L I L

241
ATTATTTGGCAACCCATTTTAAACAGTAACGATAGATAA
  L F G K P I F L K T V T D R Z

```

Fig. 1. Nucleotide sequence and deduced amino acid sequence of *atpE* of *A. woodii*. The sequence was obtained by primer walking. Shine-Dalgarno sequences are underlined, and the experimentally derived N-terminal amino acid sequence is given below the translated sequence. The sequence has been submitted to GenBank and is available under accession number U63995.

usual features in subunit *c* from *A. woodii* and *P. modestum* which we propose are relevant for Na<sup>+</sup> transport:

(1) Subunit *c* folds in the membrane like a hairpin with its N- and C-termini being located in the periplasm [14]. Interestingly, the size of subunit *c* in *A. woodii* is increased by two residues at the N-terminus and, more important, nine residues at the C-terminus; this enlarges the hydrophilic, periplasmic portion of the polypeptide (Fig. 2). As expected, such an increase is also observed with the Na<sup>+</sup> ATPase from *P. modestum*. Therefore, this feature can be considered significant in determining the ion specificity. As discussed before for *P. modestum* [15], the enlargement of subunit *c* might alter the geometry of the F<sub>0</sub> complex facing the periplasm, by altering the interaction either of multiple copies of subunit *c* or of subunit *c* with other membrane-integral subunits. Taking into consideration the presence of a selectivity filter at the mouth of the F<sub>0</sub> complex, as envisaged by Fillingame [16], the additional residues in subunit *c* of *P. modestum* and *A. woodii* might constitute (or contribute to) such a filter which

widens the mouth and allows the passage of the large sodium ion into the F<sub>0</sub> complex. This assumption is corroborated by the finding that random mutagenesis of subunit *c* of *P. modestum* led to mutants which lost Na<sup>+</sup> but retained H<sup>+</sup> transport by exchange of the bulky residues F84 and L87 against smaller residues [15]. Based on these findings we predict that deletion of the periplasmic C-terminus of *A. woodii* and *P. modestum* would result in a loss of Na<sup>+</sup> with a retention of H<sup>+</sup> pumping.

(2) Residues able to ligand the Na<sup>+</sup> ion should be present in Na<sup>+</sup> but not H<sup>+</sup> ATPases. In the membrane spanning regions of subunit *c* of *A. woodii* six residues are present which have the potential ability to ligand ions by use of their free electron pair: M19, P25, Q29, E62, T63 and T64. A multiple alignment revealed that only P25, Q29, E62, T63 and T64 are conserved or functionally substituted in Na<sup>+</sup>-F<sub>1</sub>F<sub>0</sub> ATPases from *A. woodii* and *P. modestum* and in the second half of subunit *c* of the Na<sup>+</sup>-V<sub>1</sub>V<sub>0</sub> ATPase from *Enterococcus hirae* but not in H<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub> ATPases (Fig. 2). E62 and T63 but not T64 were shown by site directed mutagenesis to be essential for Li<sup>+</sup> binding in the F<sub>1</sub>F<sub>0</sub> ATPase from *E. coli* [17]. Therefore, we predict that P25, Q29, E62 and T63 constitute (or contribute to) the Na<sup>+</sup> binding site of the Na<sup>+</sup> ATPase of *A. woodii*. These four residues are located at approximately the same distance from the surface of the cytoplasmic membrane, and therefore would also physically be able to build at least part of the cavity in which the Na<sup>+</sup> ion is coordinated (Fig. 3). Experimental evidence for a close proximity of A24 and I28 of helix one (corresponding to P25 and Q29 in *A. woodii*) to D61 of helix two (corresponding to E62 in *A. woodii*) in *E. coli* is available [18,19].

In conclusion, the data available now revealed two determinants for Na<sup>+</sup> specificity of the F<sub>1</sub>F<sub>0</sub> ATPase of *A. woodii*: an enlargement of the C-terminus of subunit *c* accompanied by a change in the geometry of the complex and the presence of the sodium ion binding motif P25, Q29, E62 and T63; residues in this motif can be substituted by functionally identical residues, i.e. those of appropriate size and bearing a free electron pair. Although one or the other determinant is

periplasm			membrane					
Awo	1	.....	..MEGLDFIK	ACSAIGAGIA	M.IAGVGPGI	GQGFAAGKGA		37
Pmo	1	.....	M DMVLAKTVVL	AASAVGAGAA	M.IAGIGPGV	GQGYAAGKAV		40
Ehi	69	GFVIAF	LIFINLGS DM	SVVQGLNFLG	ASLPFAFTGL	FSGIAQKGA		114
Eco	1	.....	...MENLNM	DLLYMAAAVM	MGLAATGA AI	GIGILGGKFL		36
Val	1	.....	...METL.L	SFSAIAVGII	VGLASLGT AI	GFALLGGKFL		35
Ps3	1	.....	.....M	SLGVLAAAIA	VGLGALGAGI	GNGLIVSRTI		31
Bme	1	.....	.....	.MGLIASAIA	IGLAALGAGI	GNGLIVSKTI		29
Sec	1	.....	....MDSTVA	AASVIAAALA	VGLGATGPGI	GQGNASGQAV		36
			cytoplasm			membrane		
Awo	38	EAVGRQPEAQ	SDIIRTMLLG	AAVAETGTIY	GLIVALILLF	GKPIFLKTVT	DR	89
Pmo	41	ESVARQPEAK	GDIISTMVLG	QAIAESTGIY	SLVIALILLY	ANP.FVGLLG		89
Ehi	115	AAGIQILAKK	PEHATKGIIF	AAMVETYAIL	GFVISFLLVL	NA.....		156
Eco	37	EGAARQPDLI	PLLRTQFFIV	MGLVDAIPMI	AVGLGLYVMF	AVA.....		79
Val	36	EGAARQPEMA	PMLQVKMFII	AGLLDAVPMI	GIVIALLFTF	ANPFVQQLG.		84
Ps3	32	EGIARQPELR	PVLQTTMFIG	VALVEALPII	GVVFSFIYLG	R.....		72
Bme	30	EGTARQPEAR	GTLTSMMFVG	VALVEALPII	AVVIAFMVQG	K.....		70
Sec	37	SGIARQPEAE	GKIRGTLLLT	LAFMESLTIY	GLVIALVLLF	ANPFA.....		81

Fig. 2. Alignment of subunit *c* from *A. woodii* with subunit *c* from *P. modestum* [20], *E. hirae* [21], *E. coli* [22], *V. alginolyticus* [23], thermophilic bacterium PS3 [24], *Bacillus megaterium* [25], and *Synechocystis* 6803 [26]. The topology of subunit *c* is indicated.




Fig. 3. Hairpin folding model of subunit *c* of the  $\text{Na}^+$ - $\text{F}_1\text{F}_0$  ATPase of *A. woodii*. A potential sodium ion binding motif is indicated.

present in certain other enzymes (i.e. an enlarged C-terminus in *V. alginolyticus* and a sodium ion binding motif in *Synechocystis* 6803), both are present only in  $\text{Na}^+$ - $\text{F}_1\text{F}_0$  ATPases from *A. woodii* and *P. modestum*.

**Acknowledgements:** This work was supported by a grant from the Deutsche Forschungsgemeinschaft. We are indebted to Prof. G. Gottschalk, Göttingen, for support and stimulating discussions, and to H. Ebner for skilful technical assistance.

## References

- [1] Schneider, E. and Altendorf, K. (1987) *Microbiol. Rev.* 51, 477–497.
- [2] Senior, A.E. (1990) *Annu. Rev. Biophys. Biophys. Chem.* 19, 7–41.
- [3] Dimroth, P., Laubinger, W., Kluge, C., Kaim, G., Ludwig, W. and Schleifer, K.H. (1992) *Ann. NY Acad. Sci.* 671, 310–322.
- [4] Reidlinger, J. and Müller, V. (1994) *Eur. J. Biochem.* 223, 275–283.
- [5] Forster, A., Daniel, R. and Müller, V. (1995) *Biochim. Biophys. Acta* 1229, 393–397.
- [6] Laubinger, W., Deckers-Hebestreit, G., Altendorf, K. and Dimroth, P. (1990) *Biochemistry* 29, 5458–5463.
- [7] Kluge, C. and Dimroth, P. (1993) *J. Biol. Chem.* 268, 14557–14560.
- [8] Spruth, M., Reidlinger, J. and Müller, V. (1995) *Biochim. Biophys. Acta* 1229, 96–102.
- [9] Tschuch, A. and Pfennig, N. (1984) *Arch. Microbiol.* 137, 163–167.
- [10] Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning, A Laboratory Manual*, 2nd edn., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- [11] Brosius, J. (1992) *Methods Enzymol.* 216, 469–483.
- [12] Takeshita, S., Sato, M., Toba, M., Masahashi, W. and Hashimoto-Gotoh, T. (1987) *Gene* 61, 63–74.
- [13] Marmur, J. (1961) *J. Mol. Biol.* 3, 208–218.
- [14] Girvin, M.E. and Fillingame, R.H. (1993) *Biochemistry* 32, 12167–12177.
- [15] Kaim, G. and Dimroth, P. (1995) *J. Mol. Biol.* 253, 726–738.
- [16] Fillingame, R.H. (1990) in *The Bacteria* (Kruschwitz, T.A., Ed.), pp. 345–391, Academic Press, San Diego, CA.
- [17] Zhang, Y. and Fillingame, R.H. (1995) *J. Biol. Chem.* 270, 87–93.
- [18] Miller, M.J., Oldenburg, M. and Fillingame, R.H. (1990) *Proc. Natl. Acad. Sci. USA* 87, 4900–4904.
- [19] Girvin, M.E. and Fillingame, R.H. (1993) *Biochemistry* 33, 665–674.
- [20] Ludwig, W., Kaim, G., Laubinger, W., Dimroth, P., Hoppe, J. and Schleifer, K.H. (1990) *Eur. J. Biochem.* 193, 395–399.
- [21] Kakinuma, Y., Kakinuma, S., Takase, K., Konishi, K., Igerashi, K. and Yamoto, I. (1993) *Biochem. Biophys. Res. Commun.* 195, 1063–1069.
- [22] Kanazawa, H., Mabuchi, K., Kayano, T., Tamura, F. and Futai, M. (1981) *Biochem. Biophys. Res. Commun.* 100, 219–225.
- [23] Krumholz, L.R., Esser, U. and Simoni, R.D. (1989) *Nucleic Acids Res.* 17, 7993–7994.
- [24] Hoppe, J. and Sebald, W. (1980) *Eur. J. Biochem.* 107, 57–65.
- [25] Brusilow, W.S.A., Scarpetta, M.A., Hawthorne, C.A. and Clark, W.P. (1989) *J. Biol. Chem.* 264, 1528–1533.
- [26] Lill, H. and Nelson, N. (1991) *Plant Mol. Biol.* 17, 641–652.