

# In vitro characterization of rice importin $\beta$ 1: molecular interaction with nuclear transport factors and mediation of nuclear protein import

Chang-Jie Jiang<sup>a</sup>, Naoko Imamoto<sup>b</sup>, Rikyu Matsuki<sup>c</sup>, Yoshihiro Yoneda<sup>b</sup>,  
Naoki Yamamoto<sup>a,\*</sup>

<sup>a</sup>Photosynthesis Laboratory, Plant Physiology Department, National Institute of Agrobiological Resources, 2-1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan

<sup>b</sup>Department of Anatomy and Cell Biology, Osaka University Medical School, Suita, Osaka 565-0871, Japan

<sup>c</sup>Central Research Institute of Electric Power Industry, Abiko, Chiba 270-1194, Japan

Received 14 August 1998; received in revised form 7 September 1998

**Abstract** We recently isolated two cDNAs encoding importin  $\beta$  homologues (rice importin  $\beta$ 1 and  $\beta$ 2), the first such homologues identified in plants. To address the function of rice importin  $\beta$ 1 in the process of nuclear import of proteins, we carried out in vitro binding and nuclear import assays. Recombinant protein of rice importin  $\beta$ 1 assembled a complex (PTAC) with rice importin  $\alpha$ 1 and NLS protein, and also bound to the nuclear envelope of tobacco BY-2 cells. Ran-GTP, but not Ran-GDP, interacted with rice importin  $\beta$ 1 and dissociated the heterodimer formed between rice importin  $\alpha$ 1 and rice importin  $\beta$ 1. An in vitro nuclear import assay using digitonin-permeabilized HeLa cells revealed that rice importin  $\beta$ 1 can mediate nuclear envelope docking of NLS proteins and their subsequent translocation into the nucleus. These data strongly suggest that rice importin  $\beta$ 1 functions as a component of the NLS receptor in plant cells.

© 1998 Federation of European Biochemical Societies.

**Key words:** Importin  $\beta$ ; Nuclear localization signal; Nuclear protein import; Rice (*Oryza sativa*)

## 1. Introduction

Nucleo-cytoplasmic transport of diverse proteins and RNA-protein complex takes place across the nuclear pore complex (NPC) [1–5]. The import of proteins containing a nuclear localization signal (NLS) into the nucleus is one of the best characterized transport pathways and is initiated by recognition of the NLS by the NLS receptor in cytoplasm. The NLS receptor exists as a heterodimer of importin  $\alpha$  and importin  $\beta$ . Importin  $\alpha$  recognizes and binds to the NLS of a nuclear protein, forming a stable pore-targeting complex (PTAC) in the cytoplasm [6–8], while importin  $\beta$  interacts directly with nucleoporins that contain FXFG or GLFG repeats, docking the PTAC to the cytoplasmic face of the NPC [9–11]. Translocation of the docked PTAC into the nucleus is an energy-dependent process, mediated by small GTPase Ran along with a homodimeric factor known as p10 or NTF2 [12–15]. Ran's nucleotide-bound state is regulated by two different proteins, the chromatin-bound exchange factor, RCC1, generating Ran-GTP in the nucleus [16,17], and the cytoplasmic GTPase activating protein, RanGAP1, which depletes Ran-GTP from the cytoplasm [12,18–22]. Ran-GTP binding to importin  $\beta$  has been shown to release importin  $\alpha$ -substrate complex into the nucleus [23–25].

Utilizing both homology search and biochemical ap-

proaches, cDNAs for homologues of importin  $\alpha$  [26–30], importin  $\beta$  [31], Ran [32–34] and Ran binding protein (RanBP1) [35] from diverse plant species have been identified and partially characterized. In addition, plant in vitro assays for the nuclear import of proteins have been developed using permeabilized, evacuated protoplasts from tobacco BY-2 cell [26,36]. In these assays, however, in vitro import did not require exogenous cytosolic extract and ATP, suggesting that all necessary factors are retained within the permeabilized protoplasts in sufficient amount for efficient nuclear import. This makes such an assay unsuitable for elucidating the role of a specific factor in the process of nuclear import of proteins. In order to solve this problem, in our previous work we employed a vertebrate in vitro assay system using digitonin-permeabilized HeLa cells, and successfully demonstrated that the rice importin  $\alpha$ 1, in combination with vertebrate nuclear transport factors, can specifically bind to functional NLS-containing proteins and direct them into the nucleus [30].

Recently we isolated two cDNAs encoding the first known plant importin  $\beta$  homologues, rice importin  $\beta$ 1 and  $\beta$ 2 [31]. Recombinant rice importin  $\beta$ 1 protein showed specific interaction with both the rice importin  $\alpha$ 1 and  $\alpha$ 2 [31]. In the present work, we show that the rice importin  $\beta$ 1 interacts with importin  $\alpha$ , the Ran GTPase and the nuclear envelope, and can mediate the nuclear import of NLS proteins in digitonin-permeabilized HeLa cells.

## 2. Materials and methods

### 2.1. DNA construction

GST-rice importin  $\beta$ 1: a fragment of rice importin  $\beta$ 1 cDNA [31] with an artificial *Eco*RI site at the 5' end was generated by polymerase chain reaction (PCR) and inserted between the *Eco*RI and *Xho*I sites of pGEX-6p-3 (Pharmacia Biotech Inc.) to obtain an in-frame translation fusion with GST.

GST-NLS-GFP: oligonucleotides encoding either the NLS sequence of the SV40 large T-antigen (T-NLS: CTPPKKKRKV) or its point mutant, in which the sixth lysine residue of the T-NLS was replaced by a threonine (Tm-NLS: CTPPKTKRKV), were coupled to the 5' end of the GFP gene by PCR using pS65T-C1 (Clontech Laboratories, Palo Alto, CA) as template DNA. The 5' end primer used for the PCR incorporates an *Eco*RI site at the 5' end followed in frame by the NLS-encoding nucleotide sequence. The 3' end primer incorporates a *Xho*I site at its 5' end. The PCR product was digested with *Eco*RI and *Xho*I restriction enzymes and cloned in frame at the 3' end of GST in pGEX-4T-1 (Pharmacia Biotech Inc.). The 5' end oligonucleotide primers used for the PCR were as follows: (1) T-NLS: 5'-ACGGAATTCTGCACCCCGCCGAAAAAACGCAAGT-GATGAGTAAAGGAGAAGAACTTTTCACTGGA-3'; (2) Tm-NLS: 5'-ACGGAATTCTGCACCCCGCCGAAAAACCAACGCA-AAGTGATGAGTAAAGGAGAAGAACTTTTCACTGGA-3'. The 3' end primer used was 5'-ACGCTCGAGTTATTTGTAGAGCT-

\*Corresponding author. Fax: (81) (298) 38-7073.

E-mail: naoyam@abr.affrc.go.jp

CATCCATGCCATGTGT-3'. The nucleotide sequences encoding T-NLS and Tm-NLS are underlined and the endonuclease restriction sites are indicated by boldface.

## 2.2. Expression and purification of fusion proteins

The fusion proteins were expressed in *Escherichia coli* BL21 by growing in the presence of 0.5 mM IPTG for 4–6 h at 20°C and the proteins were purified essentially according to the manufacturer's instruction (Pharmacia Biotech). All the procedures were carried out at 4°C, and 1 mM EGTA and 2 mM dithiothreitol (DTT) were included in the solutions through out the purification procedures. The GST portion of the GST-rice importin  $\alpha$ 1 and rice importin  $\beta$ 1 fusion protein was cleaved by incubation of the fusion protein bound to glutathione-Sepharose 4B resin with 80 U/ml resin of Preission protease (Pharmacia Biotech) for 4 h at 5°C. The Preission protease is a recombinant fusion protein with GST and hence can be easily removed by glutathione-Sepharose 4B. The purified proteins were concentrated by Millipore Ultrafree-MC (Millipore, Bedford, MA) and finally suspended in 20 mM HEPES buffer (pH 7.3) containing 1 mM EGTA and 1 mM DTT.

Recombinant rice importin  $\alpha$ 1 [30] and mouse Ran [20,37,38] were prepared as described previously.

## 2.3. In vitro binding assay

In vitro protein binding was examined by native gel electrophoresis as described previously [30]. 20 pmol of each protein was mixed in 15  $\mu$ l of transport buffer (TB) (20 mM HEPES, pH 7.3, 110 mM potassium acetate, 2 mM magnesium acetate, 5 mM sodium acetate, 0.5 mM EGTA, 2 mM DTT, 1  $\mu$ g/ml each of aprotinin, leupeptin and pepstatin A) supplemented with 250 mM sucrose and incubated for 1 h at room temperature. 7.5% polyacrylamide gels were run in the presence of 1 mM DTT, 1 mM EGTA and 2 mM magnesium acetate in both the gels and the running buffer. Protein bands on the gels were visualized by staining with Coomassie blue.

## 2.4. Fluorescence labeling of rice importin $\beta$ 1 and nuclear envelope binding assay

Rice importin  $\beta$ 1 was labeled with Cy3 Dye (Amersham Life Science) essentially according to the manufacturer's instruction. The unconjugated dye was removed by Millipore Ultrafree-MC (Millipore, Bedford, MA) with five washes, and the labeled protein was suspended in 20 mM HEPES buffer (pH 7.3) containing 1 mM EGTA and 1 mM DTT.

Evacuolated protoplasts of BY-2 cells were isolated and permeabilized with Triton X-100 as described by Merkle et al. [36]. The Cy3-labeled rice importin  $\beta$ 1, at final concentration of 10  $\mu$ g/ml, either alone or with a four-fold excess of unlabeled rice importin  $\beta$ 1, was mixed with the Triton-permeabilized, evacuolated BY-2 protoplasts and incubated for 20 min at room temperature. After fixation with 3.7% formaldehyde in TB for 30 min, the nuclei were stained with 4',6-diamidino-2-phenylindole dihydrochloride (DAPI) at 0.5  $\mu$ g/ml. Following twice washes with TB, an aliquot of the sample was transferred onto a slide glass and viewed under a microscope equipped with epifluorescence illumination (Olympus AX70). The microscopic images were taken with a MicroMAX digital CCD camera (Princeton Instruments, Trenton, NJ) and processed with the IP Lab image processing system (Scanalytics, Fairfax, VA).

## 2.5. HeLa cell culture and in vitro import assay

HeLa cells were cultured in 5% CO<sub>2</sub> at 37°C in Dulbecco's modified Eagle's essential medium supplemented with 5% fetal bovine serum (Life Technologies). Digitonin-permeabilized cells were prepared as described previously [39,40]. 10  $\mu$ l sample solution contained 1  $\mu$ l of GST-NLS-GFP (0.2  $\mu$ g/ $\mu$ l) and appropriate transport factors were diluted with TB. For the nuclear binding assay, the incubation was performed on ice for 20 min in the presence of rice importin  $\alpha$ 1 (6 pmol) and rice importin  $\beta$ 1 (6 pmol), the concentration of which was adjusted with TB containing 2% BSA. For the nuclear import assay, the incubation was performed at 25°C for 20 min in the presence of rice importin  $\alpha$ 1 (12 pmol), rice importin  $\beta$ 1 (3 pmol), mouse Ran-GDP (42 pmol), 1 mM ATP, ATP regeneration system (20 units/ml creatine phosphokinase, 5 mM creatine phosphate), and 1 mM GTP, the concentration of which was adjusted with TB containing 2% BSA. After incubation, cells were fixed with 3.7% formaldehyde in TB. NLS-GFP was detected by Axiophot microscopy (Carl Zeiss).

## 2.6. Miscellaneous

SDS-polyacrylamide gel electrophoresis (SDS-PAGE) was performed by the method of Laemmli [41]. Concentration of proteins were determined by the method of Bradford [42] using Bio-Rad dye reagent (Bio-Rad) and BSA as the standard.

## 3. Results and discussion

### 3.1. Complex assembly of importin $\alpha$ , $\beta$ and NLS substrate

The NLS of a nuclear protein to be imported is recognized and complexed in cytoplasm by NLS receptor consisting importin  $\alpha$  and  $\beta$ , thereby forming PTAC [6–8]. To address the ability of rice importin  $\beta$ 1 to bind to importin  $\alpha$  and assemble a PTAC in the presence of NLS substrate, an in vitro binding assay was carried out using native gel electrophoresis. On a native gel, complex formation between two proteins gives a new band with mobility different from either of the proteins alone. GST-T-NLS-GFP (T-GFP) and GST-Tm-NLS-GFP (Tm-GFP) fusion proteins were used as NLS substrates throughout the study, as positive and negative controls, respectively.

Rice importin  $\beta$ 1, rice import  $\alpha$ 1, T-GFP and Tm-GFP migrate as single bands on a non-denaturing gel (Fig. 1, lanes 1, 2, 3 and 9). A mixture of rice importin  $\beta$ 1 with rice importin  $\alpha$ 1 gave a new, major band of retarded mobility, with some unbound proteins migrating at the position of the controls (Fig. 1, lane 4), indicating a direct binding of the two proteins. Addition of T-GFP (Fig. 1, lane 5), but not Tm-GFP (Fig. 1, lane 7), to the mixture resulted in formation of a large complex with lower mobility on the gel relative to the complex formed either between rice importin  $\beta$ 1 and  $\alpha$ 1 (Fig. 1, lane 4) or between rice importin  $\alpha$ 1 and T-GFP (Fig. 1, lane 6). This indicates that PTAC assembly occurred in the mixture. A mixture of rice importin  $\beta$ 1 and T-GFP did not give any new visible band (Fig. 1, lane 10), suggesting that the PTAC assembly (lane 5) occurred through the rice importin  $\alpha$ 1 simultaneously binding to rice importin  $\beta$ 1 at one site and T-GFP at another.




Fig. 1. Complex assembly of rice importin  $\beta$ 1 with rice importin  $\alpha$ 1 and T-GFP. Native gel electrophoresis showing rice importin  $\beta$ 1 (lane 1), rice importin  $\alpha$ 1 (lane 2), T-GFP (lane 3) and Tm-GFP (lane 9) run separately or as a mixture. The mixture of rice importin  $\beta$ 1 and rice importin  $\alpha$ 1 (lane 4) gives a new band of retarded mobility, indicating binding between the two proteins. In the presence of T-GFP (lane 5), but not Tm-GFP (lane 7), it gives a major, new band of mobility lower than the complex composed of rice importin  $\alpha$ 1 and T-GFP (lane 6). These data show that rice importin  $\beta$ 1 is capable of forming a complex with rice importin  $\alpha$ 1 and T-GFP. No direct interaction between rice importin  $\beta$ 1 and T-GFP was observed (lane 10).




Fig. 2. Association of rice importin  $\beta 1$  with the nuclear envelope of tobacco BY-2 cells. Rice importin  $\beta 1$  was labeled with Cy3 dye and applied, either alone (a,b) or with a four-fold excess of unlabeled rice importin  $\beta 1$  (c,d), to Triton-permeabilized, evacuated protoplasts of BY-2 cells. The Cy3-labeled rice importin  $\beta 1$  accumulated predominantly at the nuclear rim (a). Such accumulation was drastically reduced by the addition of unlabeled rice importin  $\beta 1$  to the nuclear envelope, indicating specific binding of rice importin  $\beta 1$  to the nuclear envelope. Panels b and d show nuclei stained with DAPI.

### 3.2. Association of rice importin $\beta 1$ to plant nuclear envelope

Importin  $\beta$  interacts directly with nucleoporins at the nuclear envelope, docking the PTAC to the cytoplasmic face of the NPC [9–11]. To assess if rice importin  $\beta 1$  can bind to the nuclear envelope, we labeled the rice importin  $\beta 1$  with the fluorescent dye, Cy3, and applied to Triton-permeabilized, evacuated BY-2 protoplasts. As shown in Fig. 2, the rice importin  $\beta 1$  accumulated predominantly at the nuclear rim (Fig. 2a). Such accumulation was rice importin  $\beta 1$ -specific, as addition of a four-fold excess unlabeled rice importin  $\beta 1$  drastically reduced the accumulation of Cy3 Dye at the nuclear rim (Fig. 2c). This observation suggests a specific binding of rice importin  $\beta 1$  to the nuclear envelope, probably via interaction with some yet unidentified nucleoporins.

### 3.3. Interaction between rice importin $\beta 1$ and Ran

The GTPase Ran has been shown to be involved in multiple reactions in the translocation of the PTAC into the nucleus. In its GTP-bound state, Ran has been shown to bind directly to importin  $\beta 1$  and dissociate the importin  $\alpha/\beta$  heterodimer





Fig. 3. Interaction between rice importin  $\beta 1$  and Ran. A: Native gel electrophoresis showing rice importin  $\beta 1$ , Ran-GTP and Ran-GDP run separately or as a mixture. Ran-GDP gives a single band (lane 5), while the Ran-GTP sample appears as doublet bands (lane 1) due to the presence of some GDP-bound form in the sample. The lower band represents Ran-GTP, which is more negatively charged, and the upper band represents Ran-GDP. A mixture of rice importin  $\beta 1$  and the Ran-GTP sample appears as doublet bands (lane 2), indicating interaction between the two proteins. No such interaction was observed with Ran-GDP (lane 4). B: Native gel electrophoresis showing Ran-GTP (lane 2), but not Ran-GDP (lane 3), dissociates the heterodimer of rice importin  $\beta 1$  and rice importin  $\alpha 1$ .

## A: Nuclear-Binding



## B: Nuclear-Import




Fig. 4. Rice importin  $\beta 1$  supports the nuclear import of NLS-GFP in a vertebrate in vitro assay system. A: Rice importin  $\alpha 1$  alone does not dock (c,d), whereas in combination with rice importin  $\beta 1$ , it can dock the T-GFP to the nuclear envelope (a,b). B: Rice importin  $\beta 1$ , in conjunction with rice importin  $\alpha 1$  and mouse Ran-GDP, mediates translocation of T-GFP (a,b), but not Tm-GFP (e,f), into the nucleus. The translocation did not occur in the absence of rice importin  $\beta 1$  (c,d). Panels a, c and e: fluorescence images; panels b, d and f: phase contrast images.

[23–25]. We therefore examined the interaction between rice importin  $\beta 1$  and Ran, as well as the effect of Ran on the interaction between rice importin  $\beta 1$  and rice importin  $\alpha 1$ .

Ran-GDP migrates as single band on the gel (Fig. 3A, lane 5), while Ran-GTP sample appears as doublet bands (Fig. 3A, lane 1). The lower band represents Ran-GTP, which is more

negatively charged, while the upper band represents Ran-GDP. In our Ran-GTP preparation, about 60–70% is in the GTP-bound form, while the other 30–40% is in the GDP-bound form. A mixture of rice importin  $\beta$ 1 with Ran-GTP gave a major band of high mobility (Fig. 3A, lane 2). In contrast, no such new major band was observed in a mixture of rice importin  $\beta$ 1 with Ran-GDP (Fig. 3A, lane 4). This result indicates that rice importin  $\beta$ 1 specifically interacts with Ran-GTP but not with Ran-GDP. There was a new minor band with the same mobility of rice importin  $\beta$ 1/Ran-GTP complex in lane 4. However, this was most likely due to the presence of trace amounts of Ran-GTP in the Ran-GDP sample.

The specific interaction of Ran-GTP with the rice importin  $\beta$ 1 completely dissociated the heterodimer of rice importin  $\alpha$ 1 and rice importin  $\beta$ 1 (Fig. 3B, lane 2). In contrast, no apparent effect was observed with the Ran-GDP (Fig. 3B, lane 3).

### 3.4. Activity of the rice importin $\beta$ 1 in the *in vitro* import assay using digitonin-permeabilized HeLa cells

The PTAC, formed in the cytoplasm by the binding of importin  $\alpha/\beta$  heterodimer to NLS protein, docks at the cytoplasmic surface of NPC via binding of importin  $\beta$  to nucleoporins. The PTAC is then transported as a single entity into the nucleus in a process mediated by the GTPase Ran. To assess the functional activity of the rice importin  $\beta$ 1 in the process of nuclear import of proteins, we performed an *in vitro* nuclear import assay using digitonin-permeabilized HeLa cells.

In the nuclear binding assay as shown in the Fig. 4A, the rice importin  $\alpha$ 1 alone was not sufficient to direct the substrate to the nuclear rim (Fig. 4Ac). However, addition of the rice importin  $\beta$ 1 to the mixture resulted in efficient accumulation of T-GFP at the nuclear rim (Fig. 4Aa).

In the presence of the rice importin  $\beta$ 1 in the transport solution, T-GFP was translocated efficiently into the nucleus (Fig. 4Ba). The translocation of the substrate was rice importin  $\beta$ 1-dependent (Fig. 4Bc) and NLS-specific (Fig. 4Be).

In summary, we have shown here that rice importin  $\beta$ 1 shares most of the molecular interactions common to all known importin  $\beta$ s from vertebrates and yeast, specifically, interaction with importin  $\alpha$ , the nuclear envelope and Ran-GTP. In addition, the *in vitro* transport assay demonstrated that rice importin  $\beta$ 1, in conjunction with importin  $\alpha$  and Ran factor, can mediate the import of NLS protein into the nucleus. These data strongly suggest that rice importin  $\beta$ 1 functions as a component of the NLS receptor in plant cells.

**Acknowledgements:** This work was supported by the Special Coordination Fund for Promoting Science and Technology, Enhancement of Center-of-Excellence, to the National Institute of Agrobiological Resources from the Science and Technology Agency of Japan, Grant-in-Aid for Scientific Research C (No. 0868076) from the Japanese Ministry of Education, Science, Sports and Culture (N.I., Y.Y.), and in part by a fund from the Ministry of Agriculture, Forestry and Fisheries of Japan (BDP-98-I-1-2).

## References

- [1] Yoneda, Y. (1996) Trends Microbiol. 4, 1–2.
- [2] Görlich, D. (1997) Curr. Opin. Cell Biol. 9, 412–419.
- [3] Ullman, K.S., Powers, M.A. and Forbes, D.J. (1997) Cell 90, 967–970.
- [4] Imamoto, N., Kamei, Y. and Yoneda, Y. (1998) Eur. J. Histochem. 42, 9–20.
- [5] Mattaj, I.W. and Englmeier, L. (1998) Annu. Rev. Biochem. 67, 265–306.
- [6] Imamoto, N., Shimamoto, T., Takao, T., Tachibana, T., Kose, T., Matsubae, M., Sekimoto, T., Shimonishi, Y. and Yoneda, Y. (1995) EMBO J. 14, 3617–3626.
- [7] Imamoto, N., Tachibana, T., Matsubae, M. and Yoneda, Y. (1995) J. Biol. Chem. 270, 8559–8565.
- [8] Görlich, D., Kostka, S., Kraft, K., Dingwell, C., Laskey, R.A., Hartmann, E. and Prehn, S. (1995) Curr. Biol. 5, 383–392.
- [9] Iovine, M.K., Watkins, J.L. and Wente, S.R. (1995) J. Cell Biol. 131, 1699–1713.
- [10] Kraemer, D.M., Strambio-de-Castillia, C., Blobel, G. and Rout, M.P. (1995) J. Biol. Chem. 270, 19017–19021.
- [11] Radu, A., Moore, M.S. and Blobel, G. (1995) Cell 81, 215–222.
- [12] Melchior, F.B., Paschal, J., Evans, J. and Gerace, L. (1993) J. Cell Biol. 123, 1649–1659.
- [13] Moore, M.S. and Blobel, G. (1993) Nature 365, 661–663.
- [14] Moore, M.S. and Blobel, G. (1994) Proc. Natl. Acad. Sci. USA 91, 10212–10216.
- [15] Paschal, B.M. and Gerace, L. (1995) J. Cell Biol. 129, 925–937.
- [16] Ohtsubo, M., Okazaki, H. and Nishimoto, T. (1989) J. Cell Biol. 109, 1389–1397.
- [17] Bischoff, F.R. and Ponstingl, H. (1991) Nature 354, 80–82.
- [18] Hopper, A.K., Traglia, H.M. and Dunst, R.W. (1990) J. Cell Biol. 111, 309–321.
- [19] Bischoff, F.R., Klebe, C., Kretschmer, J., Wittinghofer, A. and Ponstingl, H. (1994) Proc. Natl. Acad. Sci. USA 91, 2587–2591.
- [20] Bischoff, F.R. and Ponstingl, H. (1995) Methods Enzymol. 257, 135–144.
- [21] Matunis, M.J., Coutavas, E. and Blobel, G. (1996) J. Cell Biol. 135, 1457–1470.
- [22] Mahajan, R., Delphin, C., Guan, T., Gerace, L. and Melchior, F. (1997) Cell 88, 97–107.
- [23] Rexach, M. and Blobel, G. (1995) Cell 83, 683–692.
- [24] Görlich, D.N.P., Kutay, U., Aebi, U. and Bischoff, F.R. (1996) EMBO J. 15, 5584–5594.
- [25] Chi, N.C., Adam, E.J.H. and Adam, S.A. (1997) J. Biol. Chem. 272, 6818–6822.
- [26] Hicks, G.R., Smith, H.M.S., Lobreaux, S. and Raikhel, N.V. (1996) Plant Cell 8, 1337–1352.
- [27] Ballas, N. and Citovsky, V. (1997) Proc. Natl. Acad. Sci. USA 94, 10723–10728.
- [28] Shoji, K., Iwasaki, T., Matsuki, R., Uchimiya, H., Miyao, M. and Yamamoto, N. (1998) Genes 212, 279–286.
- [29] Iwasaki, T., Matsuki, R., Shoji, K., Sanmiya, K., Miyao, M. and Yamamoto, N. (1998) FEBS Lett. 428, 259–262.
- [30] Jiang, C.J., Imamoto, N., Matsuki, R., Yoneda, Y. and Yamamoto, N. (1998) J. Biol. Chem. 273, 24083–24087.
- [31] Matsuki, R., Iwasaki, T., Shoji, K., Jiang, C.J. and Yamamoto, N. (1998) Plant Cell Physiol. 39, (8) 879–884.
- [32] Ach, R.A. and Gruissem, W. (1994) Proc. Natl. Acad. Sci. USA 91, 5863–5867.
- [33] Merkle, T., Haizel, T., Matsumoto, T., Harter, K., Dallmann, G. and Nagy, F. (1994) Plant J. 6, 555–565.
- [34] Saalbach, G. and Christov, V. (1994) Plant Mol. Biol. 24, 969–972.
- [35] Haizel, T., Merkle, T., Pay, A., Fejes, E. and Nagy, F. (1997) Plant J. 11, 93–103.
- [36] Merkle, T., Leclerc, D., Marshallsay, C. and Nagy, F. (1996) Plant J. 10, 1177–1186.
- [37] Melchior, F., Sweet, D.J. and Gerace, L. (1995) Methods Enzymol. 257, 279–291.
- [38] Sakamoto, K. and Nagatani, A. (1996) Plant J. 10, 859–868.
- [39] Adam, S.A., Sterne-Marr, R. and Gerace, L. (1990) J. Cell Biol. 111, 807–816.
- [40] Miyamoto, Y., Imamoto, N., Sekimoto, T., Tachibana, T., Seki, T., Tada, S., Enomoto, T. and Yoneda, Y. (1997) J. Biol. Chem. 272, 26375–26381.
- [41] Laemmli, U.K. (1970) Nature 227, 680–685.
- [42] Bradford, J.M. (1976) Anal. Biochem. 72, 248–254.