

## Erratum and corrigendum

---

FEBS 21443

Erratum to: **The catalytic mechanism of amidase also involves nitrile hydrolysis** (FEBS 21159)

[*FEBS Letters* 439 (1998) 325–328]<sup>1</sup>

Michihiko Kobayashi\*, Masahiko Goda, Sakayu Shimizu

*Division of Applied Life Sciences, Graduate School of Agriculture, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan*

On p. 326, last line, left column, 8.15 mmol should read 8.15  $\mu$ mol.

The publisher apologizes for this error.

---

\*Corresponding author. Fax: (81) (75) 753-6128.

<sup>1</sup> PII S0014-5793(98)01406-9

---

FEBS 21495

Corrigendum to: **GD1 $\alpha$ -replica peptides functionally mimic GD1 $\alpha$ , an adhesion molecule of metastatic tumor cells, and suppress the tumor metastasis** (FEBS 21260)

[*FEBS Letters* 441 (1999) 20–21]<sup>1</sup>

Dai Ishikawa<sup>a</sup>, Hironori Kikkawa<sup>b</sup>, Koichi Ogino<sup>a</sup>, Yoshio Hirabayashi<sup>c</sup>, Naoto Oku<sup>b</sup>, Takao Taki<sup>a,\*</sup>

<sup>a</sup>*Cellular Technology Institute, Otsuka Pharmaceutical Co. Ltd., 463-10 Kagasuno, Kawauchi, Tokushima 771-0192, Japan*

<sup>b</sup>*Department of Radiobiochemistry, School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Shizuoka 422-8526, Japan*

<sup>c</sup>*Institute of Physical and Chemical Research, RIKEN, Hirosawa 2-1, Wako, Saitama 351-0198, Japan*

The authors wish to inform readers that in section 2.9 of the above-mentioned article (page 21, line 13), and in the legend of Table 2 (page 24, line 1), the peptide solution volume should read 200  $\mu$ g/0.1 ml.

---

\*Corresponding author. Fax: (81) (886) 65-5662.

E-mail: t\_taki@research.otsuka.co.jp

<sup>1</sup> PII S0014-5793(98)01511-7













































































































