

Erratum and corrigendum

FEBS 21443

Erratum to: **The catalytic mechanism of amidase also involves nitrile hydrolysis** (FEBS 21159)

[*FEBS Letters* 439 (1998) 325–328]¹

Michihiko Kobayashi*, Masahiko Goda, Sakayu Shimizu

Division of Applied Life Sciences, Graduate School of Agriculture, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

On p. 326, last line, left column, 8.15 mmol should read 8.15 μ mol.

The publisher apologizes for this error.

*Corresponding author. Fax: (81) (75) 753-6128.

¹ PII S0014-5793(98)01406-9

FEBS 21495

Corrigendum to: **GD1 α -replica peptides functionally mimic GD1 α , an adhesion molecule of metastatic tumor cells, and suppress the tumor metastasis** (FEBS 21260)

[*FEBS Letters* 441 (1999) 20–21]¹

Dai Ishikawa^a, Hironori Kikkawa^b, Koichi Ogino^a, Yoshio Hirabayashi^c, Naoto Oku^b, Takao Taki^{a,*}

^a*Cellular Technology Institute, Otsuka Pharmaceutical Co. Ltd., 463-10 Kagasuno, Kawauchi, Tokushima 771-0192, Japan*

^b*Department of Radiobiochemistry, School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Shizuoka 422-8526, Japan*

^c*Institute of Physical and Chemical Research, RIKEN, Hirosawa 2-1, Wako, Saitama 351-0198, Japan*

The authors wish to inform readers that in section 2.9 of the above-mentioned article (page 21, line 13), and in the legend of Table 2 (page 24, line 1), the peptide solution volume should read 200 μ g/0.1 ml.

*Corresponding author. Fax: (81) (886) 65-5662.

E-mail: t_taki@research.otsuka.co.jp

¹ PII S0014-5793(98)01511-7