

Isolation and characterization of a calendic acid producing (8,11)-linoleoyl desaturase¹

Kathrin Fritsche^{a,2}, Ellen Hornung^{a,2}, Nicola Peitzsch^a, Andreas Renz^b, Ivo Feussner^{a,*}

^aLeibniz Institut für Pflanzenbiochemie, Weinberg 3, D-06120 Halle, Germany

^bBASF AG, ZHPIT - A 030, Pflanzenbiotechnologie, D-67056 Ludwigshafen, Germany

Received 4 October 1999; received in revised form 29 October 1999

Edited by Ulf-Ingo Flügge

Abstract For the biosynthesis of calendic acid a (8,11)-linoleoyl desaturase activity has been proposed. To isolate this desaturase, PCR-based cloning was used. The open reading frame of the isolated full-length cDNA is a 1131 bp sequence encoding a protein of 377 amino acids. For functional identification the cDNA was expressed in *Saccharomyces cerevisiae*, and formation of calendic acid was analyzed by RP-HPLC. The expression of the heterologous enzyme resulted in a significant amount of calendic acid presumably esterified within phospholipids. The results presented here identify a gene encoding a new type of (1,4)-acyl lipid desaturase.

© 1999 Federation of European Biochemical Societies.

Key words: Acyl-group desaturase; Seed oil; *Calendula officinalis*; *Saccharomyces cerevisiae*

1. Introduction

The most common octadecatrienoic fatty acid in plants is α -linoleic acid (9Z,12Z,15Z-18:3), which is the main constituent of chloroplastic membranes [1]. Moreover, triacylglycerols from seeds provide an easily accessible source of additional conjugated octadecatrienoic acids having (Z,E,E) or (Z,E,Z) geometries [2]. At least five different regio isomers have been reported within plants with double bond systems in the following positions: (Z,E,Z)- and (E,E,Z)-8,10,12-18:3 and (Z,E,Z)-, (Z,E,E)- and (E,E,Z)-9,11,13-18:3. One of these, calendic acid (8E,10E,12Z-18:3), is the major constituent of the seed oil of *Calendula officinalis*, *Calendula stellata*, *Osteospermum spinescens* and *Osteospermum hyoseroides* [1]. These seed oils are of industrial interest, because the oil is used for cosmetic purposes and may be used as a drying oil in paintings.

A number of chemical mechanisms have been proposed to describe the biosynthesis of conjugated octadecatrienoic acids. These include the formation of an epoxy derivative of linoleic acid as an intermediate [3], the oxidation of linoleic acid [4], the isomerization of linolenic acid [5] and the formation of a radical of linoleic acid due to a lipoxygenase-type reaction [6–8]. However, in a recent publication on the biosynthesis of

α -eleostearic acid (9Z,11E,13E-18:3), which is a regio isomer of calendic acid, it was shown that the conversion of linoleic acid to this conjugated octadecatrienoic acid occurs while this acyl moiety is esterified to phosphatidylcholine [9]. Based on these data one may assume that the conversion of esterified linoleic acid to conjugated octadecatrienoic acids is catalyzed by a new type of (1,4)-linoleoyl acyl lipid desaturase.

To obtain additional information on the biosynthesis of conjugated octadecatrienoic acids we decided to continue the analysis of the biosynthesis of calendic acid (8E,10E,12Z-18:3) in the developing seeds of *C. officinalis*. This species is amenable to such studies, because it is easy to grow, its seeds contain large quantities of calendic acid, and most of the studies on the biosynthesis of conjugated octadecatrienoic acids have been performed with seeds from this plant. Here, we describe the cloning of a new type of (1,4)-acyl lipid desaturase, a (8,11)-linoleoyl desaturase that catalyzes the formation of a conjugated triene, calendic acid.

2. Materials and methods

C. officinalis was grown from seeds in a greenhouse under 16 h of artificially supplemented illumination at a temperature ranging between 22 and 30°C. For all experiments developing seeds at a fully expanded green stage were used.

For RNA isolation 20 g of plant material was ground in liquid nitrogen, 100 ml of extraction buffer I (100 mM Tris-HCl, pH 7.5, 25 mM EDTA, 2% (w/v) laurylsarcosyl, 4 M guanidinium thiocyanate, 5% (w/v) PVPP, 1% (v/v) β -mercaptoethanol) was added and the homogenate was shaken for 15 min. After centrifugation at 4000 \times g for 15 min the floating solid lipid phase was discarded and the remaining liquid phase was extracted twice for 15 min with PCI (phenol:chloroform:isoamyl alcohol, 25:24:1). After centrifugation at 4000 \times g for 15 min the upper hydrophilic phase was loaded on a CsCl cushion (5 M CsCl) of 8 ml and centrifuged at 18°C and 100 000 \times g for 18 h. After discarding the supernatant the RNA precipitate was dried, washed with 70% ethanol and extracted for 15 min with a mixture consisting of 7.5 ml of extraction buffer II (100 mM Tris-HCl, pH 8.8, 100 mM NaCl, 5 mM EDTA, 2% (w/v) SDS) and 10 ml PCI. After centrifugation at 4000 \times g for 15 min the upper hydrophilic phase was washed with chloroform and the remaining RNA was precipitated overnight at 4°C by adding an equal volume of 5 M LiCl. After centrifugation for 60 min at 12 000 \times g at 4°C the precipitate was washed twice with 70% ethanol, dried and dissolved in 500 μ l water. From this total RNA fraction mRNA was isolated using the Poly-Attract-Kit (Promega, Mannheim, Germany).

Single stranded cDNA (ss-cDNA) was isolated from mRNA of developing marigold seeds by reversed transcription with Superscript-II (Gibco BRL, Eggenstein, Germany). This ss-cDNA was used as template for PCR-based cloning. A 470 bp PCR fragment was amplified with the degenerate sense primer A 5'-CCD TAY TTC TCI TGG AAR WWH AGY CAY CG-3' and antisense primer B 5'-CCA RTY CCA YTC IGW BGA RTC RTA RTG-3' derived from the amino acid sequences PYFSWK(Y/I)SHR and HYDS(S/T)EW-(D/N)W, respectively. The PCR reaction was carried out with Tfl-

*Corresponding author. Fax: (49)-345-5582 162.
E-mail: ifeussne@ipb.uni-halle.de

¹ The nucleotide sequence reported in this paper has been submitted to the GenBank/EMBL data bank with accession number AJ245938.

² The first two authors contributed equally to this work.

DNA-Polymerase (Biozym, Hess. Oldendorf, Germany) using an amplification program of 2 min denaturation at 94°C, followed by 10 cycles of 30 s at 94°C, 45 s at 50°C, 1 min at 72°C, followed by 20 cycles of 30 s at 94°C, 45 s at 50°C, 1 min at 72°C (time increment 5 s) and terminated by 2 min extension at 72°C. PCR products of the expected length were cloned in pCR2.1-TOPO (Invitrogen, Carlsbad, CA, USA) and sequenced. The fragment CoDes2.2 was chosen for the isolation of a full-length cDNA clone using the Marathon cDNA amplification kit (Clontech, Heidelberg, Germany). To amplify the 5'- and 3'-ends of CoDes2.2 by PCR specific primers were used: 5'-race: primer C 5'-GTG AGG GAG TGA GAG ATG GGT GTG GTG C-3', primer D 5'-AAC ACA CTT ACA CCT AGT ACT GGA ATT-3'; 3'-race: primer E 5'-TAT TCC AAA CTT CTT AAC AAT CCA CCC G-3', primer F 5'-CAA TTC CAG TAC TAG GTG TAA GTG TGT T-3'. The PCR reaction was carried out according to the supplier's instructions. The fragments were cloned in pCR2.1-TOPO and sequenced. To obtain the full-length cDNA clone by PCR, specific primers of the expected open reading frame of the entire cDNA of 1.2 kb were used for amplification: sense primer G 5'-ATT AGA GCT CAT GGG TGC TGG TGG TCG GAT GTC G-3' and anti-sense primer H 5'-ATT ACT CGA GTG ACA TAC ACC TTT TTG ATT ACA TCT TG-3'. The PCR reaction was carried out with the Expand High Fidelity System (Boehringer, Mannheim, Germany) using an amplification program of 2 min denaturation at 94°C, followed by 10 cycles of 30 s at 94°C, 35 s at 63°C, 2 min at 72°C, followed by 15 cycles of 30 s at 94°C, 35 s at 63°C, 1 min at 72°C (time increment 5 s) and terminated by 2 min extension at 72°C. The fragment was cloned into pGEM-T and the resulting plasmid pCoDes8.11 was sequenced. For expression in *Sacharomyces cerevisiae* the open reading frame of pCoDes8.11 was cloned as a *SacI/XhoI* fragment behind the galactose-inducible promoter *GAL1* into the shuttle vector pYES2 (Invitrogen, Carlsbad, CA, USA) to yield plasmid pYESDes8.11. Plasmid pYESDes8.11 was transformed into the yeast strain INVSC1 (Invitrogen, Carlsbad, CA, USA) by the polyethylene glycol method

[10]. The expression of cells harboring plasmid pYESDes8.11 was performed essentially as described before including the exogenous application of linoleic acid [11]. Cells reaching the stationary phase were washed with minimal medium, harvested by centrifugation and stored at -20°C until they were analyzed.

For analysis of the calendic acid content of transformed yeast cells, the cell pellets were disrupted using a sonifier tip and lipids were extracted with hexane/isopropanol according to [12]. After recovery of the organic phase, solvents were evaporated by vacuum and the lipids were reconstituted in methanol, then either directly analyzed by HPLC or subjected to alkaline hydrolysis as described [13]. HPLC analysis of the free fatty acid derivatives was carried out by RP-HPLC as described before [13]. Calendic acid eluted with a retention time of 11.8 min exhibiting the characteristic conjugated UV spectrum with a maximum at 268 nm [14]. Linoleic acid eluted with a retention time of 12.7 min exhibiting maximal absorbance at 202 nm. Co-injections with authentic standards were performed. The standard of calendic acid was prepared from the triacylglycerol fraction of lipid extracts from marigold seeds by alkaline hydrolysis. Alternatively, the corresponding fatty acid methyl esters were analyzed by gas chromatography/mass spectrometry (GC/MS). For derivation the free fatty acid derivatives were incubated with diazomethane. GC/MS analysis was performed with a Finigan GCQ GC/MS system equipped with a capillary Rtx-5 column (5% diphenyl-95% polydimethyl siloxane, 30 m×0.25 mm; 0.25 µm coating thickness; Restek, Germany). Helium was used as carrier gas (40 cm/s). An electron energy of 70 eV, an ion source temperature of 150°C, and a temperature of 275°C for the transfer line were used. The samples were measured in the EI mode, and the splitless injection mode (opened after 1 min) with an injector temperature of 250°C. The temperature gradient was 60°C–110°C at 25°C/min, 110°C for 1 min, 110°C–270°C at 10°C/min, and 270°C for 10 min. Calendic acid eluted with a retention time of 17.16 min, linoleic acid eluted with a retention time of 15.92 min, and α -linolenic acid eluted with a retention time of 15.97 min.

CoDes8,11	MGAGGR MSDPSE GGK ----NILERVFPVDP P --FTLS DLKKAIP THCFERSVIRSSYYV VHDL 55
CaAce	MGGGGR GR--TS QK ----PLMERVSVDP P --FTVSD LKQAIP PHCFKRSVIRSSYY VHDA 53
CpEpo	MGAGGR GR--TSE K ----SVMERVSVDPV T FSLS ELKQAIP PHCFQRSVIRSSYYV VQDL 54
BoDes12	MGGGGR MPVPTKGKSKSDV FQ RVSEK P FTVGD LKKVIP PHCFQRSVLHSF SVVYDL 60
CoDes8,11	IVAYVFYLLANTYIPLIPT PLAYLAWPVY WFCQAS IL TGLWVIG HECGHHAF SDYQL IDD 115
CaAce	IIAYIFYFLADKYIPILPAP PLAYLAWPLY WFCQAS IL TGLWVIG HECGHHAF SDYQ WVDD 113
CpEpo	IIAYIFYFLANTYIPTLPTSLAY LAWPVY WFCQAS VL TGLWILG HECGHHAF SNYT WFDD 114
BoDes12	VIAALFFYTASRYIHLQ PHPLSYVAWPLY WFCQGS VL TGVWVIA HECGHHAF SDYQ WLDD 120
CoDes8,11	IVGFVLH SALLTPYFSWKY SHR NHH ANTNSLDNDEVYIPKRKSKVKIYSK LLNNPPGR VF 175
CaAce	TVGFILH SFLMTPYFSWKY SHR NHH ANTNSLDNDEVYIPKSKAKVALYK VLNHPPGR LL 173
CpEpo	TVGFILH SFLLT PYFSWK FSHRNHH SNTSSIDNDEVYIPKSKSKLARIYK LLNNPPGR LL 174
BoDes12	TVG LLHLSALLVPYFSWKY SHR RHH SNTGSLERDEVFV PKRSGISWSSEY LNN PPGR VL 180
CoDes8,11	TLVFR LT LG FPLYLL TNISGKKYGRFAN HFDPM SP I FND RER VQ LLSD FGL LA VFYA IK 235
CaAce	IMFIT TLG F PLYLL TNISGKKYERFAN HFDPM SP I FKERERFQ VLLSD LGL LA VLYGVK 233
CpEpo	VLIIM FT LG FPLYLL TNISGKKYDRFAN HFDPM SP I FKERERFQ VFLSD LGL LA VFYGIK 234
BoDes12	VLLVQ LT LGW PLYLMF NVSGRPYDR FACH FD PKSP IYND RER LQ IYIS DAGIVAVMYGLY 240
CoDes8,11	LLVA AKGA AWVINMYAIPVLGVSVFFVLIT YLHHT LSLPHYD STEWN WIKGAL STIDRD 295
CaAce	LVA AKGA AWVT CIY GPVLGVFIFFDIIT YLHHT LSLPHYD SEWN WLRGAL STIDRD 293
CpEpo	VAV ANKGA AWVACMYGV PVLGV TF FDVIT FLHHTQ SSPHYD STEWN WIRGAL SAID RD 294
BoDes12	RLVA AKGVA WVCYGV P LLV VNG FLVLIT YLQHTQ PSLPHYD SEWD WLKGAL ATVDRD 300
CoDes8,11	FGFLNRV FHDV TH THVHL LSYIPHYHAK EARDAIK PVLGEY YKIDRT PIFKAM YREAK 355
CaAce	FGFLNSVL HDV TH THVMH LSYIPHYHAK EARDAINT VLGDFY KIDRT PI LKAM W REAK 353
CpEpo	FGFLNSV FHDV TH THVMH LSYIPHYHAK EARDAIK PI L GD FY MI DRTP IL KAM W REGR 354
BoDes12	YGFLNKVL HNITD TH THVH LS FTM PHYHAK EATKA KPI L GD Y Q CDRT PI VFKAM Y REVK 360
CoDes8,11	ECIYIEPDEDSEHKG VFWY --HK M 377
CaAce	ECIFIEPEKGRESKG VY WY--NK F 375
CpEpo	ECMYIEPDS--KLKG VY WY--HK L 374
BoDes12	ECIYVEADEGDNKG VFWY KN KL 383

Fig. 1. Sequence alignment of the (8,11)-linoleoyl desaturase (CoDes8,11) from marigold with the Δ^{12} fatty acid acetylenase from *Crepis alpina* (CaAce, accession number: 081931), the Δ^{12} fatty acid epoxigenase from *Crepis palaestina* (CpEpo, accession number: 065771) and the Δ^{12} acyl lipid desaturase from borage (BoDes12, accession number: 082729). The three characteristic and highly conserved histidine regions are indicated by boxes, and identical amino acids are marked as bold letters.




Fig. 2. Phylogenetic tree analysis of plant acyl lipid desaturases. Alignments were generated by the CLUSTAL-X program, and the phylogenetic tree was made with 'TreeView'. ¹Y16283, ²Y16285, ³U22378, ⁴L26296, ⁵O82729, ⁶Y16284, ⁷AJ224161, ⁸U79010, ⁹D16547, ¹⁰D88536, ¹¹P48627, ¹²P48628, ¹³L22961, ¹⁴L22964.

3. Results

3.1. PCR-based cloning and isolation of a full-length cDNA

For PCR-based cloning degenerate primers, deduced from conserved regions of acyl lipid desaturases, were synthesized according to [11,15,16]. The template used was ss-cDNA from *C. officinalis*, which was reverse-transcribed from mRNA of developing seeds at a fully expanded green stage. PCR products of the expected length were cloned and sequenced. Data base searches and alignments with these fragments indicated similarities to plant acyl lipid desaturases with different regio specificities. Except for one (CD2.2), all were classified either as putative members of the classical Δ^{12} desaturases or as putative sphingolipid desaturases based on identities more than 75% [16]. However, the 470 bp fragment CD2.2 exhibited highest identities to the Δ^{12} fatty acid acetylenase from *Crepis alpina* and the Δ^{12} fatty acid epoxygenase from *Crepis palaestina* [15]. To isolate the full-length cDNA clone, PCR with specific primers was used to amplify the 5'- and 3'-ends of CD2.2. The fragments were cloned and sequenced. With specific primers for the expected open reading frame the entire cDNA of 1.2 kb was amplified by PCR and cloned into pGEM-T. The resulting fragment CoDes8.11 was sequenced. The full-length cDNA had a length of 1131 bp encoding a protein of 377 amino acids with a calculated molecular mass of 43.6 kDa. This desaturase sequence showed highest identities with the Δ^{12} acetylenase from *C. alpina* (74%) [15], the Δ^{12} epoxygenase from *C. palaestina* (73%) [15], and the Δ^{12} desaturase from borage (62%) [17] (Fig. 1). The three characteristic and highly conserved histidine regions of the active site of the Δ^{12} acyl lipid desaturases are indicated by boxes. These findings were further supported by phylogenetic analysis ac-

cording to [17]. As shown in Fig. 2, the newly identified (8,11)-linoleoyl desaturase groups with the Δ^{12} acyl lipid desaturase family. However, it forms one subfamily with the Δ^{12} acetylenase and the Δ^{12} epoxygenase. All members of this subfamily are clearly separated from the classical Δ^{12} desaturases and 12-hydroxylases.

3.2. Functional expression in *S. cerevisiae* and fatty acid analysis

In order to elucidate the catalytic properties of this new desaturase, CoDes8.11 was expressed in yeast cells under the control of the galactose-inducible promoter *GALI*. Since yeast cells do not contain dienoic fatty acids as substrates for the (8,11)-linoleoyl desaturase [8], the expression was performed with supplementation of linoleic acid. Galactose-induced yeast cells were used for the analysis either of fatty acid methyl esters by GC/MS or of free fatty acids by HPLC. The methyl ester of calendic acid was separated from the classical dienoic and trienoic fatty acids by GC/MS. It eluted with a retention time of 17.16 min, whereas the methyl ester of linoleic acid eluted with a retention time of 15.92 min, and that of α -linolenic acid eluted with a retention time of 15.97 min (data not shown). However, using this method calendic acid was only identified by its characteristic retention time, because the corresponding mass spectra of the methyl esters of calendic acid and α -linolenic acid showed no characteristic fragmentation patterns when compared against each other. The spectra differed only in their relative intensities of the obtained fragments (data not shown). Therefore, we decided to analyze the formation of calendic acid within yeast cells overexpressing CoDes8.11 by RP-HPLC, because of the characteristic UV spectrum of calendic acid and the substantial loss of cal-




Fig. 3. Reversed-phase HPLC analysis of calendric acid either isolated from the triacylglycerol fraction of marigold seeds or isolated from yeast cells transformed with the marigold (8,11)-linoleoyl desaturase. The lipids were extracted and analyzed by HPLC as described [12]. Calendric acid and linoleic acid were characterized by co-elution of authentic standards. The insets show the UV spectra of calendric acid exhibiting a maximum at 268 nm (maximum for conjugated triene system).

endic acid during derivation into its methyl ester for GC/MS analysis. As a standard calendric acid was isolated from the triacylglycerol fraction of marigold seeds and analyzed by RP-HPLC (Fig. 3, upper panel). The retention time of calendric acid was 11.8 min and as shown in the inset it exhibited a characteristic UV spectrum of conjugated trienes with the three relative maxima at 262 nm, 268 nm and 282 nm [14]. Yeast cells harboring only the empty vector synthesized no calendric acid (data not shown). However, *S. cerevisiae* cells expressing the CoDes8.11 cDNA were capable of forming a fatty acid which co-elutes with calendric acid at 11.8 min and exhibits the characteristic UV spectrum of conjugated trienes (Fig. 3, lower panel, first peak at 11.8 min). The second peak, eluting close behind calendric acid at 12.7 min, was identified as linoleic acid. This indicates that the newly isolated cDNA encodes a desaturase

which is capable of converting linoleic acid into calendric acid upon expression in yeast cells. Moreover, no calendric acid was found in the fraction of free fatty acids (data not shown). This supports the idea that calendric acid formation may occur while the acyl moiety is esterified to phosphatidylcholine, as has been suggested before [9].

4. Discussion

Here, we report the isolation and characterization of a cDNA encoding a new type of desaturase catalyzing a (1,4)-dehydrogenation reaction. The cDNA encoding an (8,11)-linoleoyl desaturase was cloned by a PCR-based approach as has been described before [11,16]. So far, this cDNA could not be identified by EST sequencing of cDNA libraries from




Fig. 4. Speculative mechanism for (8,11)-linoleoyl desaturases (adapted from [20]) based on the proposed catalytic mechanism for the Δ^9 desaturase from castor bean [21]. Not shown is the resting level of the (8,11)-linoleoyl desaturase harboring two diferrous Fe(II) atoms. To reach this level an additional reduction by two more electrons is required.

developing marigold seeds, indicating that the expression of this (8,11)-linoleoyl desaturase gene is either very low or developmentally regulated. The deduced protein shared highest homologies with the recently identified Δ^{12} acetylenase and Δ^{12} epoxigenase from *Crepis* [15]. All three enzymes convert the double bond in the Δ^{12} position of linoleic acid esterified to phosphatidylcholine into groups with new functionalities. In addition, within a phylogenetic analysis these three enzymes could be clearly separated from the other members of the Δ^{12} acyl lipid desaturase family (Fig. 2).

As shown by heterologous expression in yeast cells, the enzyme converts linoleic acid into a conjugated octadecatrienoic acid (Fig. 3). Based on the observation that linoleic acid is the precursor of calendic acid one may propose a new reaction mechanism for this Δ^{12} -type desaturase. Whereas within the classical Δ^{12} acyl lipid desaturases a (1,2)-*syn* dehydrogenation at the carbon atoms C-12 and C-13 of oleic acid may occur [18], one may assume in the case of this new desaturase a (1,4)-dehydrogenation at the carbon atoms C-8 and C-11 of linoleic acid as shown in a speculative reaction scheme in Fig. 4. Within linoleic acid the elimination of protons at these two positions might be sterically favored, because the carbon atoms at the positions C-8 and C-11 are localized near each other in the three-dimensional structure of the fatty acid molecule. For the conversion of linoleic acid into α -eleostearic acid one may assume a homologous enzyme catalyzing a (1,4)-dehydrogenation at the carbon atoms C-14 and C-11 of linoleic acid. Structural determinants which point to this new regio specificity in proton abstraction can hardly be determined by sequence comparison. However, conjugated octadecatrienoic acids are widespread in plant seed oils [19]. Therefore, sequence comparisons between this enzyme and novel yet to be isolated (1,4)-acyl lipid desaturases might help to identify structural characteristics or differences in these enzymes which lead to the observed regio and stereochemical specificities within this subfamily of Δ^{12} desaturases. Even the differences between this (8,11)-linoleoyl desaturase catalyzing the formation of calendic acid (8*E*,10*E*,12*Z*-18:3) and the corresponding (8,11)-linoleoyl desaturase catalyzing the formation of jacaric acid (8*Z*,10*E*,12*Z*-18:3) from *Jacaranda mimosifolia* [1] is an interesting but open question. With the isolation of this enzyme the first example has been given for a (1,4)-dehydrogenation reaction catalyzed by a desaturase on a dienoic fatty acid. However, this reaction has been suggested recently for a slightly different substrate as well, the

conversion of a monoenoic fatty acid into a conjugated fatty acid within the biosynthesis of bombykol, a sex pheromone of the female moth *Bombyx mori* [20].

Acknowledgements: The authors are grateful to S. Barth and A. Zeuner for expert technical assistance. The authors thank M.A. Fuller and Prof. Dr. C. Wasternack for critical reading of the manuscript. This work was supported by the BASF AG.

References

- [1] Gunstone, F.D., Harwood, J.L. and Padley, F.B. (1994) The Lipid Handbook, Chapman and Hall, London.
- [2] Tulloch, A.P. (1982) Lipids 17, 544–550.
- [3] Gunstone, F.D. (1965) Chem. Ind. 1965, 1033–1034.
- [4] Hamberg, M. (1992) Biochem. Biophys. Res. Commun. 188, 1220–1227.
- [5] Wise, M.L., Hamberg, M. and Gerwick, W.H. (1994) Biochemistry 33, 15223–15232.
- [6] Noda, M. (1984) in: Structure, Function and Metabolism of Plant Lipids (Siegenthaler, P.-A. and Eichenberger, W., Eds.), pp. 29–32, Elsevier, Amsterdam.
- [7] Crombie, L. and Holloway, S.J. (1984) J. Chem. Soc. Chem. Commun. 953–955.
- [8] Crombie, L. and Holloway, S.J. (1985) J. Chem. Soc. Perkin Trans. I, 2425–2434.
- [9] Liu, L., Hammond, E.G. and Nikolau, B.J. (1997) Plant Physiol. 113, 1343–1349.
- [10] von Pein, M. (1992), Ph.D. Thesis, Heinrich Heine-Universität, Düsseldorf.
- [11] Girke, T., Schmidt, H., Zähringer, U., Reski, R. and Heinz, E. (1998) Plant J. 15, 39–48.
- [12] Kohlmann, M., Bachmann, A., Weichert, H., Kolbe, A., Balkenhohl, T., Wasternack, C. and Feussner, I. (1999) Eur. J. Biochem. 260, 885–895.
- [13] Feussner, I., Balkenhohl, T.J., Porzel, A., Kühn, H. and Wasternack, C. (1997) J. Biol. Chem. 272, 21635–21641.
- [14] Chrisholm, M.J. and Hopkins, C.Y. (1962) J. Org. Chem. 27, 3137–3139.
- [15] Lee, M. et al. (1998) Science 280, 915–918.
- [16] Sperling, P., Zähringer, U. and Heinz, E. (1998) J. Biol. Chem. 273, 28590–28596.
- [17] Napier, J.A., Sayanova, O., Sperling, P. and Heinz, E. (1999) Trends Plant Sci. 4, 2–4.
- [18] Heinz, E. (1993) in: Lipid Metabolism in Plants (Moore, Jr., T.S., Ed.), pp. 33–89, CRC Press, London.
- [19] van de Loo, F.J., Fox, B.G. and Somerville, C. (1993) in: Lipid Metabolism in Plants (Moore, Jr., T.S., Ed.), pp. 91–126, CRC Press, London.
- [20] Svatos, A., Kalinova, B. and Boland, W. (1999) Insect Biochem. Mol. Biol. 29, 225–232.
- [21] Lindqvist, Y., Huang, W.J., Schneider, G. and Shanklin, J. (1996) EMBO J. 15, 4081–4092.