

Corrigendum

Corrigendum to: The role of conserved extracellular cysteine residues in vasopressin V2 receptor function and properties of two naturally occurring mutant receptors with additional extracellular cysteine residues

[*FEBS Letters* 466 (2000) 101–106]¹

Ralf Schüle^{a,*}, Kerstin Zühlke^a, Alexander Oksche^a, Ricardo Hermosilla^a, Jens Furkert^a,
Walter Rosenthal^{a,b}

^aForschungsinstitut für Molekulare Pharmakologie (FMP), Alfred-Kowalke-Str. 4, D-10315 Berlin, Germany

^bInstitut für Pharmakologie, Freie Universität Berlin, Thielallee 67–73, D-14195 Berlin, Germany

In Fig. 1 of our original article, residue G188 rather than the correct G185 was indicated as the site of the G185C mutation which we constructed. The corrected figure is given below. The authors apologize to the readers.




Fig. 1. Topological model of the human V2 receptor. The one-letter code for amino acids is used. The following putative post-translational modifications are shown: glycosylation at asparagine 22, palmitoylation at cysteine residues 341 and 342. The amino acid replacements for the conserved cysteine residues (C112S, C112A, C192S, C192A) and for the two NDI-causing mutations (G185C, R202C) are indicated. The putative disulfide bridge is depicted between the conserved extracellular cysteine residues C112 and C192.

*Corresponding author. Fax: (49)-30-51 551 291.
E-mail: schuelein@fmp-berlin.de

¹ PII of original article: S0014-5793(99)01764-0.