

Enhanced toxicity of *Bacillus thuringiensis* Cry3A δ -endotoxin in coleopterans by mutagenesis in a receptor binding loop

Sheng-Jiun Wu^a, C. Noah Koller^b, Deborah L. Miller^c, Leah S. Bauer^{b,c}, Donald H. Dean^{a,*}

^aDepartment of Biochemistry, The Ohio State University, 484 W. 12th Ave. Columbus, OH 43210-1292, USA

^bDepartment of Entomology and the Center for Integrated Plant Systems, Michigan State University, East Lansing, MI 48824, USA

^cUSDA Forest Service, North Central Research Station, East Lansing, MI 48823, USA

Received 13 December 1999; received in revised form 10 April 2000

Edited by Marc Van Montagu

Abstract We used site-directed mutagenesis to modify the *Bacillus thuringiensis* cry3A gene in amino acid residues 350–354. Two mutant toxins, A1 (R₃₄₅A,Y₃₅₀F,Y₃₅₁F) and A2 (R₃₄₅A, Δ Y₃₅₀, Δ Y₃₅₁), showed significantly improved toxicity against *Tenebrio molitor* (yellow mealworm). The mutant toxin A1 was also more potent against both *Leptinotarsa decemlineata* (Colorado potato beetle) and *Chrysomela scripta* (cottonwood leaf beetle), while A2 displayed enhanced toxicity only in *L. decemlineata*. Competitive binding assays of *L. decemlineata* brush border membrane vesicles (BBMV) revealed that binding affinities for the A1 and A2 mutant toxins were ca. 2.5-fold higher than for the wild-type Cry3 toxin. Similar binding assays with *C. scripta* BBMV revealed a ca. 5-fold lower dissociation rate for the A1 mutant as compared to that of Cry3A.

© 2000 Federation of European Biochemical Societies.

Key words: *Bacillus thuringiensis*; Cry toxin; δ -Endotoxin; Coleopteran

1. Introduction

The naturally occurring crystal proteins of *Bacillus thuringiensis* (*B.t.*), known as δ -endotoxins or Cry toxins, are insecticidal agents considered safe for users, consumers, and the environment. These proteins are long established for conventional application throughout the world in agricultural, forested, and aquatic ecosystems. Today, insect-resistant plants, genetically engineered to express high concentrations of these proteinaceous toxins, are commercially available. Development of resistance to *B.t.* toxins by target insects, such as the diamondback moth, *Plutella xylostella*, after intensive management with conventionally applied *B.t.* [1], threatens the efficacy of high expression δ -endotoxins produced by plants unless tactics to manage resistance are developed and implemented [2,3]. Detailed structural and biochemical knowledge of *B.t.* δ -endotoxins permits the redesign of these insecticidal proteins to customize host range, alter activity, improve stability, and delay or overcome resistance.

Molecular mechanisms of intoxication by δ -endotoxins are the subject of intensive research, and a complex pathway is gradually emerging, [3–5]. After ingestion by a susceptible insect, crystalline δ -endotoxin is solubilized and proteolyti-

cally cleaved from an inactive protoxin, to an active toxin form within the insect midgut. The activated toxin binds to receptors in the midgut and is believed to integrate into the lipid bilayer of the brush border membrane. Ion channels are formed, causing midgut cells to lose their membrane potential. If the rate of damage to the midgut exceeds the rate of repair, lesions form, bacteria invade the hemocoel, and death results from bacterial septicemia. The δ -endotoxins from *B.t.* comprise a group of over 100 related proteins [5], which were previously categorized by insecticidal activity [6] but currently by amino acid similarity [7]. The spectrum of toxicity for each toxin is relatively narrow, while the collective activity of this group of pesticidal toxins now spans seven orders of insect and several other invertebrate groups including nematodes, mites and protozoans [8].

The Cry3A toxin, produced by *B.t.* var. *tenebrionis* [9,10] and other strains [11,12], is toxic to coleopteran species including the yellow mealworm, *Tenebrio molitor* [9,11], the Colorado potato beetle, *Leptinotarsa decemlineata* [12,13] and the cottonwood leaf beetle, *Chrysomela scripta* [14]. The comparatively simple structure of the Cry3A toxin [15], which is remarkably similar to Cry1Aa [16], makes it a useful model for exploring the structure–function relationship between ligand and receptor. The three-dimensional structure of the Cry toxins consists of three functional domains: (I) a cluster of seven α -helices predicted to be involved in membrane interaction [15]; (II) three antiparallel β -sheets involved in receptor binding [15,17]; and (III) a β -sandwich implicated in receptor binding [18–21] and ion channel activity [22–24] in related Cry toxins. The use of oligonucleotide-directed mutagenesis to engineer or alter the three-dimensional atomic structure of this toxin is proving a powerful probe in the study of mechanisms of toxicity.

Several recent reports indicate that amino acid substitutions in the three surface-exposed loops at the molecular apex of domain II, from residue 291 to 500 of related Cry toxins, affect binding affinity and toxicity [17,18,22,25–29]. In Cry3A, alanine replacements in the three major loops of domain II result in disruption of receptor binding or structural instability [17]. The objectives of this study were to enhance toxicity of the Cry3A δ -endotoxin in beetle larvae by engineering the specific residues, R₃₄₅, Y₃₅₀ and Y₃₅₁, of loop I in the binding domain.

2. Materials and methods

2.1. Construction of loop mutants

The construction of pOS4601 carrying the cry3Aa2 gene (hereafter referred to as cry3A) expressed in *Escherichia coli* was described by

*Corresponding author. Fax: (1)-614-292 6773.

E-mail: dean.10@osu.edu

Abbreviations: *B.t.*, *Bacillus thuringiensis*; BBMV, brush border membrane vesicles

Wu and Dean [17]. We subcloned the *Bam*HI–*Hind*III fragment into M13mp18 RF DNA [30] to target the loop I region in domain II. Site-directed mutagenesis was carried out by the method of Kunkel [31] using the Bio-Rad M13 MutaGene mutagenesis kit; oligonucleotides were kindly provided by T. Yamamoto, Sandoz Agro Corporation (sequences available upon request). After mutagenesis was complete, sequencing of single-strand DNA was carried out by the method of Sanger et al. [32] following the manufacturer's instructions (United States Biochemical). The mutated fragment was then recloned into the vector, pOS4601. Other molecular cloning techniques were performed according to Sambrook et al. [33]. Restriction enzymes and fine chemicals were purchased from Boehringer Mannheim Biochemicals.

2.2. Purification and protease digestion of protoxins

Wild-type and mutant genes were expressed in *E. coli* JM103. Cells were grown for 48 h in 500 ml of LB medium [33] containing 50 µg/ml ampicillin. Crystall extracts were prepared and solubilized in 50 mM sodium carbonate buffer (pH 10.5) as described [34]. The concentration of solubilized protoxin was determined by Coomassie protein assay reagent (Pierce). Protoxin was activated with diluted *T. molitor* midgut juice with a juice protein/protoxin ratio of 1:20 (w/w) for 15 min at 37°C. The activated toxin was used immediately. Protoxin and toxins were analyzed by sodium dodecyl sulfate–polyacrylamide gel (12.5%) electrophoresis (SDS–PAGE) according to Laemmli [35].

2.3. Toxicity bioassays

Bioassays with *T. molitor* larvae were used to screen the mutant toxins for shifts in toxicity when compared to the wild-type Cry3A. The force-feeding bioassay method and data analyses for each mutant toxin screened for toxicity in *T. molitor* are described by Wu and Dean [17]. Two of the mutant toxins, A1 and A2, showed enhanced toxicity in *T. molitor*, and were further assayed in *C. scripta* and *L. decemlineata* larvae, Cry3A-sensitive species in the family Chrysomelidae. The latter bioassays were done by applying serial dilutions of solubilized and activated toxin in a 1 µl droplet of sucrose solution (22.5% final concentration), onto 4 mm poplar or potato leaf disks placed on the surface of 2% agar (Gelcarin) in 24 well tissue culture plates. Second instar *C. scripta* or *L. decemlineata* larvae were placed individually in each well, allowed to feed for 24 h, and placed on fresh poplar or potato foliage, respectively. Mortality was recorded daily. Estimates of median lethal concentrations (LC₅₀) and 95% fiducial limits (FL₉₅) for the A1, A2 mutants, and wild-type Cry3A toxin in *C. scripta* and *L. decemlineata* were determined 96 h after treatment by probit analysis [36]. The lethal dose ratio test was used to determine if there were significant differences in toxicity between toxin pairs, Cry3A:A1, Cry3A:A2, and A1:A2. This test is used when probit lines are not parallel by analyzing the LC₅₀ and slope data [36].

2.4. Preparation of brush border membrane vesicles

Brush border membrane vesicles (BBMV) were prepared from whole, third instar *C. scripta* larvae or midguts of fourth instar *L. decemlineata* larvae. The methods used for making BBMV were described by MacIntosh [37] and English and Readdy [38], respectively, with some modifications. Homogenization buffer (50 mM sucrose and 4 mM Tris at pH 7.5) included a protease inhibitor cocktail with a final concentration in the buffer of 1.0 mM phenylmethylsulfonyl fluoride, 5 mM benzamidine, 5 µg/ml each of pepstatin A, leupeptin and antipain, and 1 µg/ml E-64. The buffer was made fresh and added to each stage of purification. The volume of ice-cold homogenization buffer for use throughout the first precipitation procedure was equal to nine times the weight of tissue to be processed. The initial BBMV pellet was resuspended and reprecipitated with MgCl₂ one time for *L. decemlineata* and twice more for *C. scripta*. The final pellet was resuspended in 0.32 M sucrose, frozen in liquid nitrogen, and stored at –80°C until use. Protein concentration was determined by the BCA protein assay reagent (Pierce). Enrichment, determined by leucine aminopeptidase activity (Sigma assay), was 15–40 times greater than the initial homogenate.

2.5. Iodination of toxins

Protoxins were activated by midgut juice from *T. molitor* and analyzed for proteolysis with silver-stained SDS–PAGE gel. The purity of the activated Cry3A toxin used in the iodination was more than 95% as determined by scanning densitometry. Iodination of toxins was carried out according to Wolfersberger [39], and specific activities of the wild-type Cry3A and mutant toxins ranged from 0.6 to 1.2 µCi/µg protein.

2.6. Competition binding assays

Homologous and heterologous competition binding assays were performed as described by Lee et al. [40] with the following modifications: BBMV (100 µg/ml) were incubated with ¹²⁵I-labeled toxins (2.0 nM) in 100 µl of binding buffer (50 mM Tris, 0.9% NaCl, pH 8.0, containing 0.5% bovine serum albumin and 0.15% Tween 20). Samples were assayed in duplicate for each data point, and binding data were analyzed by the LIGAND program [41]. This program calculates the relative binding affinity based on homologous competition, *K*_{com} [17], and binding site concentration, *B*_{max}, of bound ligand as a best fit of theoretical curves to the experimental data. The term *K*_{com} for the binding constant was derived from competition studies of Cry toxin with BBMV. This term is proportional to IC₅₀ of classical kinetic analysis (median inhibition of binding by competitor).

2.7. Dissociation binding assays and calculation of *k*_{dis}

The ¹²⁵I-labeled wild-type or A1 mutant toxin was incubated with

Table 1
Construction of loop I mutants and toxicity assays to *T. molitor* larvae

Mutant	Construction	Expression	LD ₅₀ ^a	(FL ₉₅) ^a	Relative activity
Wild-type	Cry3A [17] ^b	++	11.4	(8.5–14.9)	1.0
Single mutant	R ₃₄₅ A [17] ^b	++	9.0	(5.9–13.5)	1.3
Single mutant	Y ₃₅₀ A	–			
Single mutant	Y ₃₅₁ A	+	12.2	(7.5–16.3)	0.9
Single mutant	Y ₃₅₀ F	–			
Single mutant	Y ₃₅₁ F	–			
Single mutant	N ₃₅₃ D	++	16.7	(11.6–43.5)	0.7
Single mutant	D ₃₅₄ E	++	7.2	(4.5–11.0)	1.6
Single mutant	D ₃₅₄ N	++	17.2	(12.3–43.2)	0.7
Double mutant	Y ₃₅₀ A,Y ₃₅₁ A [17] ^b	–			
Double mutant	N ₃₅₃ A,D ₃₅₄ A [17] ^b	++	20.0 ^c		<0.6
Double mutant	Y ₃₅₀ F,Y ₃₅₁ F	+ ^d	ND ^e		
Double mutant	ΔY ₃₅₀ ,ΔY ₃₅₁	+ ^d	ND		
Triple mutant	R ₃₄₅ A,Y ₃₅₀ A,Y ₃₅₁ A [17] ^b	+	13.1	(9.1–23.4)	0.9
A1 mutant	R ₃₄₅ A,Y ₃₅₀ F,Y ₃₅₁ F	+	1.0	(0.6–1.4)	11.4
A2 mutant	R ₃₄₅ A,ΔY ₃₅₀ ,ΔY ₃₅₁	+	4.2	(2.2–7.0)	2.7
Block mutant	Y ₃₅₀ A,Y ₃₅₁ A,G ₃₅₂ A				
	N ₃₅₃ A,D ₃₅₄ A				

^aLD₅₀ and FL₉₅ are expressed as µg per larva.

^bThe bioassay data for wild-type Cry3A and mutants have been reported [17]. Those data are listed in this table for comparison.

^cDouble mutant N₃₅₃A,D₃₅₄A was not toxic to *T. molitor* larvae even at the highest dose tested, 20 µg per larva.

^dDouble mutants Y₃₅₀F,Y₃₅₁F and ΔY₃₅₀,ΔY₃₅₁ were degraded soon after solubilization in 50 mM sodium carbonate buffer, pH 10.5.

^eND, not determined.



Fig. 1. Proteolytic digestion of wild-type Cry3A and loop I mutants. Coomassie blue-stained 7.5% SDS-PAGE gel is shown. Lane 1, protein standards (Bio-Rad). Positions of molecular weight markers (in kDa) are indicated on the left; lane 2, wild-type Cry3A toxin purified from *B.t.* var. *tenebrionis*; lane 3, Cry3A purified from *E. coli*; lane 4, Cry3A treated with *T. molitor* midgut juice; lane 5, A1 mutant, R₃₄₅A,Y₃₅₀F,Y₃₅₁F, purified from *E. coli*; lane 6, A1 mutant treated with *T. molitor* midgut juice; lane 7, A2 mutant, R₃₄₅A,ΔY₃₅₀,ΔY₃₅₁, purified from *E. coli*; lane 8, A2 mutant treated with *T. molitor* midgut juice. Each lane contained 8–10 μg of protein.

BBMV to achieve saturation binding. The reaction mixture was then chased with corresponding unlabeled toxin. These dissociation binding assays were similar to the methods of Ihara et al. [42] with the following modification: 2.0 nM ¹²⁵I-labeled toxin (wild-type or mutant toxin) was incubated with 100 μg/ml BBMV at room temperature for 60 min to achieve association binding. After association binding, the mixtures were diluted two-fold in binding buffer containing 0.5 μM corresponding unlabeled toxin. The reaction was stopped at different time points (5–80 min) by centrifugation. The pellet was washed three times with binding buffer and the final pellet was counted in a gamma counter (Beckman). Non-specific binding was subtracted from total binding.

The post-incubation time course of these dissociation reactions was transformed to fit the single exponential equation [43]:

$$y = \{\Delta T\} \{1 - \exp(-k_{\text{dis}}x)\} + z \quad (1)$$

where y is fitted to the concentration of dissociated toxin, and ΔT is the maximum dissociated toxin; k_{dis} is the observed first-order dissociation rate constant; x is time; and z is the correction factor, which is near zero. Dissociation rate constants were determined using Sigma Plot software (Jandel Scientific Co.) to solve the rate equation, Eq. 1.

3. Results

3.1. Expression of loop I mutant proteins

A total of 15 novel mutant toxins were produced by site-directed mutagenesis in the loop I region of domain II and at nearby R₃₄₅ (Table 1). Except for the single mutant Y₃₅₁A, mutagenesis of two surface-exposed Y₃₅₀ and Y₃₅₁ did not produce structurally stable protein (Table 1). Although the

Table 3

Median lethal concentration (LC₅₀) ratio test^a on Cry3A, A1 and A2 mutant proteins for young second instar *C. scripta* and *L. decemlineata*

Toxin comparison	<i>C. scripta</i>		<i>L. decemlineata</i>	
	Ratio	CL ₉₅	Ratio	CL ₉₅
Cry3A:A1 mutant	2.56 ^b	1.35–4.88	1.91 ^b	1.28–2.85
Cry3A:A2 mutant	1.73	0.94–3.21	2.20 ^b	1.44–3.36
A1 mutant:A2 mutant	0.68	0.38–1.20	1.15	0.77–1.73

^aThe lethal dose ratio test [36] was used to determine significance because the probit lines for the three toxins were not parallel.

^bThe LC₅₀s of each toxin comparison are not significantly different at the $P < 0.01$ level of significance if the CL₉₅ interval for their ratio includes 1.

double mutant, Y₃₅₀F,Y₃₅₁F, and the double deletion, ΔY₃₅₀,ΔY₃₅₁, expressed proteins in *E. coli*, these mutant proteins were degraded soon after solubilization in 50 mM sodium carbonate buffer, pH 10.5. To suppress the unstable mutants, a nearby mutation R₃₄₅A was made as reported earlier [17]. Based on these results, the triple mutants A1 (R₃₄₅A,Y₃₅₀F,Y₃₅₁F) and A2 (R₃₄₅A,ΔY₃₅₀,ΔY₃₅₁) were constructed to investigate the functional role of Y₃₅₀ and Y₃₅₁. As a control, the single mutant, R₃₄₅A, was constructed for comparison with triple mutants and wild-type Cry3A δ-endotoxin. These triple mutants produce structurally stable proteins, with a yield of about 5 mg from 500 ml LB medium (Table 1).

To complete the analysis of loop I residues, alanine replacements were generated for N₃₅₃ and D₃₅₄. For comparison, N₃₅₃D, D₃₅₄E and D₃₅₄N were constructed. These mutants produced stable δ-endotoxins with a yield similar to wild-type Cry3A (Table 1). Protein stabilities of wild-type A1 and A2 mutants were determined by incubation with *T. molitor* midgut juice, followed by analysis of the products by SDS-PAGE (Fig. 1).

3.2. Toxicity assays of loop I mutants in coleopterans

Both triple mutants A1 and A2 were significantly more toxic to larvae of *T. molitor* than the wild-type Cry3A (Table 1). The A1 mutant, with phenylalanine substitutions of the two tyrosine residues in loop I, was ca. 10-fold more toxic than the wild-type Cry3A. The A2 mutant, with deletions of the two tyrosine residues in loop I, was ca. three-fold more toxic. Mutations at positions 353 and 354 had no significant effect on toxicity in *T. molitor*, except for a reduction in toxicity for the double mutant, N₃₅₃A,D₃₅₄A (Table 1).

Toxicity of the A1 and A2 mutant toxins compared to Cry3A in *C. scripta* and *L. decemlineata* revealed a trend toward lower LC₅₀s and steeper slopes for the mutants when compared to Cry3A (Tables 2 and 3).

Table 2

Maximum likelihood estimates of median lethal concentrations (LC₅₀) of Cry3A, A1 and A2 mutant proteins for young second instar *C. scripta* and *L. decemlineata*

Protein ^a	<i>C. scripta</i>			<i>L. decemlineata</i>		
	LC ₅₀ ^b	FL ₉₅ ^b	Slope ± S.E.M.	LC ₅₀	FL ₉₅	Slope ± S.E.M.
Cry3A	1.70	0.85–2.97	1.65 ± 0.34	2.07	1.41–3.44	1.90 ± 0.27
A1 mutant	0.66	0.38–0.97	2.29 ± 0.47	1.08	0.64–1.63	2.35 ± 0.41
A2 mutant	0.98	0.63–1.42	2.14 ± 0.37	0.94	0.57–1.48	1.82 ± 0.26

^aEach protein assay was replicated three times. Each replicate assay included four to six protein concentrations and a control applied to the appropriate leaf disks (poplar or potato) with 10–20 insects per concentration.

^bLC₅₀ and FL₉₅ are expressed as ng/mm².




Fig. 2. Competition binding of ^{125}I -labeled Cry3A toxin to *C. scripta* (A) and *L. decemlineata* (B) BBMV as a function of the concentration of unlabeled loop I mutant proteins. Vesicles (100 $\mu\text{g}/\text{ml}$) were incubated with labeled wild-type toxin (2.0 nM) in the presence of increasing concentrations of unlabeled competitors. Binding is expressed as percentage of the amount bound upon incubation with labeled toxin alone. A: $\circ$, wild-type Cry3A toxin; $\bullet$, A1 mutant ($\text{R}_{345}\text{A}, \text{Y}_{350}\text{F}, \text{Y}_{351}\text{F}$); $\diamond$, A2 mutant ($\text{R}_{345}\text{A}, \Delta\text{Y}_{350}, \Delta\text{Y}_{351}$). B: $\circ$, wild-type Cry3A toxin; $\bullet$, A1 mutant ($\text{R}_{345}\text{A}, \text{Y}_{350}\text{F}, \text{Y}_{351}\text{F}$); $\diamond$, A2 mutant ($\text{R}_{345}\text{A}, \Delta\text{Y}_{350}, \Delta\text{Y}_{351}$). Results are obtained from the average of three experiments. Each point represents the mean of duplicate samples.

3.3. Effects of mutations on membrane receptor binding

Competitive binding of ^{125}I -labeled Cry3A in the presence of unlabeled wild-type Cry3A was reduced to 13% for *C. scripta* (Fig. 2A) and 27% for *L. decemlineata* (Fig. 2B). Homologous competition binding assays, using ^{125}I -labeled wild-type Cry3A toxin, determined that *C. scripta* had a >3-fold higher binding affinity (K_{com}) than *L. decemlineata* (Table 4). In contrast, the binding site concentration (B_{max}) for *L. decemlineata* was >3-fold higher than *C. scripta* (Table 4).

Binding affinities for both the mutant toxins (A1 and A2) on *C. scripta* BBMV were similar and slightly lower than that for the wild-type Cry3A, although not significantly different (Table 4). The binding site concentration of *C. scripta* BBMV was significantly greater for the A1 mutant than that for either the wild-type Cry3A or the A2 mutant (Table 4). The binding assays with *L. decemlineata* BBMV revealed both greater affinity and a lower concentration of binding sites for the mutant toxins compared to the wild-type Cry3A (Table 4).

3.4. Dissociation analysis of A1 mutant on *C. scripta* BBMV

Since the A1 mutant toxin had minimal effect on K_{com} and B_{max} , but substantially increased toxicity to *C. scripta* larvae, dissociation assays were performed to further examine the kinetic aspects of membrane receptor binding. We observed that about 44% of wild-type Cry3A could be dissociated from the BBMV, reaching equilibrium after 10–20 min (Fig. 3A). In contrast, only 24% of A1 mutant toxin was dissociated from its binding site in *C. scripta* by the addition of excess unlabeled ligand, reaching equilibrium after 30 min (Fig. 3A). The dissociation rate calculated for A1 mutant, $k_{\text{dis}} = 0.05 \text{ min}^{-1}$, was nearly five-fold lower when compared to the dissociation rate for the wild-type Cry3A, $k_{\text{dis}} = 0.24 \text{ min}^{-1}$ (Fig. 3B).

4. Discussion

Our results support that site-specific changes in loop I of the receptor binding domain of Cry3A δ -endotoxin altered the toxicity of this protein in coleopteran larvae. Of particular interest were two mutant toxins, A1 and A2, which were more toxic than the wild-type Cry3A in bioassays of three coleopteran species (Tables 1 and 2). These two mutants differed only slightly, in A1 phenylalanines were substituted for tyrosines at residues 350 and 351 ($\text{R}_{345}\text{A}, \text{Y}_{350}\text{F}, \text{Y}_{351}\text{F}$), and in A2 those tyrosines were deleted ($\text{R}_{345}\text{A}, \Delta\text{Y}_{350}, \Delta\text{Y}_{351}$). Furthermore, receptor binding studies demonstrated that phenylalanine substitutions for Y_{350} and Y_{351} mostly affect dissociation, apparently accounting for the enhancement of toxicity. Ultimately, we hope to understand the structural role of loop I residues in insect toxicity and parameters of binding to insect midgut receptors.

The similar toxicity of wild-type Cry3A in *C. scripta* and *L. decemlineata*, both within the same family of Chrysomelidae, was achieved through quite different receptor binding parameters. The wild-type Cry3A binding affinity (K_{com}) to *C. scripta* BBMV was greater than that for *L. decemlineata* BBMV, whereas the Cry3A binding site concentration (B_{max}) was higher for *L. decemlineata* than that for *C. scripta* (Table 4).

Table 4
Binding properties of Cry3A and mutant proteins incubated with BBMV of *C. scripta* and *L. decemlineata*

Protein	<i>C. scripta</i>		<i>L. decemlineata</i>	
	K_{com} (nM) ^a	$B_{\text{max}}^{\text{a,b}}$	K_{com} (nM) ^a	$B_{\text{max}}^{\text{a,b}}$
Cry3A	37.9 ± 4.2 a	23.3 ± 3.3 b	135.9 ± 11.8 a	79.3 ± 6.4 a
A1 mutant	49.7 ± 5.6 a	32.7 ± 2.1 a	54.4 ± 6.8 b	50.3 ± 2.9 c
A2 mutant	46.3 ± 5.1 a	25.6 ± 1.5 b	53.8 ± 6.5 b	63.2 ± 1.9 b

Values for K_{com} or B_{max} within one species and followed by the same letter are not significantly different using Student's *t*-test with paired comparisons.

^aThe binding constants (K_{com}) and binding site concentrations (B_{max}) represent the calculated mean $\pm$ S.E.M. of three competition binding experiments.

^bThe B_{max} values are expressed as pmol/mg of BBMV protein.

The mutant toxins, A1 and A2, also displayed different receptor binding parameters in the two chrysomelid species. In BBMV prepared from *L. decemlineata*, both mutant strains showed enhanced binding affinity and fewer binding sites (Table 4). The differences in toxicity between the two mutants were similar to differences in binding site concentration. In contrast, receptor binding parameters measured in *C. scripta* BBMV revealed slightly lower binding affinities for both mutants, and a significantly greater number of binding sites for the more toxic mutant A1 (Table 4).

Bioassays for the A2 mutant indicated that deletion of the two tyrosines in loop I showed greater toxicity in both *T. molitor* and *L. decemlineata* (2.7- and 2.2-fold, respectively), but only a minimal effect on toxicity in *C. scripta* (Tables 1 and 3). It is not clear how the adjacent N₃₅₃ and D₃₅₄ residues contribute to receptor binding and toxicity, but previous results indicated that alanine substitutions for N₃₅₃ and D₃₅₄ caused the loss of binding affinity and toxicity [17]. In addition, toxicity of the D₃₅₄E mutant was similar to that of the wild-type Cry3A (Table 1). Perhaps, the deletions of Y₃₅₀ and Y₃₅₁ allow the aliphatic side chains of N₃₅₃ and D₃₅₄ to make better contact with the receptor, enhancing toxicity.

The results of our mutational, bioassay and receptor binding studies suggest that hydrophobic side chains for the residues in loop I interact with the receptor. Hence, the fact that phenylalanine can effectively substitute for Y₃₅₀ and Y₃₅₁ in toxicity enhancement suggests that it is not the phenolic hydroxyl group of tyrosine at these positions, but rather its aromatic phenol ring that is important for toxicity (Tables 1 and 2). This observation implies that the hydrophobic interaction of aromatic residues with the contact interface on the receptor side may be necessary for tight binding affinity. The well-packed hydrophobic contact residues may play an important role in the insecticidal activity and binding affinity.

The involvement of hydrophobic interactions between receptors and ligands is well documented. Mutagenesis and crystallographic studies have revealed that aromatic amino acids, tyrosine, phenylalanine and tryptophan, were the major determinants of binding affinity in human lysozyme [44], diphtheria toxin [45], human growth hormone and the extracellular binding domain of its receptor [46,47], and other insecticidal Cry toxins [48]. In our study, the binding analysis of A1 and A2 mutants to *C. scripta* and *L. decemlineata* BBMV verified the correlation between hydrophobic interaction in binding affinity and insecticidal activity. In *L. decemlineata* larvae, competition binding analysis indicated that both the deletions and phenylalanine substitutions for Y₃₅₀ and Y₃₅₁ resulted in increased binding affinity, a ca. 2.5-fold lower K_{com} value than wild-type toxin (Fig. 2B and Table 4). These results suggest that the more hydrophobic phenylalanine rings or removal of the phenolic hydroxyl groups at residues 350 and 351 can lead to higher binding affinity and higher toxicity.

Interestingly, there was no significant difference in the competition analysis between wild-type, A1 and A2 mutant toxins to *C. scripta* (Fig. 2A and Table 4) or *T. molitor* (unpublished data), despite enhanced toxicity of the A1 mutant toxin in *T. molitor* and *C. scripta* larvae (ca. 11.4- and 2.6-fold, respectively). Our investigations of the dissociation of the A1 mutant toxin suggested that the hydrophobic phenylalanines substituted for Y₃₅₀ and Y₃₅₁ reduced the dissociation rate by nearly five-fold, as compared with wild-type Cry3A (Fig. 3B). This slower dissociation rate may account for its greatly




Fig. 3. Dissociation of ^{125}I -labeled Cry3A wild-type and A1 mutant toxins from *C. scripta* BBMV. A: 2.0 mM ^{125}I -labeled toxin (wild-type or A1 mutant) was incubated with 100 $\mu\text{g}/\text{ml}$ vesicles at room temperature for 60 min. After incubation, the mixtures were diluted two-fold in binding buffer containing 0.5 μM corresponding unlabeled toxins. Non-specific binding was subtracted. ●, wild-type Cry3A toxin; ■, A1 mutant (R₃₄₅A, Y₃₅₀F, Y₃₅₁F). Each point represents the mean $\pm$ S.D. of triplicate experiments. B: Single exponential fits for dissociation binding curves give dissociation rate constants, k_{dis} , for wild-type and mutant toxins. ●, wild-type Cry3A toxin; ■, A1 mutant (R₃₄₅A, Y₃₅₀F, Y₃₅₁F).

enhanced toxicity in *C. scripta*. This result may have direct relevance to the enhancement of insecticidal activity. Our results are consistent with the tight binding affinity of Cry1Ab toxin to *Manduca sexta* BBMV, also caused by the hydrophobic phenylalanine, F₃₇₁, in the surface-exposed loop [29].

Overall binding affinity can be considered a dynamic balance of association and dissociation reactions. Since the affinity constant (K_{com}) is determined by competition binding experiments, it reflects all binding events. Distinction of association and dissociation will be necessary to explain the subtle quantitative differences in the toxicity of δ -endotoxins in their hosts.

The implications of our findings are important for rational design of δ -endotoxins that overcome insect resistance to these toxins and for altered or improved insecticidal activity, as has been achieved in other Cry toxins [4,27]. For instance, the hydrophobic residues around the loop I region in domain II, F₃₁₃NYW₃₁₆ of Cry1Aa, E₃₁₃YYW₃₁₆ of Cry1Ab and Y₃₁₃YYW₃₁₆ of Cry1Ac toxins, may be identified as the homologue of Y₃₅₀YGND₃₅₄ of Cry3A by functional mapping and sequence alignment [49]. Identification of these hydrophobic contact residues provides the potential for constructing

genetically enhanced and durable toxins for use as biological pesticides.

Acknowledgements: The initial culture of *T. molitor* was generously provided by G. Keeney (Department of Entomology, The Ohio State University). We are grateful to Mr. David Matthews and Dr. Takashi Yamamoto for providing the oligonucleotides used for mutagenesis, Drs. Dan Zeigler, Francis Rajamohan and Mi K. Lee for comments on the manuscript and helpful discussions. This work was supported by a grant from the National Institute of Health, NIH RO1 A1 29092 to D.H.D., USDA Cooperative State Research Service 93-37-302-9360 to C.N.K., and USDA Forest Service NCFES Cooperative Agreement 23-92-06.

References

- [1] Tabashnik, B.E., Cushing, N.L., Finson, N. and Johnson, M.W. (1990) *J. Econ. Entomol.* 83, 1671.
- [2] Tabashnik, B.E. (1994) *Annu. Rev. Entomol.* 39, 47–79.
- [3] Bauer, L.S. (1995) *Florida Entomol.* 78, 414–443.
- [4] Rajamohan, F., Lee, M.K. and Dean, D.H. (1998) *Prog. Nucleic Acid Res. Mol. Biol.* 60, 1–27.
- [5] Schnepf, E., Crickmore, N., Van Rie, J., Lereclus, D., Baum, J., Feitelson, J., Zeigler, D.R. and Dean, D.H. (1998) *Microbiol. Mol. Biol. Rev.* 62, 775–806.
- [6] Höfte, H. and Whiteley, H.R. (1989) *Microbiol. Mol. Rev.* 53, 242–255.
- [7] Crickmore, N., Zeigler, D.R., Feitelson, J., Schnepf, E., Van Rie, J., Lereclus, D., Baum, J. and Dean, D.H. (1998) *Microbiol. Biol. Rev.* 62, 807–813.
- [8] Feitelson, J.S. (1993) in: *Advanced Engineered Pesticides* (Kim, L., Ed.), pp. 63–71, Marcel Dekker, New York.
- [9] Krieg, A., Huger, A., Langenbruch, G. and Schnetter, W.B.H. (1983) *J. Appl. Entomol.* 96, 500–508.
- [10] Krieg, V.A., Huger, A.M. and Schnetter, W. (1987) *J. Appl. Entomol.* 104, 417–424.
- [11] Donovan, W.P., González Jr., J.M., Gilbert, M.P. and Dankocsik, C. (1988) *Mol. Gen. Genet.* 214, 365–372.
- [12] Lambert, B., Höfte, H., Annys, K., Jansens, S., Soetaert, P. and Peferoen, M. (1992) *Appl. Environ. Microbiol.* 58, 2536–2542.
- [13] Krieg, V.A., Huger, A.M., Langenbruch, G.A. and Schnetter, W. (1984) *Anz. Schädlingskde. Pflanzenschutz Umweltschutz* 57, 145–150.
- [14] Bauer, L.S. (1990) *Environ. Entomol.* 19, 428–431.
- [15] Li, J., Carroll, J. and Ellar, D.J. (1991) *Nature* 353, 815–821.
- [16] Grochulski, P., Masson, L., Borisova, S., Pusztai-Carey, M., Schwartz, J.-L., Brousseau, R. and Cygler, M. (1995) *J. Mol. Biol.* 254, 447–464.
- [17] Wu, S.-J. and Dean, D.H. (1996) *J. Mol. Biol.* 255, 628–640.
- [18] Lee, M.K., Young, B.A. and Dean, D.H. (1995) *Biochem. Biophys. Res. Commun.* 216, 306–312.
- [19] de Maagd, R.A., van der Klei, H., Bakker, P.L., Stiekema, W.J. and Bosch, D. (1996) *Appl. Environ. Microbiol.* 62, 2753–2757.
- [20] Burton, S.L., Ellar, D.J., Li, J. and Derbyshire, D.J. (1999) *J. Mol. Biol.* 287, 1011–1022.
- [21] Lee, L.K., You, T.H., Gould, F.L. and Dean, D.H. (1999) *Appl. Environ. Microbiol.* 65, 4513–4520.
- [22] Chen, X.J., Lee, M.K. and Dean, D.H. (1993) *Proc. Natl. Acad. Sci. USA* 90, 9041–9045.
- [23] Wolfersberger, M.G., Chen, X.J. and Dean, D.H. (1996) *Appl. Environ. Microbiol.* 62, 279–282.
- [24] Schwartz, J.L., Potvin, L., Chen, X.J., Brousseau, R., Laprade, R. and Dean, D.H. (1997) *Appl. Environ. Microbiol.* 63, 3978–3984.
- [25] Lu, H., Rajamohan, F. and Dean, D.H. (1994) *J. Bacteriol.* 176, 5554–5559.
- [26] Rajamohan, F., Alcantara, E., Lee, M.K., Chen, X.J., Curtiss, A. and Dean, D.H. (1995) *J. Bacteriol.* 177, 2276–2282.
- [27] Rajamohan, F., Alzate, O., Cottrill, J.A., Curtiss, A. and Dean, D.H. (1996) *Proc. Natl. Acad. Sci. USA* 93, 14338–14343.
- [28] Rajamohan, F., Cottrill, J.A., Gould, F. and Dean, D.H. (1996) *J. Biol. Chem.* 271, 2390–2397.
- [29] Smith, G.P. and Ellar, D.J. (1994) *Biochem. J.* 302, 611–616.
- [30] Yanisch-Perron, C., Vieira, J. and Messing, J. (1985) *Gene* 33, 103–119.
- [31] Kunkel, T.A. (1985) *Proc. Natl. Acad. Sci. USA* 82, 488–492.
- [32] Sanger, R., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
- [33] Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, Cold-Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- [34] Ge, A.Z., Shivarova, N.I. and Dean, D.H. (1989) *Proc. Natl. Acad. Sci. USA* 86, 4037–4041.
- [35] Laemmli, U.K. (1970) *Nature* 227, 680–685.
- [36] Robertson, J.L. and Preisler, H.K. (1992) *Pesticide Bioassays with Arthropods*, pp. 17–34, CRC Press, Boca Raton, FL.
- [37] MacIntosh, S.C., Lidster, B.D. and Kirkham, C.L. (1994) *J. Invertebr. Pathol.* 63, 97–98.
- [38] English, L.H. and Readdy, T.L. (1989) *Insect Biochem.* 19, 145–152.
- [39] Wolfersberger, M.G. (1990) *Experientia* 46, 475–477.
- [40] Lee, M.K., Milne, R.E., Ge, A.Z. and Dean, D.H. (1992) *J. Biol. Chem.* 267, 3115–3121.
- [41] Munson, P.J. and Rodbard, D. (1980) *Anal. Biochem.* 107, 220–239.
- [42] Ihara, H., Kuroda, E., Wadano, A. and Himeno, M. (1993) *Bio-sci. Biotechnol. Biochem.* 57, 200–204.
- [43] Johnson, K.A. (1992) *Enzymes* 20, 1–61.
- [44] Muraki, M., Harata, K. and Jigami, Y. (1992) *Biochemistry* 31, 9212–9219.
- [45] Blanke, S.R., Huang, K. and Collier, R.J. (1994) *Biochemistry* 33, 15494–15500.
- [46] Cunningham, B.C. and Wells, J.A. (1993) *J. Mol. Biol.* 234, 554–563.
- [47] Clackson, T. and Wells, J.A. (1995) *Science* 267, 383–386.
- [48] Rajamohan, F., Hussain, S.-R.A., Cottrill, J.A., Gould, F. and Dean, D.H. (1996) *J. Biol. Chem.* 271, 25220–25226.
- [49] Kwak, I.-S., Lu, H. and Dean, D.H. (1995) *Mem. Inst. Oswaldo Cruz* 90, 75–79.