

TRIAZOLO[4,5-d]PYRIMIDINES. VII. THE PHOTOCHEMICAL TRANSFORMATION OF 3-PHENYL-3H-1,2,3-TRIAZOLO[4,5-d]PYRIMIDINES INTO 9H-PYRIMIDO[4,5-b]INDOLES

Takeo Higashino* and Eigaku Hayashi

Shizuoka College of Pharmacy, 2-2-1 Oshika, Shizuoka-shi,
422 Japan

Hideaki Matsuda and Tatsuhiko Katori

Central Research Laboratory, S S Pharmaceutical Co., Ltd.,
1143 Nanpeidai, Narita-shi, 286 Japan

Abstract — The photolysis of 3-phenyl-3H-1,2,3-triazolo-[4,5-d]pyrimidines in dioxane, benzene and methanol gave 9H-pyrimido[4,5-b]indoles in moderate yields. Similarly, 3-phenyl-3H-1,2,3-triazolo[4,5-b]pyridines were easily transformed by the photolysis into 9H-pyrido[2,3-b]indoles.

It was reported by Ohashi and his co-workers that on irradiation, 5-chloro-1-(4-chlorophenyl)-1H-1,2,3-benzotriazole (A) lost a nitrogen molecule to give 3,6-dichloro-9H-carbazole (B) as a sole product.¹⁾






Chart 1

Anticipating that similar photochemical transformation would take place, we examined the photolysis of 3-phenyl-3H-1,2,3-triazolo[4,5-d]pyrimidines (1), and succeeded in obtaining 9H-pyrimido[4,5-b]indoles (2) in moderate yields.


On irradiation with a high pressure mercury lamp (100 W) in a solvent such as dioxane, benzene, and methanol for 10 to 15 hr, 3-phenyl-3H-1,2,3-triazolo-[4,5-d]pyrimidine (1a)²⁾ and its derivatives (1b)³⁾ and 1c)⁴⁾ having electron




<u>1</u>	-R
<u>1a</u>	-H
<u>1b</u>	-Cl
<u>1c</u>	-CN
<u>1d</u>	-NHCH ₃
<u>1e</u>	-NHCH ₂ C ₆ H ₅
<u>1f</u>	-N(CH ₃) ₂
<u>1g</u>	-OCH ₃
<u>1h</u>	-OC ₂ H ₅
<u>1i</u>	-OC ₆ H ₅
<u>1j</u>	-CH ₃




<u>2</u>	-R ₁	-R ₂
<u>2a</u>	-H	-H
<u>2b</u>	-Cl	-H
<u>2c</u>	-CN	-H
<u>2d</u>	-NHCH ₃	-H
<u>2e</u>	-NHCH ₂ C ₆ H ₅	-H
<u>2f</u>	-N(CH ₃) ₂	-H
<u>2g</u>	-OCH ₃	-H
<u>2h</u>	-OC ₂ H ₅	-H
<u>2i</u>	-OC ₆ H ₅	-H
<u>2j</u>	-CH ₃	-H
<u>2k</u>	-	-H
<u>2l</u>	-CH ₂ OH	-H
<u>2m</u>	-CN	-OCH ₃




<u>5</u>	-R ₁	-R ₂
<u>5a</u>	-H	-H
<u>5b</u>	-CH ₃	-COOC ₂ H ₅
<u>5c</u>	-CH ₃	-COCH ₃
<u>5d</u>	-(CH ₂) ₄ -	
<u>5e</u>	-(CH ₂) ₃ -	
<u>5f</u>	-C ₆ H ₅	-H



<u>6</u>	-R ₁	-R ₂
<u>6a</u>	-H	-H
<u>6b</u>	-CH ₃	-COOC ₂ H ₅
<u>6c</u>	-CH ₃	-COCH ₃
<u>6d</u>	-(CH ₂) ₄ -	
<u>6e</u>	-(CH ₂) ₃ -	
<u>6f</u>	-C ₆ H ₅	-H



<u>3</u>	-R
<u>3a</u>	-H
<u>3b</u>	-Cl
<u>3c</u>	-CN



<u>4</u>	-R
<u>4b</u>	-Cl
<u>4c</u>	-CN
<u>4d</u>	-OCH ₃

Chart 2

attracting substituents such as chloro and cyano groups at the 7-position gave the corresponding ring transformation products, 9H-pyrimido[4,5-b]indoles (2a)⁷⁾ to 2c) as main products together with by-products (2k to 2m, 3a)⁵⁾ to 3c, and 4b)⁶⁾ to 4d). Thus, in the case of 1a in dioxane 2k as a by-product was formed by the photoreaction of the resulting 2a with dioxane used, introducing a substituent

Table I. The Photolysis of 1 and 5

Starting material	Solvent	Product					
		No.	Mp (°C)	Yield (%)	No.	Mp (°C)	Yield (%)
<u>1a</u>	Dioxane	<u>2a</u>	237-240	33	<u>2k</u>	248-249	9
<u>1a</u>	Benzene	<u>2a</u>		50	<u>3a</u>	113-114	11
<u>1a</u>	Methanol	<u>2a</u>		54	<u>2l</u>	250-252	15
<u>1b</u>	Dioxane	<u>2b</u>	274-277	14	<u>4b</u>	158-159	2
<u>1b</u>	Benzene	<u>2b</u>		32	<u>3b</u>	167-169	8
<u>1b</u>	Methanol	<u>2b</u>		7	<u>4d</u>	149-151	2
<u>1c</u>	Dioxane	<u>2c</u>	284-286	25	<u>4c</u>	150-152	12
<u>1c</u>	Benzene	<u>2c</u>		54	<u>3c</u>	218-220	19
<u>1c</u>	Methanol	<u>2c</u>		85	<u>2m</u>	260-261	4
<u>1d</u>	Dioxane	<u>2d</u>	249-250	69			
<u>1e</u>	Dioxane	<u>2e</u>	256-257	62			
<u>1f</u>	Dioxane	<u>2f</u>	238-239	91			
<u>1g</u>	Dioxane	<u>2g</u>	208-210	85			
<u>1h</u>	Dioxane	<u>2h</u>	190-191	55			
<u>1i</u>	Dioxane	<u>2i</u>	199-200	48			
<u>1j</u>	Dioxane	<u>2j</u>	260-262	23			
<u>5a</u>	Dioxane	<u>6a</u>	115-116	61			
<u>5b</u>	Dioxane	<u>6b</u>	227-228	38			
<u>5c</u>	Dioxane	<u>6c</u>	255-257	34			
<u>5d</u>	Dioxane	<u>6d</u>	254-255	48			
<u>5e</u>	Dioxane	<u>6e</u>	294-295	69			
<u>5f</u>	Dioxane	<u>6f</u>	245-246	52			

into the 4-position. Similarly, the reaction of 1a in methanol gave 4-hydroxy-methyl derivative of 2a (2l) as a by-product. The reaction of 1b and 1c in dioxane gave the corresponding ring fission products (4b and 4c) as by-products. In the case of 1b in methanol, 4d as a by-product was formed by the nucleophilic substitution of the resulting 4b with methanol. In the case of 1c in methanol, 2m as a by-product was formed by the photoreaction of the resulting 2c with methanol used, introducing a methoxy group into the 9-position. In the cases of 1a to 1c in benzene, 3a to 3c as by-products were formed by the photoreaction of the ring fission intermediates, like A', with benzene, introducing a phenyl group into the 5-position. How to depend on a solvent used for formation of by-products is not clear as yet.

The photolysis of triazolopyrimidines (1d to 1j)⁴⁾ having electron donating substituents such as amino, alkoxyl, and alkyl groups at the 7-position resulted in the formation of the corresponding 4-substituted pyrimidoindoles (2d to 2j) as sole products.

Similarly, 3-phenyl-3H-1,2,3-triazolo[4,5-b]pyridines (5a to 5f)^{8, 9)} were subjected to the photolysis in dioxane, the corresponding 9H-pyrido[2,3-b]indoles (6a¹⁰⁾ to 6f) were obtained in moderate yields.

The structures of the products, 2, 3, 4, and 6 were assigned on the basis of the spectral data as shown in Table II.

Table II. The Spectral Data for 2, 3, 4, and 6

Compd.	MS: M ⁺ m/e	IR $\nu_{\text{max}}^{\text{KBr}}$ cm ⁻¹	NMR (in CDCl ₃): ppm			
			C ² -H ^s	NH ^{bs}	C ₆ H ₅ ^m	Other
<u>2a</u>	169	3132(NH)	9.00	12.3	-	9.50 ^s (C ⁴ -H), 7.2-8.4 ^m (C ⁵⁻⁸ -H) a)
<u>2b</u>	203	3120(NH)	8.70	12.6	-	7.2-8.4 ^m (C ⁵⁻⁸ -H)
<u>2c</u>	194	3120(NH) 2230(CN)	8.79	-	-	7.2-8.4 ^m (C ⁵⁻⁸ -H) b)
<u>2d</u>	198	3450, 3110(NH)	8.45	11.9 7.1	-	3.20 ^d (N-CH ₃), 7.2-8.5 ^m (C ⁵⁻⁸ -H) a)
<u>2e</u>	274	3455, 3120(NH)	8.45	12.0	7.1-8.6	5.00 ^d (NHCH ₂), 7.1-8.6 ^m (C ⁵⁻⁸ -H) a)
<u>2f</u>	212	3180(NH)	8.35	-	-	3.42 ^s (N(CH ₃) ₂), 7.2-8.2 ^m (C ⁵⁻⁸ -H) b)
<u>2g</u>	199	3120(NH)	8.20	11.7	-	4.30 ^s (OCH ₃), 7.3-8.3 ^m (C ⁵⁻⁸ -H)
<u>2h</u>	213	3120(NH)	8.73	11.5	-	4.79 ^q , 1.62 ^t (OC ₂ H ₅), 7.3-8.3 ^m (C ⁵⁻⁸ -H)
<u>2i</u>	261	3120(NH)	8.53	12.2	7.69 ^s	7.2-8.4 ^m (C ⁵⁻⁸ -H) a)
<u>2j</u>	183	3100(NH)	8.80	-	-	3.05 ^s (CH ₃), 7.2-8.3 ^m (C ⁵⁻⁸ -H) b)
<u>2k</u>	255	3110(NH)	8.82	12.3	-	3.1-5.5 ^m (7H, CH), 7.2-8.4 ^m (C ⁵⁻⁸ -H)
<u>2l</u>	199	3130(NH or OH)	8.92	12.2	-	5.20 ^s (CH ₂ O), 5.2 ^{bs} (OH), 7.2-8.4 ^m (C ⁵⁻⁸ -H) a)
<u>2m</u>	224	2225(CN)	8.50	-	-	4.20 ^s (OCH ₃), 7.2-7.9 ^m (C ⁵⁻⁸ -H)
<u>3a</u>	247	3230(NH)	8.26	6.9	7.1-7.8	8.76 ^s (C ⁶ -H)
<u>3b</u>	281	3220(NH)	8.50	6.5	6.9-7.8	-
<u>3c</u>	272	3275(NH) 2230(CN)	8.70	6.7	7.1-7.7	-
<u>4b</u>	205	3230(NH)	8.41	8.2	7.35 ^s	6.71 ^s (C ⁵ -H)
<u>4c</u>	196	3230(NH) 2240(CN)	8.61	9.7	7.0-7.8	7.12 ^s (C ⁵ -H)
<u>4d</u>	201	3210(NH)	8.50	8.7	7.2-7.8	6.24 ^s (C ⁵ -H), 4.00 ^s (OCH ₃)
<u>6a</u>	168	3130(NH)	-	12.0	-	7.1-8.7 ^m (7H, C ²⁻⁸ -H)
<u>6b</u>	254	3130(NH) 1710(CO)	-	11.5	-	8.90 ^s (C ⁴ -H), 7.2-8.2 ^m (C ⁵⁻⁸ -H), 4.50 ^q , 1.45 ^t (OC ₂ H ₅), 3.05 ^s (CH ₃)
<u>6c</u>	224	3120(NH) 1670(CO)	-	-	-	8.72 ^s (C ⁴ -H), 7.2-8.2 ^m (C ⁵⁻⁸ -H), 2.89 ^s (CH ₃), 2.73 ^s (COCH ₃) b)
<u>6d</u> ^{c)}	222	3130(NH)	-	-	-	8.00 ^s (C ⁴ -H), 7.0-8.1 ^m (C ⁵⁻⁸ -H),

$\underline{6e}^c)$	208	3130 (NH)	-	11.4	-	1.6-3.2 ^m ((CH ₂) ₄) ^{b)} 8.17 ^s (C ⁴ -H), 7.0-8.3 ^m (C ⁵⁻⁸ -H), 2.0-3.4 ^m ((CH ₂) ₃) ^{a)}
$\underline{6f}$	244	3130 (NH)	-	11.7	-	7.0-8.5 ^m (11H, C ³⁻⁸ -H, C ₆ H ₅)

a) In CDCl₃-(CD₃)₂SO.

b) IN CDCl₃-CD₃OD.

c) The numbering used is that of 9H-pyrido[2,3-b]indole ring system.

ACKNOWLEDGEMENT — A part of this work was supported by Grant-in-Aid for Special Project Research in Development and Utilization of Nitrogen-Organic Resources from the Ministry of Education, Science and Culture.

REFERENCES

- 1) M. Ohashi, K. Tsujimoto, and T. Yonezawa, Chem. Commun., 1970, 1089.
- 2) T. Higashino, T. Katori, and E. Hayashi, Chem. Pharm. Bull. (Tokyo), 27, 2431 (1979).
- 3) D. J. Brown and M. N. Paddon-Row, J. Chem. Soc. (C), 1967, 1856.
- 4) T. Higashino, T. Katori, H. Kawayaya, and E. Hayashi, Chem. Pharm. Bull. (Tokyo), 28, 337 (1980).
- 5) G. Tsatsaronis and F. Effenberger, Chem. Ber., 94, 2876(1961).
- 6) H. C. Carrington, F. H. S. Curd, and D. N. Richardson, J. Chem. Soc., 1955, 1858.
- 7) R. G. Glushkov, V. A. Volokova, and O. Yu. Magidson, Khim. Farm. Zh., 1, 25 (1967) (Russ). [Chem. Abstr., 68, 105143f (1968)].
- 8) R. L. Clark, A. A. Pessolano, and T.Y. Shen, U.S. Patent 4,000,286 (1976). [Chem. Abstr., 86, P189951^a (1977)].
- 9) T. Higashino, T. Katori, and E. Hayashi, Chem. Pharm. Bull. (Tokyo), 27, 2861 (1979).
- 10) V. M. Clark, A. Cox, and E. J. Herbert, J. Chem. Soc. (C), 1968, 831.

Received, 16th July, 1980