

Gandharamine: A New Benzylisoquinoline Alkaloid from Berberis baluchistanicaMusa H. Abu Zarga,<sup>1</sup> Ghulam A. Miana<sup>2</sup> and Maurice Shamma\*,Department of Chemistry, The Pennsylvania State University,  
University Park, Pennsylvania 16802


*Gandharamine* (**1**) is a new quaternary oxobenzylisoquinolinium alkaloid found in *Berberis baluchistanica* Ahrendt (Berberidaceae), as well as in *Thalictrum fendleri* Engelm. ex Gray (Ranunculaceae). Its synthesis from dihydrobenzylisoquinoline **4** has been achieved. The only known naturally occurring close analog of *gandharamine* is *N*-methylpapaveraldine (?).

*Berberis baluchistanica* Ahrendt (Berberidaceae), native to Pakistan, has proven a rich repository of novel tertiary isoquinoline alkaloids including the proaporphine-benzylisoquinoline pakistanamine,<sup>3</sup> the aporphine-benzylisoquinoline pakistanine,<sup>3,4</sup> and the isoquinolone-benzylisoquinoline baluchistanamine.<sup>5</sup> Presently, investigation of the polar quaternary alkaloid fraction from 15 kg of dried plant material has yielded 17 mg of the new amorphous and optically inactive oxobenzylisoquinoline alkaloid *gandharamine* (**1**) chloride,  $C_{19}H_{18}O_4NCl$ ,  $\lambda_{max}^{EtOH}$  255, 316 and 365 nm (log  $\epsilon$  4.07, 3.65 and 3.73),  $\lambda_{min}^{EtOH}$  278 and 335 nm (log  $\epsilon$  3.43 and 3.62),  $\lambda_{max}^{EtOH, H^+}$  253 and 313 nm (log  $\epsilon$  3.98 and 3.72),  $\lambda_{min}^{EtOH, H^+}$  273 nm (log  $\epsilon$  3.96). The alkaloid is phenolic since the uv spectrum in base shows a bathochromic shift,  $\lambda_{max}^{EtOH, OH^-}$  253, 323sh and 365 nm (log  $\epsilon$  4.09, 3.63 and 3.89),  $\lambda_{min}^{EtOH, OH^-}$  278 nm (log  $\epsilon$  3.43).


The nmr spectrum in TFA-d includes three singlets at  $\delta$  3.96, 4.27 and 4.32, which can be assigned to one *N*-methylimino and two aromatic methoxyl groups, respectively. H-3 and H-4 appear as a doublet of doublets at  $\delta$  8.29 and 8.34 ( $J = 6.8$  Hz), while the remaining six aromatic protons are represented by a broad singlet centered at  $\delta$  7.21 and a sharp singlet at 7.63. The mass spectrum exhibits peaks  $m/e$  324 ( $M^+$ ,  $C_{19}H_{18}NO_4$ ) (1.6), 323 ( $M - 1$ )<sup>+</sup> (7.4), and 44 (100).

Sodium borohydride in methanol reduction of *gandharamine* chloride at room temperature overnight afforded the amorphous diastereomeric alcohols **2a** and **2b**,  $C_{19}H_{23}O_4N$ , in a 1:3 ratio. The mass spectra of both alcohols show an ( $M - H_2O$ )<sup>+</sup> peak  $m/e$  311, and a base peak  $m/e$  206. The respective nmr spectra in  $CDCl_3$  have been summarized in expressions **2a** and **2b**.


Acetylation of the major alcohol **2b** with acetic anhydride in pyridine overnight furnished diacetate **3**,  $C_{23}H_{27}O_6N$ . The  $CDCl_3$  nmr downfield shift of the ring C protons of **3** relative to




1




2a




2b




3




4



5



6



7

those of alcohol 2b locates the phenolic group of gandharamine (1) at C-4' rather than in ring A.

The structural assignment for gandharamine was confirmed by its synthesis from the known dihydrobenzylisoquinoline 4.<sup>6</sup> Oxidation of an ethanolic solution of this imine by bubbling a stream of air for three days<sup>7</sup> provided a 24% yield of the imino ketone 5,  $C_{18}H_{17}O_4N$ , mp 142-144° C ( $CH_2Cl_2$ ),  $\nu_{max}^{CHCl_3}$  1600, 1660 and 3450  $cm^{-1}$ , which was dehydrogenated using Pd/C in refluxing xylene overnight to furnish amorphous benzylisoquinoline 6 (66%),  $C_{18}H_{15}O_4N$ ,  $\nu_{max}^{CHCl_3}$  1660 and 1690  $cm^{-1}$ .<sup>8</sup> N-Methylation of 6 with methyl iodide in refluxing acetonitrile overnight afforded yellow needles of gandharamine (1) iodide, 75%,  $C_{19}H_{18}O_4NI$ , mp 185-186° C (methanol-ether).<sup>9</sup> Conversion to the amorphous chloride salt was achieved by passing a solution of the iodide in 1:1 methanol-acetone through an Amberlite CG-400 chloride anion exchange column. The chloride salt proved to be identical with gandharamine chloride obtained from the plant isolation sequence.

In some early isolation work in our laboratory carried out in the 1960's,<sup>10</sup> a quaternary alkaloidal chloride salt was obtained from Thalictrum fendleri Engelm. ex Gray (Ranunculaceae), native to Utah, which was not fully characterized at the time due to paucity of material. We have now found that the T. fendleri alkaloid is identical in terms of nmr and uv spectra, as well as tlc  $R_f$  values, with gandharamine chloride.

It is interesting to note that gandharamine (1) is the first known naturally occurring quaternary oxobenzylisoquinolinium salt belonging to the coclaurine series. Its sole counterpart in the reticuline series is N-methylpapaveraldine (7) found in Stephania sasakii Hayata (Menispermaceae).<sup>11</sup>

Acknowledgment: This research was supported by grant CA-11450 from the National Cancer Institute, National Institutes of Health, USPHS, DHHS.

#### References and Footnotes

1. Permanent address: Department of Chemistry, University of Jordan, Amman, Jordan.
2. Permanent address: Department of Chemistry, Gomal University, Dera Ismail Khan, Pakistan.
3. M. Shamma, J.L. Moniot, S.Y. Yao, G.A. Miana and M. Ikram, J. Am. Chem. Soc., **95**, 5742 (1973).
4. S.F. Hussain, M.T. Siddiqui and M. Shamma, Tetrahedron Lett., **21**, 4573 (1980).
5. M. Shamma, J.E. Foy and G.A. Miana, J. Am. Chem. Soc., **96**, 7809 (1974).
6. S. Teitel and A. Brossi, J. Heterocycl. Chem., **5**, 825 (1968); and J. Sam and A.J. Bej, J. Pharm. Sci., **56**, 906 (1967).
7. N.H. Martin, S.L. Champion and P.B. Belt, Tetrahedron Lett., 2613 (1980).
8. The  $\text{CDCl}_3$  nmr spectrum of benzylisoquinoline 6 exhibits  $\delta$  3.96 (s, 3H,  $\text{OCH}_3$ ), 4.07 (s, 3H,  $\text{OCH}_3$ ), 6.73 (d, 2H,  $J = 8.5$  Hz, H-3' and 5'), 7.16 and 7.52 (2s, 2H, H-5 and 8), 7.73 (d, 2H,  $J = 8.5$  Hz, H-2' and 6'), 7.73 (d, 1H,  $J = 4.3$  Hz, H-4), and 8.45 (d, 1H,  $J = 4.3$  Hz, H-3).
9. Gandharamine iodide shows ms  $m/e$  323 ( $M - 1$ )<sup>+</sup> (3.5), and 142 (base);  $\nu_{\text{max}}^{\text{KBr}}$  1590, 1645, 3070 and 3380  $\text{cm}^{-1}$ ; nmr ( $\text{DMSO}-d_6$ )  $\delta$  3.71 (s, 3H,  $\text{NCH}_3$ ), 4.07 (s, 3H, C-7  $\text{OCH}_3$ ), 4.14 (s, 3H, C-6  $\text{OCH}_3$ ), 6.87 and 7.87 (2s, 2H, H-5 and 8), 6.94 (d, 2H,  $J = 8.6$  Hz, H-3' and 5'), 7.80 (br d, 2H, H-2' and 6'), 8.43 (d, 2H,  $J = 6.9$  Hz, H-4) and 8.68 (d, 2H,  $J = 6.9$  Hz, H-3).
10. B.S. Dudock, Ph.D. thesis, The Pennsylvania State University (1966).
11. J. Kunitomo, E. Yuge and Y. Nagai, J. Pharm. Soc. Japan, **86**, 456 (1966).
12. NMR spectra are at 200 MHz (FT) with TMS as internal standard. All tlc was on Merck silica gel F-254 glass plates. TLC  $R_f$  values are as follows: Gandharamine (1) chloride 0.48 (chloroform-methanol- $\text{HCl}$  (50:50:trace), gandharamine iodide 0.28 (*n*-butyl alcohol-acetic acid-water 60:10:27), minor alcohol 2a 0.58 (diethylamine-chloroform 10:90), major alcohol 2b 0.52 (diethylamine-chloroform 10:90), diacetate 3 0.40 (methanol-chloroform 2:98).

Received, 9th September, 1981