

STUDIES ON 4-QUINAZOLINONES. PART IX¹.

RANEY NICKEL DESULPHURISATION OF 4-THIOQUINAZOLINES:

ALUMINA CATALYSED HYDRATION OF 1,4- AND 3,4-DIHYDROQUINAZOLINES

Manas Chakrabarty, Ajit K. Chakravarty and Satyesh C. Pakrashi*


Indian Institute of Chemical Biology, Calcutta-700 032, India

Abstract: A number of 2-substituted 3,4-dihydroquinazolines and 1,2-disubstituted 1,4-dihydroquinazolines but not the 3-mono or 2,3-disubstituted derivatives, prepared by Raney nickel desulphurisation of the corresponding 4-thioquinazolines (IV), underwent unusual alumina catalysed hydration across the C=N bond followed by ring opening through ring-chain tautomerism leading to substituted acetamide derivatives (III). A plausible mechanism has been put forward.


We earlier reported a few unusual and unpredictable course of reactions of 4-quinazolinones with some oxidising¹ and reducing²⁻⁶ agents influenced by the substitution pattern and the lone pair of nitrogen. We observed, *inter alia*, that when the C=N of N-alkyl substituted 4-quinazolinones could be smoothly reduced, attempted metal hydride reduction of the otherwise resistant amide carbonyl under forcing condition resulted in the ring opening by cleavage at the C-N bond⁴. We now wish to record yet another type of ring opening of the 3,4-dihydroquinazolines obtained by thiolation of the amide carbonyl followed by Raney nickel desulphurisation.

We recently observed that the product of LAH reduction of 2-diphenylmethyl-4-quinazolinone (Ia) varies with the column material used. Thus, while chromatography of the crude product over silica gel afforded⁵ a number of quinazoline and 3,4-dihydroquinazoline derivatives including a dimer, that over alumina furnished, besides two common products, *viz.* 2-(1'-hydroxydiphenylmethyl)-3,4-dihydroquinazoline (IIa with 1'-OH) and 2-diphenylmethylquinazoline (Va), a new compound, C₂₁H₂₀N₂O (M⁺, m/z 316.1583), m.p. 165°C, in 19% yield. The base peak at m/z 106 (C₇H₈N, species a), and the prominent peaks at m/z 167 (C₁₃H₁₁, species b) and 149 (C₈H₉N₂O, species c) coupled with a strong IR absorption band at 1655 cm⁻¹ (amide C=O) in CHCl₃ led us to N-(p-amino-benzyl)-diphenylacetamide (IIa) structure for the compound, corroborated by ¹H NMR spectrum (vide Exptl.).


Conceivably, the ring cleaved amide (IIa) arises from the 3,4-dihydroquinazoline (IIIa). In order to ascertain the validity of this assumption as also the role of alumina surface, we required some 3,4- and 1,4-dihydroquinazolines. However, in view of the difficulty in reducing the amide function of N-alkylated 4-quinazolinones⁴ and the reported^{7,8} susceptibility of the dihydroquinazolines to aëro-oxidation, we decided to chromatograph directly the crude products of Raney nickel desulphurisation⁹ of some dihydro-4-thioquinazolines (IVa - IVg), derived^{10,11} from the corresponding 4-oxo analogues (Ia - Ig).



a, m/z 106



b, m/z 167



c, m/z 149

The results shown in the Table revealed that irrespective of the column material, 1,4-dihydroquinazolines (III_f and III_g) could not be isolated. Nevertheless, silica gel column allowed the separation of all the 3,4-dihydroquinazolines (III_a - III_e) though only the N₃-substituted thiones furnished the corresponding dihydroquinazolines (III_d and III_e) over alumina. Again, the open-chain amides (II) were obtained from the 2-alkyl/aryl substituted thiones (IV) irrespective of alkyl substitution at N-1 only when alumina was used. The formation of other products, however, depended on the N-substitution rather than the adsorbent.


That I, II and V are derived indeed through the intermediacy of dihydroquinazolines (III) initially formed could be more clearly demonstrated as in the sequel. When pure III_b and III_d were separately kept adsorbed on alumina for 36 hr and eluted thereafter, the former gave the amide (II_b) and quinazoline derivative (V_b) in respective yields of 20 and 60%, and the latter furnished 4-quinazolinone derivative (Id) in 75% yield as the sole isolable product.

Since the formation of open-chain amides (II) was not observed with the N₃-substituted dihydroquinazolines in which the double bond is located between N₁ and C₂, it appeared likely that a double bond between C₂ and N₃ is a pre-requisite for this transformation. Presumably, the $\Delta^{2,3}$ -tautomer is involved in case of the 2-substituted compounds. Furthermore, the ring opening requires

Table. Products of desulphurised 4-thioquinazolines on chromatography.


4-Thio-quinazolines	Adsorbent*	Dihydro-quinazolines(%)	Amides (%)	Quinazolines (%)	4-Quinazolinones (%)
IVa	S	IIIa(11)	-	Va(26)	-
	A	-	IIa(7.5)	(31)	-
IVb	S	IIIb(10)	-	Vb(44)	-
	A	-	IIb(7.7)	(29)	-
IVc	S	IIIc(Unstable)**	-	Vc**	-
	A	-	IIc(7.3)	(33)	-
IVd	S	IIId(29)	-	-	Id(21)
	A	(10)	-	-	(22)
IVe	S	IIIe(6)	-	-	Ie(30)
	A	(28)	-	-	(22)
IVf	S	-	-	-	If(50)
	A	-	IIf(7.4)	-	(21)
IVg	S	-	-	-	Ig(48)
	A	-	IIg(7.7)	-	(20)

* S = silica gel, A = alumina ** Even during crystallisation, IIIc was partially converted to Vc. As such the yields could not be accurately estimated.




I a-e, X = O

IV a-e, X = S




II f, g, X = O


IV f, g, X = S




III a-c, f, g




V



VI a-e



VII f, g



VIII a-c

a R = CHPh₂, R' = H; b R = CH₂Ph, R' = H; c R = CH₃, R' = H; d R = CH₂Ph, R' = CH₃
e R = H, R' = Ph; f R = R' = CH₃; g R = CH₂Ph, R' = CH₃

a ring-chain tautomerism of an intermediate carbinolamine that necessitates hydration at the C₂-N₃ double bond (Scheme). As the open-chain amide was not obtained by chromatography through silica gel or by acid/base treatment of the dihydro derivative, obviously the hydration must have been induced by alumina which is known¹²⁻¹⁵ to cause a variety of surface catalysed secondary reactions.

The conversion of dihydroquinazolines to quinazolines evidently requires an available hydrogen at N₃ for elimination since quinazolinones were obtained instead when either of the nitrogen atoms was substituted.

The oxidation of the reactive methylene group at C-4 conceivably involves a free


IIa: m.p. 165°C; ν 1615 (1655 in CHCl_3); ^1H nmr 4.27 (2H, d, $\bar{J}$ = 6Hz, ArCH_2NHCO), 4.57 (2H, s, ArNH_2), 4.95 (1H, s, CHPh_2), 6.3-7.4 (14H, m, ArH); MS m/z 316 (M^+), 167, 165, 152, 149, 121, 106 (100%), 77.
 IIb: m.p. 119°C; ν 1623; ^1H nmr 3.53 (2H, s, COCH_2Ph), 4.25 (2H, br, ArNH_2).

products are given below:

Alumina (according to Brockmann; S. Merck, India) and silica gel (60-120 mesh; B. D. H., India) were used for chromatography. The ν (cm^{-1}) and ^1H nmr (δ ppm from TMS) spectra were taken in nujol and CDCl_3 respectively. General procedure for desubstituted: An alcoholic solution of the 4-thioquinazoline was treated with excess Raney nickel (activity W2), heated under reflux till the indicated disappearance of the starting material. After usual work-up, the crude mixture was chromatographed. The physical data of the

EXPERIMENTAL

appear to be peculiar to this system. on alumina would trigger radical reaction though the facile aero-oxidation^{7,8} It is, therefore, unlikely that mere adsorption of the 3,4-dihydroquinazolines dimer on being refluxed with deactivated Raney nickel in alcoholic solution. products, viz. quinazoline (Vb) or the dihydroquinazoline (IIb) yielded the zoline (IVb) over silica gel supports this contention. None of the other chromatography of desubstituted product of 2-benzyl-3,4-dihydro-4-thioquina-



radical mechanism¹⁶ plausibly through a hydroperoxide intermediate^{3,5}. The isolation of a hitherto unknown dimer, m.p. 180°C, characterised as 2,2'-dibenzyl-4,4'-biquinazolinyl (VI) by spectral analysis (vide Experimental), on

4.32 (2H, d, $J = 6\text{Hz}$, ArCH_2NHCO), 5.75 (1H, br, NHCO), 6.50-7.30 (9H, m, ArH); MS m/z 240 (M^+), 149, 121, 106 (100%), 91, 77.

IIc: m.p. 114°C (Lit.¹⁸ m.p. 113.5°C); ir 1658, 1641, 1615; ^1H nmr 2.00 (3H, s, COCH_3), 4.30 (2H, br, ArNH_2), 4.38 (2H, d, $J = 6\text{Hz}$, ArCH_2NHCO), 5.82 (1H, br, NHCO), 6.66-7.34 (4H, m, ArH); MS m/z 164 (M^+), 149, 121 (100%), 106, 76, 43.

IIf: m.p. 101°C ; ^1H nmr 2.01 (3H, s, COCH_3), 2.79 (3H, s, NHCH_3), 4.39 (2H, d, $J = 6\text{Hz}$, ArCH_2NHCO), 5.73 (1H, br, NHCO), 6.45-7.35 (4H, m, ArH); MS m/z 178 (M^+ , 100%), 135, 120, 119, 118, 91.

IIg: m.p. $91-93^\circ\text{C}$; ir 1653; ^1H nmr 2.78 (3H, s, NHCH_3), 3.56 (2H, s, COCH_2Ph), 4.33 (2H, d, $J = 6\text{Hz}$, ArCH_2NHCO), 4.90 (1H, br, ArNH), 5.65 (1H, br, NHCO), 6.00-7.30 (9H, m, ArH); MS m/z 254 (M^+), 163, 135, 120, 119, 118 (100%), 91, 77.

IIla: m.p. 171°C ; ir 1650, 1618; ^1H nmr 4.61 (2H, s, ArCH_2N), 5.13 (1H, s, CHPh_2), 6.66-7.50 (14H, m, ArH); MS m/z 298 (M^+), 297 (100%), 219, 193, 167, 165, 152, 131, 105, 104, 77.

IIb: m.p. 213°C (dec.); ir (KCl) 1660, 1640, 1630; ^1H nmr 4.05 (2H, s, ArCH_2Ph), 4.65 (2H, s, ArCH_2NH), 6.80-8.00 (9H, m, ArH); MS m/z 222 (M^+), 221 (100%), 194, 143, 131, 105, 104, 91, 77.

IIId: m.p. $245-247^\circ\text{C}$ (dec.); ir 1658, 1638; ^1H nmr 3.14 (3H, s, NCH_3), 4.45 (2H, s, ArCH_2Ph), 4.73 (2H, s, ArCH_2N), 6.80-7.80 (9H, m, ArH); MS m/z 236 (M^+), 235, 220, 219 (100%), 91, 77.

IIe: m.p. 122°C (Lit.¹⁹ m.p. 119°C); ir 1617; ^1H nmr 4.96 (2H, s, ArCH_2N), 8.16 (1H, s, $\text{C}_2\text{-H}$), 6.92-8.48 (9H, m, ArH); MS m/z 208 (M^+), 207, 130, 77.

Vb: m.p. 70°C (Lit.²⁰ m.p. $59-60^\circ\text{C}$); ir 1620; ^1H nmr 4.43 (2H, s, ArCH_2Ph), 7.20-8.00 (9H, m, ArH), 9.29 (1H, s, $\text{C}_4\text{-H}$); MS m/z 220 (M^+ , 100%), 219, 193, 166, 130, 117, 103, 91, 77, 76.

Vc: Oil (Lit.²¹ b.p. $255^\circ\text{C}/\text{atm}$); ir 1619; ^1H nmr 2.95 (3H, s, ArCH_3), 7.54-8.18 (4H, m, ArH), 9.38 (1H, s, $\text{C}_4\text{-H}$); MS m/z 144 (M^+ , 100%), 143, 129, 117, 103, 102, 76.

VI: m.p. 180°C ; ir 1616; ^1H nmr 4.51 (4H, s, $2 \times \text{ArCH}_2\text{Ph}$), 7.20-8.20 (18H, m, ArH); MS m/z 438 (M^+ , 100%), 437, 256, 219, 218, 91.

ACKNOWLEDGEMENT: One of the authors (M.C.) is grateful to the CSIR, New Delhi, for the grant of a Pool Officership and a Research Associateship later.

REFERENCES AND NOTES

1. Part VIII: S. C. Pakrashi, S. Chattopadhyay and A. K. Chakravarty, J. Org. Chem., 1976, 41, 2108.
2. S. C. Pakrashi and A. K. Chakravarty, Chem. Commun., 1969, 1443.
3. S. C. Pakrashi, J. Bhattacharyya and A. K. Chakravarty, Indian J. Chem., 1971, 9, 1220.
4. S. C. Pakrashi and A. K. Chakravarty, J. Org. Chem., 1972, 37, 3143.
5. S. C. Pakrashi and A. K. Chakravarty, Indian J. Chem., 1973, 11, 122.
6. S. C. Pakrashi and A. K. Chakravarty, J. Org. Chem., 1974, 39, 3828.
7. W. L. F. Armarego, 'Fused Pyrimidines, Pt. I' in 'The Chemistry of Heterocyclic Compounds', Ed. A. Weissberger, Interscience, New York, 1976, p. 405.
8. M. Lora-Tamayo, R. Madronero and G. G. Munoz, Chem. Ber., 1961, 94, 208.
9. B. C. Lawes and H. C. Scarborough, U. S. Patent 3, 127, 401 (Cl. 260-251), March 31, 1964; Chem. Abstr., 1964, 60, 14525.
10. D. J. Fry, J. D. Kendall and A. J. Morgan, J. Chem. Soc., 1960, 5062.
11. C. S. Rao, M. P. Dave, P. N. Mody and A. D. Pandya, Indian J. Chem., 1976, 14B, 999.
12. E. Lederer and M. Lederer, 'Chromatography' 2nd Ed., Elsevier, Amsterdam, 1957, p. 61 and Refs. cited therein.
13. G. H. Posner, Angew. Chem. Internat. Edn. Engl., 1978, 17, 487.
14. A. McKillop and D. W. Young, Synthesis, 1979, 401, 481.
15. S. C. Pakrashi, A. De and S. Chattopadhyay, Indian J. Chem., 1968, 6, 472.
16. G. R. Pettit and E. E. van Tamelen, 'Organic Reactions', Ed. A. C. Cope, John Wiley and Sons, New York, 1962, 12, p. 356.
17. H. O. House, 'Modern Synthetic Reactions', 2nd Ed., W. A. Benjamin, California, 1972, p. 35.
18. 'Beilsteins Handbuch der Organischen Chemie', 4th Ed., Julius-Springer, Berlin, 1930, 13, p. 169.
19. E. C. Wagner and A. Eisner, J. Amer. Chem. Soc., 1937, 59, 879.
20. Y. Mizuno, K. Adachi and K. Ikeda, Pharm. Bull. (Japan), 1954, 2, 225.
21. Ref. 7, p. 61.

Received, 18th August, 1982