


## SYNTHESIS OF 7,7-DIMETHYLAPORPHINE ALKALOIDS

Hassan A. Ammar, Paul L. Schiff, Jr. and David J. Slatkin\*


Department of Pharmacognosy, School of Pharmacy, University of Pittsburgh,  
Pittsburgh, Pennsylvania 15261 U.S.A.

**Abstract** - The synthesis of N,0,0-trimethyltetrahydromelosmine (1,2,3,9-tetramethoxy-7,7-dimethylaporphine) (3) is described. The amide derivative (9), prepared from the appropriate acid (8) and amine (6) by condensation, was cyclized by a Bischler-Napieralski reaction to afford the 3,4-dihydroisoquinoline derivative (10). The N-methyltetrahydrobenzylisoquinoline (12) from the reduction of 10 and subsequent N-methylation, was cyclized with thallium trifluoroacetate/trifluoroacetic acid and boron trifluoride etherate to the target compound (3).

Two novel 7,7-dimethyltetrahydroaporphine alkaloids, melosmine (1) and melosmidine (2), have been recently isolated from *Guatteria melosma*<sup>1</sup>. In that investigation, reduction and subsequent N-methylation of these alkaloids afforded novel 7,7-dimethylaporphine alkaloids. This paper presents the total synthesis of one of these products, N,0,0-trimethyltetrahydromelosmine (1,2,3,9-tetramethoxy-7,7-dimethylaporphine) (3).




2,3,4 Trimethoxy- $\beta$ -phenethylamine (6) was prepared from 2,3,4-trimethoxybenzaldehyde (4) (Aldrich) via 5 in 2 steps [i.  $CH_3NO_2/NaOH/5-10^\circ/EtOH$ ; ii.  $LAH/Et_2O/reflux$ ] as previously described<sup>2</sup> in 65% yield.  $\alpha$ -Methyl- $\alpha$ -(3-methoxyphenyl)propionic acid (8) was prepared from sodium 3-methoxyphenylacetate (Aldrich) (7) by treatment with 2 successive equivalents of lithium diisopropylamide (LDA) (prepared *in situ*: diisopropylamine/ $n-BuLi$ ) and MeI in 61% yield<sup>3</sup>. The amine (6) and the acid (8) were condensed using a Schotten-Baumann reaction<sup>4</sup> and the resulting product (9) cyclized using the conditions of the Bischler-Napieralski reaction ( $POCl_3$ ) in 57% yield<sup>5</sup>. The resulting imine (10) was reduced ( $NaBH_4$ ) to afford the amine (11) in 82% yield. This product was N-methylated ( $CH_3O/NaBH_4$ ) in 73% yield to give 12 which was subsequently cyclized using thallium trifluoroacetate/TFA/ $BF_3 \cdot Et_2O/-40^\circ C$  to afford 3 in 9% yield.<sup>6</sup> The product was identical (UV, IR, MS) to N,0,0-trimethyltetrahydromelosmine (3).



4 R = CHO


5 R = CH=CH-NO<sub>2</sub>

6 R = CH<sub>2</sub>-CH<sub>2</sub>-NH<sub>2</sub>



7 R = H

8 R = CH<sub>3</sub>



## EXPERIMENTAL

$\alpha$ -Methyl- $\alpha$ -(3-methoxyphenyl)propionic Acid (8). 3-Methoxyphenylacetic acid (Aldrich) (10 g) was added to a slurry consisting of NaH (3 g) in mineral oil and diisopropylamine (9 ml) in anhydrous THF (200 ml). After briefly heating to reflux, the suspension was cooled to 0°C and N-butyllithium (4 g; 1 equivalent) was injected. The mixture was warmed to 30°C to complete the metallation, cooled to 0°C and MeI (9 g; 1 equivalent) was added. The mixture was stirred (2 hr) at 30°C, cooled to 0°C and a second equivalent of N-butyllithium (4 g) was injected. Following warming to 30°C to complete the metallation and subsequent cooling to 0°C, a second equivalent of MeI (9 g) was added dropwise with stirring. After stirring (3 hr) at 30°C, H<sub>2</sub>O (250 ml) was added. The aqueous layer was collected, acidified with conc. HCl and extracted with Et<sub>2</sub>O. The ether layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford 8 (7.1 g) (61%) as colorless plates; mp 48-50°C (aqueous MeOH); ir  $\nu$  max 3300-2500 (br), 1710 cm<sup>-1</sup>; uv  $\lambda$  max (MeOH) 222 nm (log  $\epsilon$  3.05), 273 (2.39), 281 (2.31); <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  10.45 (1H, brs) (COOH), 7.35-6.7 (4H, m) (aromatic), 3.75 (3H, s) (C-3 methoxy), 1.52 (6H, s) (gem-dimethyl); ms m/z 194 (M<sup>+</sup>, 49%), 193(15), 166(8), 150(29), 149(100), 135(26), 133(10), 121(61), 115(10), 109(43), 108(14), 105(12), 91(41), 77(45), 65(21).

m-Methoxy- $\alpha$ -methyl-N-(2,3,4-trimethoxyphenethyl)hydratropamide<sup>7</sup> [N-2,3,4-Trimethoxyphenethyl]- $\alpha$ -methyl- $\alpha$ -(3'-methoxyphenyl)propionamide<sup>1</sup> (9). Thionyl chloride (5 ml) was added to 8 (3 g) in dry benzene (30 ml) and the mixture was heated (50°C) for 3 hours under anhydrous conditions. The acid chloride remaining after removal of volatile substances by evaporation of the reaction mixture, was dissolved in CHCl<sub>3</sub> (50 ml) and added dropwise to a well-stirred solution of 6 (3.3 g) and triethylamine (6 g) in chloroform (50 ml) at 0°C. The mixture was stirred for 2 hours at room temperature, washed successively with 100 ml portions of water, 2% HCl and 2% NaHCO<sub>3</sub>, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford 9 (5.1 g) (85%) as a colorless oil; ir  $\nu$  max (film) 3400, 1655 cm<sup>-1</sup>; uv  $\lambda$  max (MeOH) 238 nm (log  $\epsilon$  5.25), 280 (5.30); <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  7.16 (1H, d, J=8Hz) (C-6), 6.85 (1H, d, J=8Hz) (C-5), 6.82 (3H, m) (C-4', 5', 6'), 6.50 (1H, d, J=2Hz) (C-2'), 5.7 (1H, brs) (NH), 3.83 (6H, s), 3.79 (3H, s), 3.76 (3H, s) (4 x methoxy), 3.33 (2H, q, J=6Hz) (CH<sub>2</sub>NH-), 2.63 (2H, t, J=6Hz) (ArCH<sub>2</sub>-), 1.51 (6H, s) (gem-dimethyl); ms m/z 238(2%), 237(M<sup>+</sup>, 6), 195(16), 194(100), 179(23), 166(10), 149(20), 135(10), 121(16), 109(10), 91(18), 79(9), 65(5).

3,4-Dihydro-5,6,7-trimethoxy-1-(3'-methoxy- $\alpha$ , $\alpha$ -dimethylbenzyl)isoquinoline<sup>7</sup> (10). POCl<sub>3</sub> (5 ml) was added to 9 in dry benzene (50 ml) and refluxed for 1.5 hours. The brown residue left after evaporation of the mixture was triturated twice with petroleum ether, dissolved in Me<sub>2</sub>CO (50 ml) and diluted with 2% aqueous HCl (200 ml). The aqueous acidic solution was washed twice with ether, cooled, basified with dilute NH<sub>4</sub>OH and immediately extracted with Et<sub>2</sub>O. The ether layer was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford 10 (2.2 g) (57%) as colorless plates; mp 121-2°C (ether); ir  $\nu$  max (KBr) 2920, 2820, 1600, 1580, 1570, 1560 cm<sup>-1</sup>; uv  $\lambda$  max (MeOH) 225 nm (log  $\epsilon$  4.03), 272 (3.61), 310 (2.60); <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  7.30-6.65 (4H, m) (C-2', 4', 5', 6'), 6.28 (1H, s) (C-8), 3.83 (3H, s), 3.80 (3H, s), 3.75 (3H, s) (C-5, 6, 3' methoxys), 3.30 (3H, s) (C-7 methoxy), 2.59 (2H, t, J=7Hz) (ArCH<sub>2</sub>-), 1.58 (6H, s) (gem-dimethyl); ms m/z 369(M<sup>+</sup>, 53%), 368(100), 354(27), 338(10), 206(10), 195(10), 194(60), 185(10), 169(13), 162(20), 156(10), 149(30), 136(32), 121(40), 109(26), 91(41), 77(29), 65(13).

1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1-(3'-methoxy- $\alpha,\alpha$ -dimethylbenzyl)isoquinoline<sup>7</sup> (11).

NaBH<sub>4</sub> (4g) and 10 (2g) in EtOH (50 ml) were stirred at room temperature for 2 hours. After evaporation, the residue was dissolved in H<sub>2</sub>O and extracted with CHCl<sub>3</sub>. The CHCl<sub>3</sub> layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford 11 (1.7 g) (82%) as an oil; mp of N-acetate 135-6°C (CHCl<sub>3</sub>); ir  $\nu$  max (film) 3420, 1600, 1585, 1500 cm<sup>-1</sup>; uv  $\lambda$  max (MeOH) 228 nm (log  $\epsilon$  4.20), 275 (3.56), 283 (3.54); <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  7.20-6.70 (4H,m) (C-2',4',5',6'), 5.88 (1H,s) (C-8), 4.42 (1H,brs) (NH), 3.86 (3H,s), 3.83 (3H,s), 3.80 (3H,s) (C-5,6,3' methoxys), 3.44 (3H,s) (C-7 methoxy), 1.35 (3H,s) (C-CH<sub>3</sub>), 1.24 (3H,s) (C-CH<sub>3</sub>); ms m/z 222(100), 206(12), 192(7), 161(8), 149(6), 121(8), 91(8), 77(6), 65(4).

1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1-(3'-methoxy- $\alpha,\alpha$ -dimethylbenzyl)-2-methylisoquinoline<sup>7</sup> (12).

Formalin (37%) (10 ml) was added to 11 (1 g) in EtOH and stirred at room temperature for 0.5 hours. After cooling in ice, NaBH<sub>4</sub> (3 g) was gradually added and the mixture stirred for 2 hours. The reaction mixture was evaporated, H<sub>2</sub>O added and subsequently extracted with Et<sub>2</sub>O. The Et<sub>2</sub>O layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to yield 12 (0.76 g) (73%) as plates; mp 170-1°C (ether); ir  $\nu$  max (KBr) 2960, 1635, 1600, 1510 cm<sup>-1</sup>; uv  $\lambda$  max (MeOH) 228 nm (log  $\epsilon$  4.30), 262 (4.29), 320 (2.80); <sup>1</sup>H-nmr (CDCl<sub>3</sub>)  $\delta$  7.20-6.60 (4H,m) (C-2',4',5',6') 5.65 (1H,s) (C-8), 3.80 (3H,s), 3.75 (3H,s), 3.69 (3H,s) (C-5,6,3' methoxys), 3.42 (3H,s) (C-7 methoxy), 2.49 (3H,s) (N-CH<sub>3</sub>), 1.41 (3H,s) (C-CH<sub>3</sub>), 1.20 (3H,s) (C-CH<sub>3</sub>); ms m/z 236(100), 220(10), 206(9), 194(8), 175(5), 149(6), 121(5), 91(5).

N,O,O-Trimethyltetrahydromelosmine (1,2,3,9-Tetramethoxy-7,7-dimethylaporphine) (3). Thalium trifluoroacetate (TFA) (150 mg) in TFA (120 ml) was cooled to -40°C and a solution of 12 (60 mg) in CH<sub>2</sub>Cl<sub>2</sub> (5ml) and BF<sub>3</sub>-Et<sub>2</sub>O (1 ml) added in one batch. The reaction mixture was stirred (3 hr) at -40°C. After evaporation, H<sub>2</sub>O was added and the pH adjusted to 9 with 5% NH<sub>4</sub>OH. The solution was extracted with chloroform, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to afford 3 (5 mg) (9%). The product was identical (UV, IR, MS) to the target compound<sup>1</sup>.

ACKNOWLEDGEMENTS AND REFERENCES

1. V. Zabel, W.H. Watson, C.H. Phoebe, Jr., J.E. Knapp, P.L. Schiff, Jr. and D.J. Slatkin, J. Nat. Prod., 1982, 45, 94.
2. S. Kubota, T. Masui, E. Fujita and S.M. Kupchan, J. Org. Chem., 1966, 31, 516.
3. W. Adam and H. Fick, J. Org. Chem., 1978, 43, 772.
4. N.O.V. Sonntag, Chem. Rev., 1953, 52, 258.
5. W.M. Whaley and T.R. Govindachari, Org. React., 1951, 6, 74.
6. E.C. Taylor, J.G. Andrade, G.J.H. Rall and A. McKillop, J. Am. Chem. Soc., 1980, 102, 6513.
7. The authors are grateful to Kurt L. Loening, Nomenclature Director, Chemical Abstracts Service, Columbus, Ohio for providing the correct IUPAC name of this compound.

Received, 13th September, 1982