


REINVESTIGATION OF A RECENT 1,3,4-BENZOTRIAZEPINE SYNTHESIS,
FORMATION OF 1,3,4-OXADIAZOLES

Endre Tihanyi*, Melinda Gál and Péter Dvortsák

Institute for Drug Research, Budapest, P.O.B.82, H-1325 Hungary


Abstract - It was shown by spectroscopic and synthetic methods that the reaction of anthranilic acid hydrazides with 1,1'-carbonylbisimidazole (CO/Im/2) afforded, in contrast with earlier literature data, no 1,3,4-benzotriazepines or quinazolines but 1,3,4-oxadiazole derivatives.

Langis and Charest¹, later Hromatka et al², further Bailey³ reported methods for the preparation of the 3,4-dihydro-1H-1,3,4-benzotriazepine-2,5-diones 1. Peet and Sunder⁴ reexamined these synthetic routes and proved, that Langis¹ and Bailey's³ compounds were quinazoline-2,4-diones 2 instead of 1. The synthesis starting with anthranilic acid N,N'-dimethylhydrazides and phosgene described by Hromatka² was successful only for the desired benzotriazepines 3, by analogy to the known method for the preparation of this ring system⁵. Namely in this case the N-substitution of the hydrazide moiety hindered the formation of the six-membered ring, which was generally thermodynamically favored over the seven-membered triazepine ring.




Since 1975, only a few authors have published new results in this field^{6,7}. The last one was a simple synthesis from anthranilic acid hydrazides 5 and CO/Im/2 reported by Davidson⁸ to afford 1,3,4-benzotriazepines 4.


We reproduced his method ($R_1 \rightarrow H$) and established that the structure of the product (6) was not 4 as was claimed⁸.



Though elementary analyses and mass spectra did not exclude the benzotriazepine or quinazoline structures, the IR spectrum contained no amide bands, but a very strong absorption band at 1780 cm^{-1} . The missing CO-NH band suggests that the hydrazide carbonyl may have taken part in the reaction. Dimethylation of 6 didn't give the known 7,^{4,9} but a novel compound 8, which also has no CO-NH bands in the IR spectrum. In the ^1H NMR spectrum an N-CH₃ and an NH-CH₃ group were identified, so we assigned the structure of a 1,3,4-oxadiazol-5-one 8 to this product. Hence 6 must be the corresponding non-methylated oxadiazole. When N,N-dimethylantranilic acid hydrazide 9 and CO/Im/2 were reacted analogously, the same type of reaction was observed and 10 was isolated. This was methylated to give 11. The close similarity of the IR spectra of 6, 8, 10 and 11¹⁰ have also pointed to the similar structures.



In light of the above results, in the reaction of 5 with CO/Im/2, we do suppose an intermediate 13 rather than 12 which had earlier been suggested⁸. The rearrangement of 6 (erroneously proposed⁸ to be the benzotriazepine 14) to quinazoline 15 observed by Davidson may be explained to proceed via 14.



Syntheses of 1,3,4-oxadiazoles from acid hydrazides are known¹¹, in which the one carbon fragment was from orthoesters, imidoesters, phosgene, thiophosgene, isocyanide dichlorides etc., but 1,1'-carbonylbisimidazole has not been applied for this purpose before.

ACKNOWLEDGEMENTS

The authors wish to thank Miss R. Andersen and Miss Gy. László for technical assistance.

REFERENCES AND NOTES

1. A.L. Langis and M.P. Charest, Chim. Ther., 1967, 394.
2. O. Hromatka, F. Krenmüller and M. Knollmüller, Monatsh. Chem., 1969, 100, 934.
3. D.M. Bailey. U.S. Patent 3 607 866 (1971); C.A., 1971, 75, 140910v.
4. N.P. Peet and S. Sunder, J.Org. Chem., 1975, 40, 1909.
5. M. Busch, Chem. Ber., 1894, 27, 2901.
6. S. Sunder and N.P. Peet, J. Org. Chem., 1977, 42, 2551.
7. I. Bitter, L. Szőcs and L. Tőke, Acta Chim. Hung., 1981, 107, 171.
8. J.S. Davidson, Abstracts of Eighth Intern. Congr. of Heterocyclic Chem., p. 214 (1981).
9. Methylation of both 3,4-dihydro-3-methyl- and 3,4-dihydro-4-methyl-1H-1,3,4-benzotriazepine-2,5-dione give 3⁴.

10. All compounds gave satisfactory microanalytical data.

6 mp 175-176 °C; 55 % yield; δ (DMSO- d_6) 6.2 b 2H, NH_2 ; 6.63-7.45 m 4H, Ar-H; ν 3470, 3370, 3140, 1770 cm^{-1}

8 mp 179-180 °C; 85 % yield; δ (DMSO- d_6) 2.88 d 3H, $NH-CH_3$; 3.25 s 3H, $N-CH_3$; 6.6 b 1H, NH; 6.6-7.7 m 4H, Ar-H; ν 3400, 1780 cm^{-1}

10 mp 108-110 °C; 72 % yield; δ ($CDCl_3$) 2.80 s 6H, $N(CH_3)_2$; 6.8-7.7 m 4H, Ar-H; 10.4 b 1H, NH; ν 3100 b, 1780 cm^{-1}

11 honey; 90 % yield; δ ($CDCl_3$) 2.80 s 6H, $N(CH_3)_2$; 3.50 s 3H, $N-CH_3$; 6.8-7.7 m 4H, Ar-H; ν 1780 cm^{-1}

11. D. Barton and W.D. Ollis, 'Comprehensive Organic Chemistry', ed. by P.G. Sammes, Pergamon Press, London, 4, 1024 (1979).

Received, 29th November, 1982