

THE REACTION OF 2-SUBSTITUTED QUINOLINE 1-OXIDES WITH DIMETHYL  
ACETYLENEDICARBOXYLATE: FORMATION OF 1-BENZAZEPINE DERIVATIVES

Yasuhisa Ishiguro, Kazuhisa Funakoshi, Seitaro Saeki, Masatomo Hamana<sup>1\*</sup>

Faculty of Pharmaceutical Sciences, Kyushu University

Maidashi 3-1-1, Higashi-ku, Fukuoka 812, Japan

Ikuhiko Ueda and Shigeaki Kawano

College of General Education, Kyushu University

Ropponmatsu 4-2-1, Chuo-ku, Fukuoka 810, Japan

**Abstract** — The 1,3-dipolar cycloaddition of 2-phenylquinoline 1-oxide (1) and quinaldine 1-oxide (3) with dimethyl acetylenedicarboxylate gives ring expanded products, 2-phenyl- (2) and 2-methyl-3H-3-methoxalyl-3-methoxycarbonyl-1-benzazepines (5), respectively. The reaction of quinaldonitrile 1-oxide (4) yields 1H-2-cyano-3-methoxycarbonyl-1-benzazepine (6) accompanied by further demethoxalylolation.

In the course of our continuing investigation of the 1,3-dipolar cycloaddition of substituted quinoline 1-oxides with dimethyl acetylenedicarboxylate (DMAD)<sup>2</sup>, an interesting ring expansion was observed in the reaction of some 2-substituted quinoline 1-oxides.

A solution of 2-phenylquinoline 1-oxide (1) and DMAD (1.2 equiv.) in dichloromethane was kept at room temperature for 3 days to give 3H-3-methoxalyl-3-methoxycarbonyl-2-phenyl-1-benzazepine (2), colorless prisms, mp 94-96°C, in a high yield of 87.8%. Its structure was deduced from the elemental analyses, the IR and PMR spectroscopies<sup>3</sup>, and established by an X-ray diffraction study<sup>4</sup>.

Subsequently the reaction of quinaldine 1-oxide (3) as well as that of quinaldonitrile 1-oxide (4) with DMAD was found to yield also a ring expanded product.

While treatment of 3 with DMAD at room temperature caused only resinification, the reaction proceeded at -10°C in dichloromethane, though very slowly, and the 3H-1-benzazepine (5) of the same type as 2 was obtained as an oil in 81.4% yield after 2 weeks' reaction. Although purification of 5 was not effected because of its

high instability, its structure could be confirmed by spectral data.

In contrast to the case of 3, the reaction of 4 with DMAD in dichloromethane did not occur at all at low temperature. When heated in DMF at 100°C, however, the reaction proceeded and 1H-2-cyano-3-methoxycarbonyl-1-benzazepine (6), orange needles, mp 122-125°C (dec.), was isolated in 28.2% yield after 1 h's reaction. These reactions are formulated in Chart 1.




Chart 1

The formation of 1-benzazepines (2, 5 and 6) is very significant in view of no precedent for the expansion of a pyridine ring to an azepine system in the 1,3-dipolar cycloaddition of aromatic N-oxides. The formation of 2 and 5 can be well rationalized by the following mechanism. The initially formed primary cycloadduct (A) readily undergoes a 3,5-sigmatropic shift leading to a 2,3-dihydroquinoline (B) followed by its valence isomerization to give a 3H-1-benzazepine, 2 or 5. Since heating is required for the reaction of 4 to proceed, the 3H-1-benzazepine (C) similarly formed undergoes thermal demethoxalation to give the thermodynamically more stable 1H-1-benzazepine derivative 6 (Chart 2).

It is also noticeable that these results present direct evidence for the intermediacy of the 2,3-dihydroquinoline (B) of an azanorcaradiene type as well as of the analogous dihydropyridine which was postulated in a variety of 1,3-dipolar cycloadditions of quinoline and pyridine N-oxides<sup>2c,5</sup>.




Chart 2

## REFERENCES AND NOTE

1. Present address: Central Research Laboratories, Chugai Pharmaceutical Co., Ltd., Takada 3-41-8, Toshima-ku, Tokyo 171, Japan.
2. a) M. Hamana, H. Noda, and M. Aoyama, *Heterocycles*, 1974, 2, 167; b) H. Noda, T. Yamamori, M. Yoshida, and M. Hamana, *Heterocycles*, 1976, 4, 453; c) Y. Ishiguro, K. Funakoshi, S. Saeki, M. Hamana, and I. Ueda, *Heterocycles*, 1980, 14, 179; d) Y. Ishiguro, K. Funakoshi, S. Saeki, M. Hamana, I. Ueda, and S. Kawano, *Heterocycles*, 1983, 20, 193.
3. IR (Nujol)  $\text{cm}^{-1}$ : 1735, 1760 (C=O). PMR ( $\text{CDCl}_3$ )  $\delta$ : 3.41 (3H, s,  $\text{COOCH}_3$ ), 3.58 (3H, s,  $\text{COOCH}_3$ ), 6.59 (1H, d,  $J_{4,5}=10$  Hz,  $\text{C}_4$ - or  $\text{C}_5$ -H), 6.99 (1H, d,  $J_{4,5}=10$  Hz,  $\text{C}_4$ - or  $\text{C}_5$ -H), 7.10-7.60 (9H, m, Ar-H).
4. S. Kawano, I. Ueda, Y. Ishiguro, K. Funakoshi, S. Saeki, and M. Hamana, *Act. Cryst.*, 1983, in press.
5. R.A. Abramovitch and I. Shinkai, *Accts. Chem. Res.*, 1976, 9, 192.

Received, 9th May, 1983