

SYNTHESIS OF TRISUBSTITUTED 4-OXEPANONES BY THE LEWIS ACID-PROMOTED THREE-COMPONENT RING-EXPANSION REACTION OF CYCLOPROPAPYRANONES, SILYL ENOLATES AND GLYOXYLATES

Yoshiaki Sugita,* Chikasa Kimura, and Ichiro Yokoe


Faculty of Pharmaceutical Sciences, Josai University, Sakado, Saitama 350-0295, Japan

Abstract – In the presence of SnCl_4 , cyclopropapyranones easily reacted with silyl enolates and ethyl glyoxylate to give the trisubstituted 4-oxepanones in good yields.

The seven-membered oxacycles are an important class of oxygenated heterocyclic compounds that have attracted the attention of organic chemists and biochemists due to their biological activities and occurrence in natural products.¹ Various methods for the construction of the seven-membered oxacycles have been reported.²

We recently reported the synthesis of 2,7-disubstituted 4-oxepanones by the Lewis acid-promoted ring-opening addition reactions of cyclopropapyranones (**1**) with silyl enolates as nucleophiles (eq. 1, Scheme 1).³ From these reactions, the *trans*-isomers were mainly obtained.

We expected that the treatment of a cyclopropapyranone derivative with a Lewis acid readily generates a cyclic 1,3-zwitterion,⁴ which can react with both silyl enolates and appropriate electrophiles to produce the trisubstituted 4-oxepanones (eq. 2). A cyclopropane with donor and acceptor substituents at the vicinal position on the cyclopropane ring is an equivalent of a ring-opened 1,3-zwitterion. Therefore, cyclopropapyranone (**1**) is recognized as a good reactant for electrophiles and nucleophiles.⁵ We now report the synthesis of trisubstituted 4-oxepanones by the Lewis acid-promoted three-component ring-expansion reaction of cyclopropapyranones as the synthetic equivalents of cyclic 1,3-zwitterions.



Scheme 1

In our previous study, we have found that when the ring-opening addition reaction of the cyclopropapyranone (**1a**) with ketene silyl acetal (**2a**) was carried out, the use of SnCl_4 (0.3 eq.) as a Lewis acid gave the best result and the corresponding oxepanone (**5a**) was obtained in 71% yield with

moderate *trans*-selectivity.³ The reaction of **1a**, **2a** with ethyl glyoxylate (**3a**)⁶ was chosen as the model, and several reaction conditions were examined. Cyclopropapyranone (**1a**) and ketene silyl acetal (**2a**) in CH₂Cl₂ were reacted with SnCl₄ (0.3 eq.) at -78 °C for 10 min, and the resulting mixture was treated with ethyl glyoxylate (**3a**) to give the desired oxepanone (**4**) in 22% yield. However, the adduct (**5a**) was mainly obtained because the aldol type reaction as the second step did not completely occur using a catalytic amount of SnCl₄ (Table 1, Entry 1). In contrast, it was found that the use of a stoichiometric amount of SnCl₄ for the addition of **3a** resulted in an improvement of the chemical yield of the desired oxepanone from 22% to 48% (Entry 2). We also found that the addition of powdered molecular sieves 4A⁷ (MA 4A) to the reaction mixture was effective for obtaining the desired product in good yield (77% yield) although the role of MS 4A is not clear (Entry 3).⁸ In this reaction, the order of the addition of the reagents dramatically influenced the yield. Namely, the addition of **2a** to the reaction mixture of **1a** and **3a** led to the formation of a complex mixture not containing the desired product.





Table 1. Three-component reaction of **1a**, **2a** and ethyl glyoxylate (**3a**)

Entry	SnCl ₄ (eq.)	MS 4A	Yield 4 /%	Yield 5a /%
1	0	-	22	53
2	1.0	-	48	21
3	1.0	+	77	0

The obtained product (**4**) could be converted into four stereoisomeric oxepanones (**6**) by the mesylation of the hydroxy group of **4** followed by an elimination reaction (Scheme 2). The ratio of the stereoisomers (TE : CE : TZ : CZ) was determined as 71:19:8:2 by ¹H-NMR, and the stereochemical assignment of the isomers was established by the analysis of their NOE experiments (Figure 1). It was noted that in the ¹H-NMR spectrum the olefinic proton signal of the *Z*-isomers (TZ- and CZ-**6**) appeared at high field compared with that of the *E*-isomers (TE- and CE-**6**).



Scheme 2




Figure 1

Several examples of the three-component ring-expansion reaction of **1a** were examined and these results are summarized in Table 2 (Entries 1-4). For the reaction of **1a**, **2a**, and methyl glyoxylate (**3b**),⁶ a similar tendency was observed (Entry 2). As the nucleophiles, silyl enol ethers (**2b**, **2c**) also reacted with **1a** under similar conditions to give the corresponding oxepanones (**8**,⁹ **9**), even though a slight decrease in yield was observed compared to that of the reactions of the ketene silyl acetal (**2a**) (Entries 3, 4).




Table 2. Three-component reaction of cyclopropapyranones (**1**), silyl enolates (**2**) and glyoxylates (**3**)

Entry	Cyclopropane	Nucleophile	Glyoxylate	Product	Yield ^a / % (TE : CE : TZ : CZ) ^b
1			3a		66 (71 : 19 : 8 : 2)
2	1a		3b		57 (73 : 17 : 5 : 5)
3	1a		3a		48 (78 : 13 : 8 : 1)
4	1a		3a		26 ^c (81 : 10 : 8 : 1)
5			3a		57 (81 : 12 : 5 : 2)
6	1b		3b		58 (80 : 10 : 9 : 1)
7			3b		66 (80 : 15 : 5 : - ^d)

^a Total yield from **1**. ^b The ratio of the stereoisomers was determined by 500 MHz ¹H-NMR. ^c Disubstituted 4-oxepanone (**5**) was also obtained in 45%. ^d Not detected.

In order to extend the scope and synthetic utility of this methodology, the reactions of other cyclopropapyranones under similar conditions were examined. The reactions of **1b**,³ **2a**, and alkyl glyoxylates (**3a**, **3b**) gave the corresponding oxepanones (**10**,⁹ **11**) in good yields (Entries 5, 6). Under similar conditions, the chiral cyclopropapyranone (**1c**)³ also reacted with **2a** and **3b** to give the corresponding chiral oxepanone (**12**) in 66% yield with moderate stereoselectivity (Entry 7). In all the reactions, the *trans-E*-isomer was mainly obtained.

In summary, we have demonstrated that the SnCl₄-promoted three-component ring-expansion reaction of cyclopropapyranones, silyl enolates, and glyoxylates smoothly proceeded to afford the corresponding trisubstituted 4-oxepanones in good yields with moderate stereoselectivity. Further applications of this reaction are now under investigation in our laboratory.

A typical experimental procedure is described for the reaction of cyclopropane (**1a**), ketene silyl acetal (**2a**), and ethyl glyoxylate (**3a**): To a mixture of **1a** (75 mg, 0.4 mmol), **2a** (104 mg, 0.6 mmol), and molecular sieves 4A (150 mg) in CH₂Cl₂ (3 mL) was added SnCl₄ (1.0 M sol. in CH₂Cl₂, 0.12 mL, 0.12 mmol) at –78 °C under an argon atmosphere. After the mixture was stirred for 10 min, **3a** (82 mg, 0.8 mmol) and SnCl₄ (1.0 M sol. in CH₂Cl₂, 0.4 mL, 0.4 mmol) were successively added. After being stirred for 1 h at the same temperature, the reaction was quenched with saturated aqueous NaHCO₃. The mixture was filtered through a celite pad, and the filtrate was concentrated. The residue was dissolved in THF-1*N* HCl (2:1, 4 mL) and stirred for 10 min at 0 °C. The mixture was then extracted with ether. After the combined organic layers were dried and concentrated, the residue was chromatographed on silica gel to produce **4** (121 mg, 77%). To a solution of **4** (117 mg, 0.3 mmol), Et₃N (121 mg, 1.2 mmol), and DMAP (4 mg, 0.03 mmol) in CH₂Cl₂ (4 mL) was added MsCl (69 mg, 0.6 mmol) in CH₂Cl₂ (0.5 mL) at 0 °C. After being stirred for 1 h at rt, DBU (91 mg, 0.6 mmol) was added to the mixture and refluxed for 2 h. After an extractive work up and purification by silica gel column chromatography, the corresponding oxepanone (**6**) was obtained in 85% yield (95 mg).

REFERENCES AND NOTES

1. For reviews, see: T. Yamamoto and M. Murata, *Chem. Rev.*, 1993, **93**, 1897; D. J. Faulkner, *Nat. Prod. Res.*, 1997, **14**, 259; 1998, **15**, 113.
2. For examples, see: G. C. Fu, S. T. Nguyen, and R. H. Grubbs, *J. Am. Chem. Soc.*, 1993, **115**, 9856; I. Kadota, P. Jung-Youl, N. Koumura, G. Pollaud, Y. Matsukawa, and Y. Yamamoto, *Tetrahedron Lett.*, 1995, **36**, 5777; P. A. Evans and J. D. Roseman, *J. Org. Chem.*, 1996, **61**, 2252; T. Nakata, S. Nomura, and H. Matsukura, *Tetrahedron Lett.*, 1996, **37**, 213; D. Berger, L. E. Overman, and P. A. Renhowe, *J. Am. Chem. Soc.*, 1997, **119**, 2446; J. O. Hoberg, *J. Org. Chem.*, 1997, **62**, 6615; K. Fujiwara, H. Mishima, A. Amano, T. Tokiwano, and A. Murai, *Tetrahedron Lett.*, 1998, **39**, 393.
3. Y. Sugita, C. Kimura, H. Hosoya, S. Yamadoi, and I. Yokoe, *Tetrahedron Lett.*, 2001, **42**, 1095.
4. Y. Sugita, K. Kawai, H. Hosoya, and I. Yokoe, *Heterocycles*, 1999, **51**, 2029.
5. H.-U. Reissig, *Tetrahedron Lett.*, 1981, **22**, 2981; K. Saigo, S. Shimada, and M. Hasegawa, *Chem.*

- Lett.*, 1990, 905; K. Saigo, S. Shimada, T. Shibasaki, and M. Hasegawa, *Chem. Lett.*, 1990, 1093; M. Komatsu, I. Suehiro, Y. Horiguchi, and I. Kuwajima, *Synlett*, 1991, 771; S. Shimada, Y. Hashimoto, A. Sudo, M. Hasegawa, and K. Saigo, *J. Org. Chem.*, 1992, **57**, 7126; S. Shimada, Y. Hashimoto, T. Nagashima, M. Hasegawa, and K. Saigo, *Tetrahedron*, 1993, **49**, 1589; S. Shimada, Y. Hashimoto, and K. Saigo, *J. Org. Chem.*, 1993, **58**, 5226; Y. Sugita, K. Kawai, and I. Yokoe, *Heterocycles*, 2000, **53**, 657; 2001, **55**, 135.
6. J. M. Hook, *Synth. Commun.*, 1984, **14**, 83.
 7. MS 4A was used after drying at 150 °C for 6 h under reduced pressure.
 8. S. Kobayashi, I. Hachiya, H. Ishitani, and M. Araki, *Tetrahedron Lett.*, 1993, **34**, 4535; G. H. Posner, H. Dai, D. S. Bull, J.-K. Lee, F. Eydoux, Y. Ishihara, W. Welsh, N. Pryor, and S. Petr, Jr., *J. Org. Chem.*, 1996, **61**, 671; M. Bougauchi, S. Watanabe, T. Arai, H. Sasai, and M. Shibasaki, *J. Am. Chem. Soc.*, 1997, **119**, 2329; E. Keller, J. Maurer, R. Naasz, T. Schader, A. Meetsma, and B. L. Feringa, *Tetrahedron: Asymmetry*, 1998, **9**, 2409; L. P.-Serrano, L. Casarrubios, G. Dominguez, and J. P.-Castells, *Org. Lett.*, 1999, **1**, 1187; T. Tokiwano, K. Fujiwara, and A. Murai, *Chem. Lett.*, 2000, 272.
 9. The stereochemical assignment of the compounds (TE-8, CE-8, and TZ-10) was also mainly established by analysis of their NOE experiments (see figure below).

