

REACTION OF 2-CHLORO-1-AZAAZULENE WITH DIPHENYL-CYCLOPROPENONE; DIMERIZATION OF THE CYCLOADDUCT ATTENDED BY *cine*-SUBSTITUTION AND REARRANGEMENT

Noritaka Abe,* Hiroyuki Fujii, and Akikazu Kakehi[†]

Department of Chemistry, Faculty of Science, Yamaguchi University,
Yamaguchi 753-8512, Japan

[†] Department of Chemistry and Material Engineering, Faculty of Engineering,
Shinshu University, Wakasato Nagano 380-8553, Japan

Abstract — 2-Chloro-1-azaazulene reacted with diphenylcyclopropenone to give 3,4-diphenyl-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulene-2,5-dione and 6-[(3,4-diphenyl-5-oxo-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulen-2-ylidene)-methyl]-3,4-diphenyl-2a,5-dihydro-2a-azaacenaphtylen-5-one; the latter would be produced *via* cycloaddition and a successive *cine*-substitution followed by rearrangement.

INTRODUCTION

The chemistry of azaazulenes is of interest for their physical and chemical properties as well as physiological properties,¹ in comparison with those of azulenes. Cycloadditions of diphenylcyclopropenone (DPP) with heterocycles are interesting for the synthetic viewpoint that the reaction gave a some variety of fused heterocycles;^{2–8} we reported the cycloadditions of 2-amino- and 2-hydrazino-1-azaazulenes with DPP,^{9–11} where variable cycloadducts were obtained being dependent on the nature of substituents on the 1-azaazulenes. To expand the reaction, we examined the reaction of 2-chloro-1-azaazulene (**1**) with DPP, and found that the reaction proceeded a different pathway from that of 2-amino-1-azaazulenes with DPP and gave an interesting dimer of 1 : 1-adduct.

RESULTS AND DISCUSSION

The treatment of 2-chloro-1-azaazulene (**1**) with 1.5 equivalent molar of DPP in acetonitrile under reflux for 3 h gave two kinds of cycloadducts (**2**) (4%) and (**3**) (19%) along with recovered **1** (58%). The reaction at higher temperature did not improve the results; when the reaction was performed in refluxing *tert*-butylbenzene for 1 h, the reaction showed a complex feature and the compound (**3**) was isolated in 19% yield together with recovered **1** (55%). The compound (**2**) was analyzed as C₂₄H₁₅NO₂ from its MS spectrum and elemental analysis; the compound (**2**) would be a hydrolyzed product of the 1 : 1-adduct of **1** with DPP. In the IR spectrum of **2**, two carbonyl peaks appeared at 1744 and 1629 cm^{–1}. In its ¹H

NMR spectrum, a ^1H singlet appeared at δ 5.82, and four seven-membered protons were seen at δ 6.84, (1H, dd, J 11.0, 8.3), 6.93 (1H, dd, J 11.6, 8.3), 7.20 (1H, d, J 11.0), and 7.87 (1H, d, J 11.6). From the results, the compound (**2**) was deduced as 3,4-diphenyl-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulene-2,5-dione. The compound (**3**) was analyzed as $\text{C}_{48}\text{H}_{30}\text{N}_2\text{O}_2$ from its MS spectrum and elemental analysis. Its ^1H NMR spectrum showed rather complex feature and the IR spectrum showed the signals at 1632 and 1606 cm^{-1} assignable to carbonyl group; the structure could not be deduced from the spectral analysis. Eventually, the structure of **3** was determined as 6-[(3,4-diphenyl-5-oxo-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulen-2-ylidene)methyl]-3,4-diphenyl-2a,5-dihydro-2a-azaacenaphtylen-5-one by X-Ray structure analysis. The ORTEP drawing¹² of **3** is shown in Figure 1.





Figure 1. An ORTEP drawing of **3** with thermal ellipsoids (50% probability).

The formation of **2** and **3** showed that the reaction occurred at N-1 position of **1** at first. The result was a contrast to the reaction of 2-dialkylamino-1-azaazulene with DPP,¹¹ where the reaction occurred at C-3 position at first and gave **4**. It is considered that the reactivity of C-3 position of **1** would be lower than

that of 2-diethylamino-1-azaazulene, in which C-3 carbon was contained in an enamine moiety. The results was consisted with the molecular orbital calculation by Gaussian 98 using RHF/6-31G* (Figure 2). Furthermore, 2-diethylamino-1-azaazulene have a rather large lobe at C-3 position in its HOMO.

Plausible reaction mechanism is shown in the Scheme 1. Cycloaddition of 2-chloro-1-azaazulene with DPP produced **A** at first. A hydration of **A** and a successive elimination of hydrogen chloride followed by a dehydrogenation furnished **2**. The attempt of the isolation of intermediate (**A**) was failed; the pale yellow compound turned to yellow in the silica gel chlounn. Therefore, it is thought that the elimination-addition would occur in the silica gel chlounn. A hydrogen shift of **A** led to **B**. When a *cine*-substitution of **B** with **A** occurred, the dimer (**C**) could be generated. Electrocyclic reaction of the cycloheptatriene moiety of **C** gave the norcaradiene derivative (**D**). Ring contraction of **D** attended by dehydrochlorination furnished **3**.



Scheme 1




Figure 2. The molecular orbital calculation of **3** and 2-diethylamino-1-azaazulene

EXPERIMENTAL

Mps are measured using a Yanagimoto micro-melting apparatus and uncorrected. ^1H NMR spectra were recorded on a Bruker AVANCE 400S (400 MHz) using deuteriochloroform as a solvent with tetramethylsilane as an internal standard unless otherwise stated; J values are recorded in Hz. IR spectra were recorded for KBr pellets on a Nicolet FT-IR Impact 410. MS spectra were taken with on an LC-MS Waters Integrity System. Elemental analyses were taken with a Perkin Elmer 2400II. Kieselgel 60 and was used for column chromatography and Kieselgel 60G was used for thin-layer chromatography.

Reaction of 2-chloro-1-azaazulene (1) with diphenylcyclopropenone

A solution of 2-chloro-1-azaazulene (**1**) (0.330 g, 2.02 mmol) and DPP (0.630 g, 3.05 mmol) in dry acetonitrile (30 mL) was heated under reflux for 3 h, then the solvent was evaporated. The residue was chromatographed on silica gel column with chloroform to give recovered **1** (0.190 g, 58%), 3,4-diphenyl-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulene-2,5-dione (**2**) (0.012 g, 4%), and 6-[(3,4-diphenyl-5-oxo-2a,5-dihydro-2*H*-2a-azabenz[*cd*]azulen-2-ylidene)methyl]-3,4-diphenyl-2a,5-dihydro-2a-azaacenaphthylen-5-one (**3**) (0.128 g, 19%), successively.

Similar treatment of **1** (0.330 g, 2.02 mmol) with DPP (0.420 g, 2.04 mmol) in refluxing *tert*-butylbenzene (30 mL) for 1 h gave **1** (0.182 g, 55%) and **3** (0.130 g, 19%), and the treatment of **1** (0.330 g, 2.02 mmol) with DPP (0.630 g, 3.05 mmol) in refluxing xylene (30 mL) for 3 h gave **1** (0.182 g, 42%) and **3** (0.100 g, 15%).

2: Yellow needles (from hexane-dichloromethane), mp 185–187 °C; δ_{H} 5.82 (1H, s), 6.84, (1H, dd, J 11.0, 8.3), 6.93 (1H, dd, J 11.6, 8.3), 6.97–7.07 (2H, m), 7.10–7.17 (6H, m), 7.20 (1H, d, J 11.0), 7.21–7.25 (2H, m), and 7.87 (1H, d, J 11.6); ν_{max} / cm^{-1} 1744 and 1629 (C=O); m/z 350 ($\text{M}^+ + 1$, 100), 349 (M^+ , 98), 348 (97), 332 (31), 320 (92), 291 (90), 265 (24), 161 (51), and 146 (45). *Anal.* Calcd for $\text{C}_{24}\text{H}_{15}\text{NO}_2$: C, 82.50; H, 4.33; N, 4.01. Found: C, 82.61; H, 4.55; N, 4.05.

3: Violet prisms (from hexane-dichloromethane), mp 216–218 °C; δ_{H} 5.80–5.90 (2H, m), 6.12–6.18 (1H, m), 6.27 (1H, s), 6.56 (1H, s), 6.74 (1H, d, J 3.5), 6.92–7.34 (22H, m), 7.41 (1H, d, J 8.0), and 7.82 (1H, d, J 7.9); ν_{max} / cm^{-1} 1632 and 1606 (C=O); m/z (rel intensity) 667 ($\text{M}^+ + 1$, 7), 666 (M^+ , 2), 589 (11), 487 (17), 346 (36), 333 (22), 178 (100), and 149 (36). *Anal.* Calcd for $\text{C}_{48}\text{H}_{30}\text{N}_2\text{O}_2 \cdot 1/4 \text{CH}_2\text{Cl}_2$: C, 84.23; H, 4.47; N, 4.07. Found: C, 84.56; H, 4.65; N, 4.05.

X-Ray structure determination

Crystal data of 3: Black prism, $\text{C}_{48}\text{H}_{30}\text{N}_2\text{O}_2 \cdot \text{CH}_2\text{Cl}_2$, $M=751.71$, monoclinic, space group $\text{P2}_1/\text{a}$, $a=11.743(4)$, $b=19.016(5)$, $c=17.464(4)$ Å, $\beta=103.02(3)^\circ$, $V=3799(2)$ Å³, $Z=4$, $D_{\text{calcd}}=1.314$ g/cm³, crystal dimensions 0.30 x 0.42 x 0.88 mm. Data were measured on a Rigaku AFC 5S radiation diffractometer with graphite-monochromated Mo-K α radiation. A total 9426 reflections (8997 unique) were collected using ω - 2θ scan technique with in a 2θ range of 55.0° . The structure was solved by direct methods and refined by a full-matrix least-squares methods using 550 variables refined with 2704 reflections [$I > 3\sigma(I)$]. The weighting scheme $\omega = 4Fo^2/\sigma^2(Fo^2)$ gave satisfactory agreement analyses. The final refinement converged to $R = 0.063$ and $R_w = 0.072$.

REFERENCES

1. For review see, N. Abe, 'Recent Research Developments in Organic and Bioorganic Chemistry: Recent Progress in the Chemistry of Azaazulenes,' Vol. 4, Transworld Research Network, 2001; T. Nishiwaki and N. Abe, *Heterocycles*, 1981, **15**, 547; M. Kimura, *Yuki Gosei Kagaku Kyokai Shi*, 1981, **39**, 690.
2. For review see, K. T. Potts and J. S. Baum, *Chem. Rev.*, 1974, **74**, 189; T. Eicher and J. L. Weber, *Top. Curr. Chem.*, 1975, **57**, 1; K. Matsumoto, *Yuki Gosei Kagaku Kyokai Shi*, 1972, **30**, 1035.
3. J. W. Lown and K. Matsumoto, *Can. J. Chem.*, 1971, **49**, 1165; 1971, **49**, 3119.
4. A. Kascheres and J. A. R. Rodrigues, *J. Org. Chem.*, 1975, **40**, 1440; A. Kascheres, C. Kascheres, J. A. R. Rodrigues, and A. R. A. Santana, *J. Org. Chem.*, 1976, **41**, 3546; A. Kascheres, L. J. R. Reyes, and S. M. Fonseca, *Heterocycles*, 1984, **22**, 2529.; A. Kascheres and D. Marchi, Jr., *J. Org. Chem.*, 1975, **40**, 2985; A. Kascheres, C. Kascheres, and J. A. R. Rodrigues, *Synth. Commun.*, 1984, 14, 905.
5. Y. Hayashi and H. Nozaki, *Tetrahedron*, 1971, **27**, 3085.
6. T. L. Gilchrist, C. J. Harris, C. J. Moody, and C. W. Rees, *J. Chem. Soc., Perkin Trans. 1*, **1975**, 1969.
7. T. Eicher, F. Abdesaken, G. Franke, and J. L. Weber, *Tetrahedron Lett.*, **1975**, 3915; T. Eicher, J. L. Weber, and G. Chatila, *Liebigs Ann. Chem.*, **1978**, 1203; T. Eicher and G. Franke, *Liebigs Ann. Chem.*, **1981**, 1337; T. Eicher and R. Rohde, *Synthesis*, **1985**, 619.
8. D. H. Wadsworth, S. L. Bender, D. L. Smith, and H. R. Luss, *Tetrahedron Lett.*, 1981, **22**, 3569; D. H. Wadsworth, S. L. Bender, D. L. Smith, H. R. Luss, and C. H. Weidner, *J. Org. Chem.*, 1986, **51**, 4639.
9. N. Abe and A. Kakehi, *Heterocycles*, 1993, **36**, 1961.
10. N. Abe and A. Kakehi, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 2487.
11. N. Abe, T. Murafuji, Y. Sugihara, and A. Kakehi, *Heterocycles*, 1995, **41**, 2289
12. C. K. Johnson, ORTEP II, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1976.