

HETEROCYCLES, Vol. 78, No. 1, 2009, pp.45 - 57. © The Japan Institute of Heterocyclic Chemistry
Received, 8th May, 2008, Accepted, 5th September, 2008, Published online, 8th September, 2008.
DOI: 10.3987/COM-08-11432

CYCLOTHIOMETHYLATION OF FUNCTIONAL SUBSTITUTED ANILINES BY CH₂O AND H₂S

Vnira R. Akhmetova,^{1*} Guzel R. Nadyrgulova,¹ Zalifa T. Niatshina,² Regina R. Khairullina,^{1,2} Zoya A. Starikova,³ Alexandra O. Borisova,³ Michail Yu. Antipin,³ Raihana V. Kunakova,² and Usein M. Dzhemilev¹

¹Institute of Petrochemistry and Catalysis, RAS, 450075 Ufa, Prospekt Oktyabrya, 141, Russian Federation, e-mail address: ink@anrb.ru. ²Ufa State Academy of Economy and Service, 450077 Ufa, Chernyshevskii Street, 145, Russian Federation, e-mail address: utis@bashnet.ru. ³A.N. Nesmeyanov Institute of Elementoorganic Chemistry, RAS, 11999 Moscow, Vavilov Street, 28, Russian Federation, e-mail address: star@xray.ineos.ac.ru

Abstract – *N*-Aryl substituted 1,3,5-dithiazinanes have been synthesized (20–60 °C) in 32–95% yield by cyclocondensation of *o*-, *p*-aminobenzoic, 4-, 5-aminosalicylic acids, *p*-aminobenzoic acid ethyl or (β -diethylamino)ethyl esters and *p*-aniline sulfamide with CH₂O and H₂S (1:3:2 ratio). At ambient temperature *p*-aminobenzoic acid, 5-aminosalicylic acid and *p*-aminobenzoic acid ethyl ester together with 1,3,5-dithiazinanes form appropriate *N*-aryl substituted 1,3-thiazetidines. Cyclothiomethylation of *p*-aminobenzoic acid ethyl ester (0 °C) results in *N,N*-diaryl-1,3,5-thiadiazinane. Heterocyclization of *p*-aniline sulfamide with CH₂O and H₂S at a molar ratio of 1:6:4 (pH 1.3–1.5) was found to afford *bis*-1,3,5-dithiazinanes involving two amino groups. Condensation of *p*-aniline sulfacetamide with 37% formalin (CH₂O) and H₂S (pH 2.5) led to the compound, which was built as macroheterocycle from two fragments of *p*-aniline sulfacetamide molecule bound to each other by *sym*-dimethyl sulfide chain (CH₂SCH₂).

INTRODUCTION

In organic chemistry there is an original method for the synthesis of 1,3,5-dithiazinane^{la-c} based on Wohl reaction namely multimolecular cyclocondensation of methyl amine, 37% formalin (CH₂O) and H₂S. Compounds, which contain dithiazinane ring, are used as insecticides and fungicides^{2a,b} as well as

additives modified and intensified product taste,^{2c-g} inhibitors, ferments,^{2h} complexons,^{3a-g} sorbents for gold and silver.^{4a,b}

For the last 5 years we systematically investigated the Wohl reaction using cyclothiomethylation of aliphatic^{5a} and aromatic^{5b} amines, amino acids,^{5c} aminoalcohols^{5d} aminophenols^{5e} as an example to obtain the corresponding *N*-substituted 1,3,5-dithiazinanes,^{5a-e} 1,3-thiazetidines^{5a,b,c} and 1,3,5-thiadiazinanes.^{5b} The direction of the cyclothiomethylation reaction depends on the structure of initial amines and reaction conditions (ratio and order of reagent mixing,^{5a,c} temperature,^{5a-e} medium pH), under which cyclocondensation is conducted. Thus, it was shown that in the cyclothiomethylation reaction of aliphatic^{5a} and aromatic^{5b} amines with CH₂O and H₂S (amine:CH₂O:H₂S = 1:3:2) the decrease in basicity of amines causes the increase in the yield of target 1,3,5-dithiazinanes. Further more thorough and deep analysis of cyclocondensation of functionally-substituted anilines, namely *o*-, *m*- and *p*-aminophenols, with CH₂O and H₂S has shown, that the direction of the cyclothiomethylation reaction of aminophenols depends on the position of functional amino group.^{5e} Aminophenols, *o*- and *p*-isomers, were established to interact with CH₂O and H₂S as a reactant mixture (3:2 ratio) to form 1,3,5-dithiazinanes. *m*-Aminophenol under analogous conditions undergoes intermolecular condensation with CH₂O and H₂S simultaneously involving OH and NH₂ groups to form macroheterocycle, which contains the fragments of *m*-aminophenol molecules. Different reactivity of isomeric aminophenols under interaction with CH₂O and H₂S is caused by a change of NH₂ group basicity and OH group acidity according to their arrangement in aromatic ring.

Herein we report on the results of our further investigations in the field of cyclothiomethylation of functionally substituted aromatic amines in order to develop novel procedures for a synthesis of new *N*-aryl substituted 1,3,5-dithiazinanes, 1,3,5-thiadiazinanes and macroheterocycles.


RESULTS AND DISCUSSION

CYCLOTHIOMETHYLATION OF AROMATIC AMINO ACIDS AND *p*-AMINOBENZOIC ACID ESTERS

In cyclocondensation with CH₂O and H₂S there have been involved aromatic amino acids and their derivatives namely *o*- (**1a**) and *p*-aminobenzoic **1b** acids, 4-(**1c**) and 5-aminosalicylic **1d** acids, and also *p*-aminobenzoic acid ethyl **1e** and (β -diethylamino)ethyl **1f** esters. Using *o*- (**1a**), *p*-aminobenzoic **1b** and 4- (**1c**), 5-aminosalicylic **1b** acids as an example we have studied the influence of COOH group position in aromatic ring on the activity of NH₂ and OH groups in the cyclothiomethylation reaction.

Thus, condensation of aminobenzoic acid **1a** with CH₂O and H₂S (1:3:2 ratio) at 20 °C led exclusively to *o*-(1,3,5-dithiazinan-5-yl)benzoic acid **2a** in 61% yield, whereas *p*-aminobenzoic acid **1b** gave a mixture of *p*-(1,3,5-dithiazinan-5-yl)- (**2b**) and *p*-(1,3-thiazetidin-3-yl)benzoic (**3b**) acids with good yields (51 and

34%, respectively) (Scheme 1). Dithiazinane **2b** has been selectively obtained at 60 °C in 95% yield.



Scheme 1 Cyclothiomeylation of *o*- and *p*-aminobenzoic acids, 4- and 5-aminosalicylic acids

4-Aminosalicylic acid **1c**, similar to acid **1a**, reacts with CH₂O and H₂S to form exclusively 4-(1,3,5-dithiazinan-5-yl)-2-hydroxybenzoic acid **2c** in 89% yield, and 5-aminosalicylic acid **1d**, similar to acid **1b**, gave a mixture of 5-(1,3,5-dithiazinan-5-yl)-2-hydroxybenzoic acid **2d** and 5-(1,3-thiazetidin-3-yl)-2-hydroxybenzoic acid **3d** in the yields of 32 and 22%, respectively.


4-Aminosalicylic acid **1c**, in which OH and NH₂ groups occupy *meta*- position, in contrast to *m*-aminophenol, reacts with CH₂O and H₂S exclusively at the amino group, probably, as a result of the intramolecular hydrogen bonding between OH and COOH groups (OH... O the *ortho*-effect).⁶ As is evident, the direction of cyclothiomeylation of isomeric aromatic amino acids with the aid of CH₂O and H₂S depends on the arrangement of functional groups in aromatic ring and their mutual influence.

Individual compounds **2a-d** and **3b,d** have been isolated by means of fractional crystallization. The structures of dithiazinanes **2a-d** and thiazetidines **3b,d** are proven by spectral methods: ¹H NMR and ¹³C NMR, GC/MS, and also by the data of element analysis. Molecular weight was determined by Rast cryoscopic method.⁷ Heterocycle **2a** was obtained earlier^{4a} with 81% yield by cyclocondensation of **1a** with NaHS and CH₂O, for which any spectral characteristics were absent.

Thus, aromatic amino acids were condensed with CH₂O and H₂S only at NH₂ group with the formation of *N,S*-containing heterocycles (1,3,5-dithiazinanes and 1,3-thiazetidines). One should notice that *para*-position of COOH group contributes to an increase of NH₂ group activity in the condensation reaction with CH₂O and H₂S. The greatest yield of target dithiazinanes **2a-d** was reached in temperature range from 20 to 60 °C. An increase in reaction temperature up to 80 °C led to a decrease in selectivity with formation of by-product of the reaction – 1,2,4-trithiolane.⁸


By the reaction of *p*-aminobenzoic acid ethyl **1e** or (β-diethylamino)ethyl **1f** esters with CH₂O and H₂S (1:3:2 ratio) at 20 °C the compounds of dithiazinane row **2e,f** has been obtained. Amino ester **1e** in this reaction together with ethyl-4-(1,3,5-dithiazinan-5-yl) benzoate **2e** gave ethyl-4-(1,3-thiazetidin-3-yl) benzoate **3e** and 3,5-di(4-ethylcarboxyphenyl)-1,3,5-thiadiazinane **4e** in 56, 18% and 15% yields, respectively (Scheme 2). At 60 and 80 °C the reaction resulted in a mixture of products **2e-4e**. The latter compounds were isolated by column chromatography.

The regioselective synthesis of thiadiazinane **4e** was carried out at 0 °C by cyclocondensation of *p*-aminobenzoic acid ethyl ester **1e** with CH₂O and H₂S (2:3:1 ratio) in 93% yield. Heterocyclic **4e** in the presence of three-molar excess of a thiomethylation mixture CH₂O-H₂S at ambient temperature was completely converted into the appropriate dithiazinane **2e**.



Scheme 2. Cyclothiomethylation of *p*-amino benzoic acid ethyl and (β -diethylamino)ethyl esters

According to X-ray diffraction analysis the molecule of **4e** has local plane of symmetry passing through C(2) and S(1) atoms (Figure 2a). The thiadiazinane ring has a chair conformation with axial *cis* position of (4-ethoxycarbonyl)phenylic substituent at the nitrogen atoms. In this orientation the lone pairs at the nitrogen atoms occupy *cis* equatorial position and are antiperiplanar to the lone pairs of the sulfur atom (conformer B).



The orbital repulsion of lone electron-pairs of the nitrogen and sulfur atoms stabilizes an axial *cis* position of (4-ethoxycarbonyl)phenylic substituent (anomeric effect).^{9a,b}




Figure. General view (a) and fragment of crystal packing (b) of compound **4e** in the crystal

In crystal structure above there are two types of intermolecular hydrogenous bonds (Figure 2b). The sulfur atom of one thiadiazinane molecule forms hydrogenous bonds with protons on the two aromatic


rings at C(4) and with methylene protons of another thiadiazinane molecule at C(2): HC(4)...S(1)...HC(4') и HC(2)...S(1) (where H...S distances are equal to 2.83 Å (x 2) and 2.78 Å, respectively). At the same time, the ester carbonyl groups of the first thiadiazinane molecule are connected with methylene protons of two other thiadiazinane molecules: O(1)...HC(1A) and O(1')...HC(1B) (distances O...H 2.54 Å (x 2)).

In the ^1H NMR spectra of dithiazinane cycles **2a-f** one could observe two singlets of methylene protons at δ_{H} 3.40-4.22 (SCH₂S) and 4.15-5.20 ppm (NCH₂S) with integrated intensities ratio of 1:2. The ^{13}C NMR spectra contain signals at δ_{C} 30.61-34.80 (SCH₂S) and 44.51-56.89 ppm (NCH₂S) belonging to the carbon atoms. The ^1H NMR spectrum of thiazetidine cycle (compounds **3b,d,e**) exhibits singlets of methylene protons (NCH₂S) at δ_{H} 4.30-5.15 ppm, while the ^{13}C NMR spectra contain peaks at δ_{C} 51.97-54.60 ppm. In the ^1H NMR spectra of thiadiazinane **4e** the signals for CH₂SCH₂ and NCH₂N methylene protons are observed at δ_{H} 4.94 and 5.32 ppm (2:1 ratio), while the ^{13}C NMR spectrum contains signals at δ_{C} 53.13 and 68.23 ppm assigned to the carbon atoms located between two nitrogen atoms and the sulfur and nitrogen atoms, respectively.

CYCLOMETHYLATION OF *p*-ANILINE SULFAMIDE AND ITS DERIVATIVES

We investigated cyclothiomethylation of *p*-aminosulfanyl acid amides, namely, *p*-aniline sulfamide **1g**, *p*-aminobenzene sulfacetamide **1h**, 2-(*p*-aminobenzenesulfamido)-3-methoxypyrazine and 4-(*p*-aminobenzenesulfamido)-2,6-dimethoxypyrimidine with the aid of CH₂O and H₂S.


Condensation of *p*-aniline sulfamide **1g** was carried out (**1g**: CH₂O: H₂S = 1:3:2) under different temperature conditions (0-80 °C). It was stated that cyclothiomethylation of *p*-aniline sulfamide **1g** within the temperature range from 0 to 40 °C gave rise to the dithiazinane cycle formation exclusively involving the amino group of the aromatic ring to obtain 4-(1,3,5-dithiazinane-5-yl)aniline sulfamide **2g** in 35-73% yield (Scheme 3, Table 1). The increase the temperature to 80 °C facilitates the involvement of less reactive SO₂NH₂ into the cyclocondensation reaction. So, in these experiments together with **2g** cyclodimer **4g**. The obtained mixture of compounds **2g** and **4g** (3:1 ratio) was divided by fractional crystallization (Scheme 3).



Scheme 3. Cyclothiomethylation of *p*-aniline sulfamide

Cryoscopic determinations⁷ with a value of 285 ± 10 correspond to the molecular weight (mass) of compound **2g**, and a value of 493 ± 10 corresponds to the molecular weight of macroheterocycle **4g**. The element analysis of compound **2g** confirms the molecular formula $C_9H_{12}N_2O_2S_3$ and the molecular formula $C_{16}H_{20}N_4O_4S_4$ for compound **4g** as well.

To obtain 1,3,5-dithiazinanes simultaneously involving of both amino groups of *p*-aniline sulfamide **1g** into the reaction with CH_2O and H_2S , we have increased a quantity of a thiomethylation mixture: **1g**: CH_2O : H_2S = 1:6:4 (0, 20, 40, and 80 °C). As a result, cyclocondensation proceeded via both NH_2 groups to give the four-component mixture, GC/MS spectrum of which contains peaks of molecular ions of the characteristic residual fragments with m/z 318 (**7g**), 334 (**6g**), 364 (**8g**) and 380 (**5g**), apparently corresponding to compounds with the (oxy)thiazethidine **6g,7g** and (oxy)dithiazinane **5g,8g** cycles.



Cyclocondensation of **1g** by a mixture of CH_2O and H_2S (**1g**: CH_2O : H_2S = 1:6:4) in acid medium (pH 1.3-1.5) at 40 °C led to the formation of 5-[4-(1,3,5-dithiazinane-5-sulfonyl)phenyl]-1,3,5-dithiazinane **5g** in 93% yield and with 100% selectivity (Scheme 3, Table).

Table. The influence of pH medium and initial reagent ratio on the yield and product content of *p*-aniline sulfamide **1g** cyclomethylation


Reagent ratio 1g : CH_2O : H_2S	pH	T, °C	Yield (%)		
			2g	4g	5g
1 : 3 : 2	7.3-7.4	0	35	-	-
	7.3-7.4	20	58	-	-
	7.3-7.4	40	73	-	-
	7.3-7.4	80	52	14	-
1 : 6 : 4	1.3-1.5	40	-	-	93

The 1H NMR spectrum of heterocycle **2g** exhibits proton signals of aromatic ring at δ_H 6.80-7.67 ppm. Singlet signals at δ_H 3.82 and 4.32 ppm correspond to methylene protons located between the sulfur atoms and atoms of N and S, respectively, with integrated intensity of 1:2. The ^{13}C NMR spectrum for **2g** contains signals of the carbon atoms in aromatic rings at δ_C 116.48, 127.12, 133.99, and 147.86 ppm. The signals at δ_C 32.96 and 53.30 ppm evidences the presence of the dithiazinane ring methylene groups in compound **2g**.

The ^1H NMR spectrum of heterocycle **4g** exhibits proton signals of aromatic ring at δ_{H} 6.73–7.11 ppm. The singlet signals, which correspond to the methylene protons between the nitrogen and sulfur atoms, appeared at 3.98 and 4.52 ppm at a ratio of 1:1. The proton signals at δ_{H} 7.54 and 7.63 ppm belong to NH group. The ^{13}C NMR spectrum of compound **4g** contains the carbon atom signals of aromatic rings at δ_{C} 112.73, 127.35, 131.91, and 149.39 ppm. The presence of signals at δ_{C} 44.74 and 63.49 ppm for **4g** characterizes the presence of CH_2 groups between N and S atoms.

In ^1H NMR spectrum of compound **5g** the protons of aromatic ring are observed at δ_{H} 6.73–8.02, the methylene protons of SCH_2S and SCH_2N groups of the dithiazinane ring, linked with aromatic ring, are observed at δ_{H} 3.93 and 4.72 ppm (1:2 ratio), and of sulfo group at δ_{H} 4.26 and 5.14 ppm, respectively. The ^{13}C NMR spectrum of compound **5g** exhibits the signals of dithiazinane cycle carbon atoms, bound with aromatic ring, at δ_{C} 31.48 and 56.05 ppm, and with sulfo group at δ_{C} 33.10 and 63.40 ppm.

Cyclothiomethylation of *p*-aniline sulfacetamide **1h** was found to proceed simultaneously via NH_2 and $\text{SO}_2(\text{Ac})\text{NH}$ groups in acid medium (*pH* 2,5). Under these conditions cyclodimer **4h** has been obtained in 50% yield as a result of intermolecular condensation of two molecules of **1h** with a thiomethylation reagent $\text{CH}_2\text{O}-\text{H}_2\text{S}$ (Scheme 5) with 70% conversion. In neutral and alkaline medium, *p*-aniline sulfacetamide **1h** did not react with CH_2O and H_2S and 1,2,4-trithiolane was predominantly formed.⁸



Scheme 4. Cyclothiomethylation of *p*-aniline sulfo acetamide

The molecular weight of 543.53 ± 10 (calc. M_{cr} 544) determined by Rast cryoscopy method⁷ and the data of element analysis correspond to molecular formula $\text{C}_{20}\text{H}_{24}\text{N}_4\text{O}_6\text{S}_4$ that proved the formation of macroheterocycle **4h**. The ^1H NMR spectrum shows signals of methylene sulfide protons connected with amino group at δ_{H} 4.50 ppm, and connected with the sulfo acetamide group at δ_{H} 5.11 ppm. The ^{13}C NMR spectrum contains signals at δ_{C} 43.61 and 44.28 ppm assigned to carbon atoms of CH_2NH and CH_2NSO groups, respectively.

Cyclocondensation of [(*p*-aniline)sulfamido]-3-methoxypyrazine and [4-(*p*-aniline)sulfamido]-2,6-dimethoxypyrimidine containing pyrimidine rings under conditions described above does not proceed with CH_2O and H_2S . The low activity of NH_2 group in compounds **1i** and **1k** is, apparently, connected with the formation of the intermolecular hydrogenous bonds between NH_2 group and the nitrogen atoms of pyrimidine ring.

In conclusion, cyclocondensation of *o*- (**1a**), *p*- (**1b**) aminobenzoic, 4- (**1c**), 5- (**1d**) aminosalicyclic acids, ethyl- (**1e**), (β -diethylamino)ethyl esters- (**1f**) of *p*- (**1b**) and *p*-aniline sulfamide **1g** with CH₂O and H₂S under optimized conditions leads to the formation of the corresponding dithiazinanes **2a-g**, whereas *p*-aminobenzoic **1b**, 5-aminosalicylic acid **1d** and *p*-aminobenzoic acid ethyl ester **1e** together with dithiazinanes give rise to thiazetidines. Cyclothiomethylation of *p*-aminobenzoic acid ethyl ester by the CH₂O-H₂S reagent at 0 °C was found to afford *N,N*-diaryl-1,3,5-thiadiazinane **4e** in quantitative yield (93%). The reaction for obtaining *bis*-1,3,5-dithiazinanes **5g** (93%) from *p*-aniline sulfamide **1g** are more effective in acid medium (pH 1.3-1.5) at 40 °C. It was shown that in the cyclothiomethylation at 80 °C together with the formation of dithiazinane (56%) *p*-aniline sulfamide **1g** undergoes intermolecular condensation to give cyclodimer **4g** (14%). The analogous intermolecular cyclocondensation of *p*-aniline sulfacetamide in acid medium (pH 2,5) leads to cyclodimer **4h** (50%) built from two fragments of *p*-aniline sulfacetamide linked by *sym* dimethyl sulfide chain (CH₂SCH₂).

EXPERIMENTAL

All solvents were freshly distilled. The ¹H NMR spectra of compounds **2a-g**, **3b,d,e** and **4g**, **5g** were measured on spectrometer "Tesla BS-487", ¹³C NMR - on spectrometer Jeol FX 90Q (22.50 MHz), internal standard - Me₄Si. NMR experiments of compounds **4e** and **4h** were recorded on a Bruker AVANCE-400 spectrometer. The GLC-mass spectrometry was carried out on Finigan 4021 instrument. The IR-spectra were recorded on Specord 75 IR in spectrophotometer in Nujol mulls. Elemental analysis of C, H, N, S samples was determined on element analyser of Karlo Erba, model 1106. The pH values of solutions were determined on a pH meter (pH – 340). Melting points were determined on Kofler unit. Column chromatography was performed with the use of silica gel.

Single-crystal X-ray diffraction experiments were carried out with a Bruker SMART 1000 CCD area detector, using graphite monochromated Mo-K α radiation at 100 K. All calculations were performed on an IBM PC/AT using the SHELXTL software [G. M. Sheldrick, SHELXTL-97, Version 5.10, Bruker AXS Inc., Madison, WI-53719, USA]. Atomic coordinates, bond lengths, bond angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre (CCDC), deposition numbers 679451 (**4e**).

Cyclothiomethylation of functional substituted anilines (1a-1h). The calculated amount of 37% formalin (1.1 mL, 0.015 mol) or (2.2 mL, 0.030 mol) were charged to a three-neck flask equipped with a stirrer and barbotager thermostated at the chosen temperature. Hydrogen sulfide (prepared in excess amount from Na₂S and HCl) was barbotaged to give CH₂O-H₂S mixture at a ratio of 3:2 or 6:4. Then calculated amount of anilines (**1a**, **1b**, **1c**, **1d**, **1e**, **1f**, **1g** or **1h**; 0.005 mol) was added to the reaction mixture. The mixture was stirred for 2-3 h at a chosen temperature. Compounds **2e**, **3e**, **4e** were isolated

by column chromatography with C₆H₆/EtOAc (10:1) as the eluent. The products **2a–d**, **3b**, **3d**, **2g**, **4g** were selected by fractional crystallization from CHCl₃. Compounds **5g** and **4h** were filtered.

7-(1,3,5-Dithiazinane-5-yl)-2-benzoic acid (2a). White powder (61%). mp 159-160 °C. *m/z* (%): 241 (5) [*M*]⁺, 197 (47), 163 (5), 150 (30), 120 (53), 105 (5), 92 (44), 77 (12), 61 (100). ¹H NMR (80 MHz, DMSO-*d*₆): δ (ppm) 3.75 (s, 2H, 2H-2), 4.66 (s, 4H, 2H-4, 2H-6), 6.64-7.00 (m, 2H, H-10, H-12), 7.28-7.53 (m, 2H, H-9, H-11), 8.00 (s, H, H-15). ¹³C NMR (22.5 MHz, DMSO-*d*₆): δ (ppm) 32.2 (C-2), 44.5 (C-6, C-4), 109.9 (C-8), 114.9 (C-10), 116.6 (C-12), 131.5 (C-9), 134.0 (C-11), 151.7 (C-7), 169.9 (C-13). IR (KBr): 750, 1230, 1600, 1660, 2900, 3320 cm⁻¹. *Anal.* Calcd for C₁₀H₁₁O₂NS₂: C 49.79, H 4.56, N 5.80, S 26.55. Found: C 50.11, H 4.60, N 5.79, S 26.46.

7-(1,3,5-Dithiazinane-5-yl)-4-benzoic acid (2b). White powder (95%). mp 227-229 °C. *m/z* (%): 241 (9) [*M*]⁺, 207 (12), 91 (15); 44 (100). ¹H NMR (80 MHz, DMSO-*d*₆): δ (ppm) 4.00 (d, *J*=8.0 Hz, 2H, 2H-2), 4.64 (br. s, 4H, 2H-4, 2H-6), 6.72 (d, *J*=8.0 Hz, 2H, H-8, H-12), 7.17 (d, *J*=8.0 Hz, 2H, H-9, H-11), 7.83 (s, H, H-13). ¹³C NMR (22.5 MHz, DMSO-*d*₆): δ (ppm) 33.7 (C-2), 56.3 (C-6, C-4), 113.8 (C-8, C-12), 121.0 (C-10), 131.0 (C-9, C-11), 149.1 (C-7), 167.7 (C-13). IR (KBr): 780, 1200, 1600, 1680, 2900, 3300 cm⁻¹. *Anal.* Calcd for C₁₀H₁₁O₂NS₂: C 49.79, H 4.56, N 5.80, S 26.55. Found: C 49.79, H 4.56, N 5.80, S 26.63.

5-(1,3-Thiazetidine-3-yl)-4-benzoic acid (3b). White powder (51%). mp 209-210 °C. *m/z* (%): 195 (52) [*M*]⁺, 150 (33); 120 (57); 92 (48); 61 (100). ¹H NMR (80 MHz, DMSO-*d*₆): δ (ppm) 4.36 (br. s, 4 H, 2H-2, 2H-4), 6.72 (d, *J*=8.0 Hz, 2 H, H-6, H-10), 7.17 (d, *J*=8.0 Hz, 2 H, H-7, H-9), 7.75 (s, H, H-11). ¹³C NMR (22.5 MHz, DMSO-*d*₆): δ (ppm) 53.3 (C-2, C-4), 116.4 (C-6, C-10), 120.0 (C-8), 131.0 (C-7, C-9), 147.2 (C-5), 167.7 (C-11). IR (KBr): 780, 1200, 1600, 3300 cm⁻¹. *Anal.* Calcd for C₉H₉O₂NS: C 55.38, H 4.61, N 7.17, S 16.41. Found: C 55.53, H 4.59, N 7.29, S 16.92.

4-(1,3,5-Dithiazinane-5-yl)-2-hydroxybenzoic acid (2c). White powder (89%). mp 258-260 °C. ¹H NMR (80 MHz, DMSO-*d*₆): δ (ppm) 4.06 (s, 2 H, 2H-2), 5.20 (s, 4 H, 2H-4 and 2H-6), 6.23 (s, H, H-12), 6.70 (s, H, H-8), 7.85 (s, H, H-11), 8.50 (br. s, 2 H, H-13, H-14). ¹³C NMR (22.5 MHz, DMSO-*d*₆): δ (ppm) 33.6 (C-2), 53.2 (C-4, C-6), 103.7 (C-10), 106.4 (C-8), 108.4 (C-12), 131.4 (C-11), 151.5 (C-7), 163.1 (C-9), 172.3 (C-13). IR (KBr): 720, 1170, 1450, 1600, 2900, 3360 cm⁻¹. *Anal.* Calcd for C₁₀H₁₁O₃NS₂: C 46.69, H 4.28, N 5.44, S 24.90. Found: C 47.21, H 4.33, N 5.48, S 25.16.

5-(1,3,5-Dithiazinane-5-yl)-2-hydroxybenzoic acid (2d). White powder (32%). mp 194-196 °C. ¹H NMR (80 MHz, DMSO-*d*₆): δ (ppm) 4.00 (s, 2 H, 2H-2), 5.20 (s, 4 H, 2H-4 and 2H-6), 6.80-7.50 (m, 3 H, H-8, H-11 and H-12), 8.45 (br. s, 2 H, H-13, H-14). ¹³C NMR (22.5 MHz, DMSO-*d*₆): δ (ppm) 33.5 (C-2), 56.9 (C-4, C-6), 113.0 (C-9), 114.7 (C-11), 117.9 (C-8), 123.0 (C-12), 137.7 (C-7), 155.0 (C-10), 171.8 (C-14). IR (KBr): 800, 1190, 1440, 1600, 1660, 2900 cm⁻¹. *Anal.* Calcd for C₁₀H₁₁O₃NS₂: C 46.69, H 4.28, N 5.44, S 24.90. Found: C 46.73, H 4.27, N 5.54, S 25.00.

5-(1,3-Thiazetidine-3-yl)-2-hydroxybenzoic acid (3d). White powder (22%). mp 164-166 °C. ^1H NMR (80 MHz, DMSO- d_6): δ (ppm) 4.35 (s, 4 H, 2H-2, 2H-4), 6.80-7.50 (m, 3 H, H-6, H-9, H-10) 8.47 (br. s, 2 H, H-11, H-12). ^{13}C NMR (22.5 MHz, DMSO- d_6): δ (ppm) 54.5 (C-2, C-4), 113.0 (C-7), 115.1 (C-9), 119.2 (C-6), 126.3 (C-10), 138.1 (C-5), 158.8 (C-8), 172.2 (C-12). IR (KBr): 800, 1190, 1440, 1600, 1660, 2900 cm^{-1} . *Anal.* Calcd for $\text{C}_9\text{H}_9\text{O}_3\text{NS}$: C 51.17, H 4.29, N 6.63, S 15.18. Found: C 51.53, H 4.36, N 6.42, S 15.03.

Ethyl-4-(1,3,5-dithiazinan-5-yl)benzoate (2e). White powder (56%). mp 136-138 °C. m/z (%) = 269 (40) $[M]^+$, 191 (21), 177 (70), 163 (24), 149 (33), 132 (100), 77 (36) 45 (52). ^1H NMR (80 MHz, CDCl_3): δ (ppm) 1.38 (t, $J=7.2$ Hz, 3 H, H_3C -16), 3.92 (s, 2 H, 2H-2), 4.30 (s, 4 H, 2H-4, 2H-6), 5.10 (br. s, 2 H, 2H-15), 7.05 (d, $J=9.0$ Hz, 2 H, H-8, H-12), 8.05 (d, $J=9.0$ Hz, 2 H, H-9, H-11). ^{13}C NMR (22.5 MHz, CDCl_3): δ (ppm) 14.3 (C-16), 34.8 (C-2), 54.5 (C-4, C-6), 60.4 (C-15), 116.2 (C-8, C-12), 121.9 (C-10), 131.0 (C-9, C-11), 148.5 (C-7), 164.6 (C-13). IR (KBr): 750, 1230, 1450, 1660, 3320 cm^{-1} . *Anal.* Calcd for $\text{C}_{12}\text{H}_{15}\text{O}_2\text{NS}_2$: C 53.53, H 5.58, N 5.20, S 23.79. Found: C 54.19, H 5.48, N 5.11, S 23.77.

Ethyl-4-(1,3-thiazetidin-3-yl)benzoate (3e). White powder (18%). mp 175-177 °C. m/z (%) = 225 (0.4) 223 (9) $[M]^+$, 177 (9), 150 (15), 149 (100), 132 (15), 44 (42), 46 (1.7). ^1H NMR (80 MHz, CDCl_3): δ (ppm) 1.34 (t, $J=7.2$ Hz, 3H, H_3C -14), 4.25 (k, $J=9.0$ Hz, 2 H, H_2C -13), 5.10 (br. s, 4H, 2H-2, 2H-4), 6.98 (d, $J=6.3$ Hz, 2 H, H-6, H-10), 7.68 (d, $J=6.3$ Hz, 2 H, H-7, H-9). ^{13}C NMR (22.5 MHz, CDCl_3): δ (ppm) 14.4 (C-14), 53.2 (C-2, C-4), 60.6 (C-13), 116.4 (C-10, C-6), 125.5 (C-8), 131.2 (C-7, C-9), 155.0 (C-5), 165.6 (C-11). IR (KBr): 750, 1230, 1450, 1660, 3320. *Anal.* Calcd for $\text{C}_{11}\text{H}_{13}\text{O}_2\text{NS}$: C 59.19, H 5.83, N 6.28, S 14.35. Found: C 57.20, H 5.42, N 6.11, S 15.77.

3,5-Di-(4-ethylcarboxyphenyl)-1,3,5-thiadiazinane (4e). Compound **4e** was isolated by column chromatography (yield 15%). And also the compound **4e** has selectively been received at 0 °C, the starting reagents were taken in the ratio **1e**: CH_2O : H_2S = 2:3:1 (yield 93%). White powder, mp 184-186 °C. ^1H NMR (400 MHz, CDCl_3): δ (ppm) 1.32 (t, $J=6.5$ Hz, 6 H, H_3C -22, H_3C -26), 4.24 (d, $J=6.8$ Hz, 4 H, 2H-21, 2H-25), 4.94 (s, 4 H, 2H-2, 2H-6), 5.32 (s, 2 H, 2H-4), 6.91 (d, $J=8.1$ Hz, 4 H, H-8, H-12, H-14, H-18), 7.79 (d, $J=8.1$ Hz, 4 H, H-9, H-11, H-15, H-17). ^{13}C NMR (100 MHz, CDCl_3): δ (ppm) 14.4 (C-22, C-26), 53.1 (C-2, C-6), 60.57 (C-25, C-21), 68.2 (C-4), 116.3 (C-12, C-8, C-18, C14), 122.1 (C-16, C-10), 131.2 (C-11, C-9, C-17, C-15), 150.5 (C-7, C-13), 166.4 (C-23, C-19). IR (KBr): 750, 1230, 1450, 1660, 3320. *Anal.* Calcd for $\text{C}_{11}\text{H}_{13}\text{O}_2\text{NS}$: C 62.98, H 5.04, N 6.99, S 8.01. Found: C 63.25, H 5.32, N 7.11, S 8.65.

4-(1,3,5-Dithiazinane-5-yl)-2-(diethylamino)ethylbenzoate (2f). White powder (79%). mp 51-52 °C. ^1H NMR (80 MHz, CDCl_3): δ (ppm) 0.45 (t, $J=6.3$ Hz, 6 H, H_3C -19, H_3C -22), 1.30 (t, $J=9.0\text{Hz}$, 2 H, 2H-16), 3.12 (k, $J=10.1$ Hz, 4 H, 2H-18, 2H-21), 3.80 (t, $J=9.0$ Hz, 2 H, 2H-15), 4.22 (s, 2 H, 2H-2), 4.68 (s, 4 H, 2H-4, 2H-6), 7.3 (br. s, 2H, H-8, H-12), 7.87 (s, 2 H, H-9, H-11). ^{13}C NMR (22.5 MHz, CDCl_3): δ (ppm)

8.5 (C-19, C-22), 32.4 (C-2), 47.1 (C-18, C-21), 49.5 (C-4, C-6), 53.8 (C-16), 58.2 (C-15), 116.9 (C-8, C-12), 131.1 (C-9, C-11), 150.5 (C-7), 164.5 (C-13). IR (KBr): 770, 1100, 1380-1450, 1600, 1680, 2900 cm^{-1} . *Anal.* Calcd for $\text{C}_{16}\text{H}_{24}\text{O}_2\text{N}_2\text{S}_2$: C 56.47, H 8.24, N 8.24, S 18.82. Found: C 51.97, H 7.46, N 7.70, S 19.50.

4-(1,3,5-Dithiazinane-5-yl)benzenesulfonamid (2g). White powder (73%). mp 135-137 °C. ^1H NMR (80 MHz, $\text{DMSO}-d_6$): δ (ppm) 3.90 (s, 2 H, 2H-2), 4.56 (s, 4 H, 2H-4, 2H-6), 5.16 (s, 2 H, 2H-14), 6.80 (d, $J=8.3$ Hz, 2H, H-8, H-12), 7.67 (d, $J=8.3$ Hz, 2 H, H-9, H-11). ^{13}C NMR (22.5 MHz, $\text{DMSO}-d_6$): δ (ppm) 33.0 (C-2), 53.3 (C-4, C-6), 116.5 (C-8, C-12), 127.1 (C-9, C-11), 134.0 (C-7), 147.9 (C-10). IR (KBr): 685, 820, 1090, 1140, 1305, 1450, 1600, 2905, 3365 cm^{-1} . *Anal.* Calcd for $\text{C}_9\text{H}_{12}\text{N}_2\text{S}_3\text{O}_2$: C 39.11, H 4.38, N 10.14, S 34.80. Found: C 39.47, H 4.84, N 11.84, S 35.67.

2,2,12,12-Tetraon-2 λ^6 ,5,12 λ^6 ,15-tetrathia-3,7,13,17-tetraazatricyclo[16.2.2.2 8,11]-tetracos-1(20),8,10,18,21,23-hexaene (4g). White powder (14%). mp 146-148 °C. ^1H NMR (80 MHz, $\text{DMSO}-d_6$): δ (ppm) 3.98 (s, 4 H, 2H-6, 2H-16), 4.52 (s, 4 H, 2H-4, 2H-14), 6.73 (d, $J=8.44$ Hz, 4 H, H-9, H-22, H-19, H-23), 7.11 (d, $J=8.44$ Hz, 4 H, H-10, H-20, H-22, H-24), 7.54 (m, 2 H, NH-7, NH-17), 7.63 (s, 2 H, NH-3, NH-13). ^{13}C NMR (22.5 MHz, $\text{DMSO}-d_6$): δ (ppm) 44.7 (C-6, C-16), 63.5 (C-4, 14), 112.7 (C-9, C-19, C-21, C-24), 127.4 (C-10, C-20, C-22, C-23), 131.9 (C-8, C-18), 149.4 (C-1, C-11). IR (KBr): 560, 815, 1095, 1150, 1305, 1460, 1600, 2910, 3370 cm^{-1} . *Anal.* Calcd for $\text{C}_{16}\text{H}_{20}\text{N}_4\text{S}_4\text{O}_4$: C 41.72, H 4.38, N 12.16, S 27.85. Found: C 39.35, H 4.54, N 11.82, S 33.73.

5-[4-(1,3,5-Dithiazinan-5-ylsulfonyl)phenyl]-1,3,5-dithiazinane (5g). Analogously to the above-described procedure compound (5g) was prepared with accompaniment of the 0.01 mol HCl. After 2 h the reaction mixture was neutralized with an aqueous KOH solution. White powder (93%). mp 154-156 °C. ^1H NMR (80 MHz, $\text{DMSO}-d_6$): δ (ppm) 3.93 (s, 2 H, 2H-2), 4.26 (s, 2 H, 2H-17), 4.72 (s, 4 H, 2H-4, 2H-6), 5.14 (s, 4 H, 2H-15, 2H-19), 6.73 (d, $J=7.98$, 2 H, H-8, H-12), 8.02 (d, $J=7.98$, 2 H, H-9, H-11). ^{13}C NMR (22.5 MHz, $\text{DMSO}-d_6$): δ (ppm) 31.5 (C-2), 33.1 (C-17), 56.1 (C-4, C-6), 63.4 (C-15, C-19), 114.0 (C-8, C-12), 127.0 (C-9, C-11), 128.0 (C-10), 146.6 (C-7). IR (KBr): 645-695, 1005, 1090, 1140, 1300, 1445, 1595, 2900 cm^{-1} . *Anal.* Calcd for $\text{C}_{12}\text{H}_{16}\text{N}_2\text{S}_5\text{O}_2$: C 37.87, H 4.24, N 7.36, S 42.13. Found: C 37.71, H 4.53, N 8.18, S 43.91.

5-[4-(1,3-Thiazetan-3-ylsulfonyl)phenyl]-1,3,5-dithiazinane (6g). White powder (20%). m/z (%): 334 (5) $[M]^+$, 156 (15), 108 (35), 43 (50). IR (KBr): 550, 1005, 1090, 1140, 1300, 1445, 1595, 2900 cm^{-1} .

5-[4-(1,3-Oxazetan-3-ylsulfonyl)phenyl]-1,3,5-dithiazinane (7g). White powder (20%). m/z (%): 318 (5) $[M]^+$, 156 (12), 108 (40), 76 (24), 43 (95). IR (KBr): 550, 1005, 1140, 1300, 1445, 1595, 2900 cm^{-1} .

5-[4-(1,3,5-Oxathiazinan-5-ylsulfonyl)phenyl]-1,3,5-dithiazinane (8g). White powder (20%). m/z (%): 364 (2) $[M]^+$, 200 (20), 136 (60), 78 (30), 43 (100). IR (KBr): 550, 810, 1090, 1300, 1445, 2900 cm^{-1} .

2,2,12,12-Tetraon-2 λ^6 ,5,12 λ^6 ,15-tetrathia-(3,13-diacyl)-3,7,13,17-tetraasatricyclo[16.2.2.2 8,11]-

tetracos-1(20),8,10,18,21,23-hexaene (4h). Analogously to the above-described procedure compound (4h) was prepared with accompaniment of the 0.01 mol HCl. After 3h the reaction mixture was neutralized with an aqueous KOH solution. White powder (51%). mp 126-127 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) 1.88 (s, 6 H, H₃C-26, H₃C-28), 4.5 (s, 4 H, 2H-6, 2H-16); 5.11 (s, 4 H, 2H-4, 2H-14); 6.73 (d, *J* = 6.85, 4 H, H-9, H-23, H-19, H-22); 7.63 (d, *J* = 6.85, 4 H, H-10, H-24, H-20, H-21); 21 (m, 2 H, NH-7, NH-17). ¹³C NMR (100 MHz, DMSO-*d*₆): δ (ppm) 23.4 (C-26, C-28), 43.6 (C-6, C-16); 44.3 (C-4, C-14); 116.3 (C-9, C-19, C-22, C-24); 129.6 (C-20, C-21, C-10, C-24), 147.4 (C-1, C-11), 153.8 (C-8, C-18), 169.0 (C-25, C-27). IR (KBr): 770, 1100, 1400-1450, 1570, 1620, 2900 cm⁻¹. *Anal.* Calcd for C₂₀H₂₄N₄S₄O₆: C 44.10, H 4.48, N 10.09, S 23.55. Found: C 42.78, H 5.07, N 9.88, S 23.65.

REFERENCES

1. a) A. Wohl, *Bericht*, 1886, **19**, 2344. b) E. R. Braithwaite and J. Graymore, *J. Chem. Soc.*, 1953, **1**, 143. c) D. Collins and J. Graymore, *J. Chem. Soc.*, 1953, 4089. d) D. Collins and J. Graymore, *J. Chem. Soc.*, 1958, 2893. e) E. R. Braithwaite and J. Graymore, *J. Chem. Soc.*, 1950, **2**, 208.
2. a) Brit. Patent, 943.273, 1963 (*Chem. Abstr.*, 1964, **60**, 5528a). b) N. Nohyaku, *Chem. Abstr.*, 1985, **102**, 149292d. c) T. Shibamoto, *J. Agr. Food Chem.*, 1980, **28**, 237. d) H. W. Brinkman, H. Copier, Joop J. M. de Leuw and S. B. Tjan, *J. Agr. Food Chem.*, 1972, **20**, 177. e) M. Boelens, L. M. van der Linde, P. J. de Valois, H. M. van Dort and H. J. Takken, *J. Agr. Food Chem.*, 1974, **22**, 1071. f) T. Shibamoto and G. F. Russell, *J. Agr. Food Chem.*, 1976, **24**, 4, 843. g) G. MacLeod and B. M. Coppock, *J. Agr. Food Chem.*, 1977, **25**, 113. h) H. Paulsen and F. R. Heiker, *Liebigs. Ann. Chem.* 1981, 2180.
3. a) J. C. Galves-Ruiz, H. Nöth and A. Flores-Parra, *Inorg. Chem.*, 2003, **42**, 7569. b) C. J. Galvez-Ruiz, C. Guadarrama-Perez, H. Nöth and A. Flores-Parra, *Eur. J. Inorg. Chem.*, 2004, 601. c) A. Flores-Parra, S. A. Sanchez-Ruiz, C. Guadarrama, H. Nöth and R. Contreras, *Eur. J. Inorg. Chem.*, 1999, 2069. d) C. Guadarrama-Perez, G. Cadenas-Pliego and A. Flores-Parra, *Chem. Ber.*, 1997, **130**, 813. e) L. M. Martinez-Aguilera, G. Cadenas-Pliego, R. Contreras and A. Flores-Parra, *Tetrahedron: Asymmetry*, 1995, **6**, 1585. f) A. Flores-Parra, G. Cadenas-Pliego, L. M. Martinez-Aguilera, M. L. Garcia-Nares and R. Contreras, *Chem. Ber.*, 1993, **126**, 863. g) A. Flores-Parra, N. Farfan, A. I. Hernandez-Bautista, L. Fernandez-Sanchez and R. Contreras, *Tetrahedron*, 1991, **47**, 6903.
4. French Patent, 1.341.792, 1963 (*Chem. Abstr.*, 1964, **60**, 5528d).
5. a) S. R. Khafizova, V. R. Akhmetova, L. F. Korzhova, T. V. Tyumkina, G. R. Nadyrgulova, R. V. Kunakova, E. A. Kruglov and U. M. Dzhemilev, *Russ. Chem. Bull.*, 2005, **2**, 432. b) S. R. Khafizova,

- V. R. Akhmetova, R. V. Kunakova and U. M. Dzhemilev, *Russ. Chem. Bull.*, 2003, **8**, 1817. c) S. R. Khafizova, V. R. Akhmetova, G. R. Nadyrgulova, I. V. Rusakov, R. V. Kunakova and U. M. Dzhemilev, *Petrochemistry*, 2005, **5**, 345. d) V. R. Akhmetova, G. R. Nadyrgulova, T. V. Tyumkina, Z. A. Starikova, M. Yu. Antipin, R. V. Kunakova and U. M. Dzhemilev, *Russ. J. Org. Chem.*, 2007, **41**, 151. e) V. R. Akhmetova, G. R. Nadyrgulova, S. R. Khafizova, T. V. Tyumkina, A. A. Yakovenko, M. Yu. Antipin, L. M. Khalilov, R. V. Kunakova and U. M. Dzhemilev, *Russ. Chem. Bull.*, 2006, **2**, 312.
6. R. D. Lindsay in *Comprehensive Org. Chem.*, Moscow, Chemistry, 1982, **3**, 168 [S. D. Barton and W. D. Ollis, *Comprehensive Org. Chem.*, Pergamon Press, Oxford-N. Y.-Toronto, 1979].
7. C. E. Bennett, *Am. Laboratory*, 1971, **3**, 67.
8. R. S. Aleev, Yu. S. Dalnova, Yu. N. Popov, R. M. Masagutov and S. R. Rafikov, *Dokl. Acad. Nauk SSSR*, 1988, **303**, 873.
9. a) L. Angiolini, R. P. Duke, R. A. Y. Jones and A. R. Katritzky, *J. Chem. Soc., Perkin 2*, 1972, 674.
b) Yu. Yu. Samitov, *Chem. Heter. Comp.*, 1980, **11**, 1443.