

HETEROCYCLES, Vol. 78, No. 1, 2009, pp. 161 – 168. © The Japan Institute of Heterocyclic Chemistry
Received, 31st July, 2008, Accepted, 12th September, 2008, Published online, 18th September, 2008.
DOI: 10.3987/COM-08-11509

ONE-POT SYNTHESIS OF 1-ARYLINDOLE-3-CARBOXYLATES FROM 2-(2-ISOCYANOPHENYL)ACETATES

Shuhei Fukamachi, Hisatoshi Konishi, and Kazuhiro Kobayashi*

Division of Applied Chemistry, Department of Chemistry and Biotechnology,
Graduate School of Engineering, Tottori University, 4-101 Koyama-minami,
Tottori 680-8552, Japan

Abstract - An efficient method for the preparation of 1-arylindole-3-carboxylates under mild conditions has been developed. Thus, cyclization of ethyl 2-(2-isocyanophenyl)acetates using sodium hydride at room temperature was followed by 1-arylation of the resulting ethyl 1-sodioindole-3-carboxylates with fluoro(di)nitrobenzenes to give ethyl 1-(di)nitrophenylindole-3-carboxylates in generally good yields. This method also allows preparation of 1,1'-(4,6-dinitro-1,3-phenylene)diindole derivatives by using 1,3-difluoro-4,6-dinitrobenzene.


INTRODUCTION

2-(Lithiomethyl)phenyl isocyanides have been used as intermediates in syntheses of a number of indole derivatives and related heterocycles by reactions with various electrophiles followed by cyclization.¹ Methyl 2-(2-isocyanophenyl)acetate, which could be prepared by reacting 2-(lithiomethyl)phenyl isocyanide with methyl chloroformate,² was elaborated to the preparation of methyl indole-3-carboxylate and methyl 3-alkyl-3*H*-indole-3-carboxylates by copper(I) oxide-catalyzed cyclization.³ However, any other elaborations of 2-(2-isocyanophenyl)acetates have not been reported so far. In this report, we describe a facile one-pot procedure for the synthesis of ethyl 1-arylindole-3-carboxylates by cyclization of ethyl 2-(2-isocyanophenyl)acetates followed by 1-arylation with fluoro(di)nitrobenzenes under mild conditions. Indole-3-carboxylates are synthetically as well as medicinally important heterocyclic compounds,⁴ and so, several methods for their preparation have been recently reported.⁵ However, only a few methods have been reported for the synthesis of 1-aryl derivatives involving construction of the pyrrole moiety.⁶ Most of syntheses of 1-arylindole-3-carboxylates have been based on the copper-catalyzed coupling reactions between indole-3-carboxylates and halobenzenes at higher temperature.⁷

RESULTS AND DISCUSSION

The one-pot conversion of ethyl 2-(2-isocyanophenyl)acetates (**2**), which were easily prepared from

2-methylphenyl isocyanides (**1**) on a successive treatment with lithium diisopropylamide (LDA) and ethyl chloroformate according to the procedure reported by Ito et al,^{1,2} into ethyl 1-(2-[or 4-]nitro- or 2,4-dinitrophenyl)indole-3-carboxylates (**5**) was conducted as illustrated in Scheme 1. Thus, compounds (**2**) were treated with sodium hydride in dimethyl sulfoxide (DMSO)–THF at room temperature. The cyclization proceeded slowly (for about 3 h) to generate the corresponding ethyl 1-sodioindole-3-carboxylates (**3**). Subsequent 2-(or 4-)nitro- or 2,4-dinitro-phenylation was achieved by adding 1-fluoro-2-(or 4-)nitro- or 1-fluoro-2,4-dinitro-benzenes (**4**) to give, after usual workup followed by purification by preparative TLC on silica gel, the desired 1-arylindole-3-carboxylates (**5**). The results are summarized in Table 1. From these results it can be seen that the 1-arylation reactions generally proceeded very smoothly (within 30 min) and cleanly to give the desired products (**5**) in fair to good yields.⁸ When the sequence was carried out using 1-fluoro-2,4-dinitrobenzene, the dinitroarylation completed immediately (within 5 min) to give the corresponding ethyl 1-(2,4-dinitrophenyl)indole-3-carboxylates (**5d, g, i, and j**) in good yields (Entries 4, 7, 9, and 10).⁹




Scheme 1

Table 1. Preparation of 1-Arylindole-3-carboxylates (**5**)

Entry	2	4	Time/min	5 (Yield/%) ^a
1	2a	4a (R ² = NO ₂ , R ³ = H)	10	5a (84)
2	2a	4b (R ² = NO ₂ , R ³ = Me)	10	5b (82)
3	2a	4c (R ² = H, R ³ = NO ₂)	30	5c (80)
4	2a	4d (R ² = R ³ = NO ₂)	<5	5d (91)
5	2b	4a	10	5e (76)
6	2b	4c	30	5f (76)
7	2b	4d	<5	5g (84)
8	2c	4a	10	5h (86)
9	2c	4d	<5	5i (90)
10	2d	4d	<5	5j (91)

^aYields are of purified products.

We next examined the possibility of the synthesis of 1,1'-(4,6-dinitro-1,3-phenylene)diindole derivatives **7** as an application of the present cyclization-arylation sequence. As shown in Scheme 2, when 1-sodioindole-3-carboxylates (**3**) were treated with 1,3-difluoro-4,6-dinitrobenzene (**6**) (2:1 molar ratio) under the same conditions as described for the preparation of 1-arylindole-3-carboxylates (**5**), the arylation reaction also proceeded very smoothly to lead to the formation of the corresponding desired products (**7**) in good-to-excellent yields. To the best of our knowledge, this type of (phenylene)diindoles are unknown in the literature.



Scheme 2

Unfortunately, however, it should be noted that the use of 2-fluorobenzonitrile instead of fluoro(di)nitrobenzenes in the 1-arylation step using **2a** resulted in the formation of an intractable mixture of products, from which no more than a trace amount of the desired ethyl 1-(2-cyanophenyl)indole-3-carboxylates was obtained. This may be due to lability of the cyano function under the reaction conditions.

In conclusion, 2-(2-isocyanophenyl)acetates have been utilized for the one-pot synthesis of 1-[(di)nitrophenyl]indole-3-carboxylates. Notable advantages of the present synthesis include: i) simplicity of the procedure, ii) milder reaction conditions, iii) readily availability of the starting materials, and iv) good yields of products. This procedure has been applied to the synthesis of 1,1'-(1,3-phenylene)diindole derivatives. Further elaboration of 2-(2-isocyanophenyl)acetates to the preparation of other types of heterocycles is currently being pursued in our laboratory.

EXPERIMENTAL

The melting points were determined on a Laboratory Devices MEL-TEMP II melting-point apparatus and are uncorrected. The IR spectra were recorded on a Shimadzu FTIR-8300 spectrometer. ^1H NMR spectra were determined using SiMe_4 as an internal reference in CDCl_3 with a JEOL ECP500 FT NMR spectrometer operating at 500 MHz. ^{13}C NMR spectra were determined using SiMe_4 as an internal reference with a JEOL ECP500 FT NMR spectrometer operating at 125 MHz in CDCl_3 . Low- and high-resolution mass spectra (EI, 70 eV) were recorded on a JEOL JMS-AX505 HA spectrometer. Thin-layer

chromatography (TLC) was carried out using Merck Kieselgel 60 PF₂₅₄. Column chromatography was performed using Merck Kieselgel 60 (0.063–0.200 mm). All of the solvents used were dried over the appropriate drying agents and distilled under argon prior to use.

Starting Materials. 2-Methylphenyl isocyanides (**1**) were prepared by a modification¹ of Ugi's Method.¹⁰ All other chemicals used in this study were commercially available.

Typical Procedure for the Preparation of 2-(2-Isocyanophenyl)acetates (2). Ethyl 2-(2-Isocyanophenyl)acetate (2a). To a stirred deep-red solution of 2-lithiomethylphenyl isocyanide, which was generated in situ by treating **1a** (1.2 g, 10 mmol) with 2 molar amount of LDA in diglyme (25 mL) at –78 °C, was added ethyl chloroformate (1.1 g, 10 mmol) dropwise. The color of the solution immediately turned into light red. Saturated aqueous NH₄Cl (30 mL) was added, and organic materials were extracted with Et₂O three times (30 mL each). The combined extracted was washed with water five times and then brine once, and dried over anhydrous Na₂SO₄. After evaporation of the solvent, the residue was purified by column chromatography on silica gel (1:5 THF–hexane) to give **2** (1.2 g, 65%); pale-yellow needles; mp 45–46 °C (hexane); IR (KBr) 2122, 1732 cm^{–1}; ¹H NMR δ 1.28 (t, *J* = 7.3 Hz, 3H), 3.80 (s, 2H), 4.20 (q, *J* = 7.3 Hz, 2H), 7.32 (dd, *J* = 7.3, 1.8 Hz, 1H), 7.34–7.42 (m, 3H). Anal. Calcd for C₁₁H₁₁NO₂: C, 69.83; H, 5.86; N, 7.40. Found: C, 69.68; H, 6.00; N, 7.25.

Ethyl 2-(2-Isocyano-5-methylphenyl)acetate (2b): a yellow oil; *R_f* 0.40 (1:7 THF–hexane); IR (neat) 2120, 1737 cm^{–1}; ¹H NMR δ 1.29 (t, *J* = 7.3 Hz, 3H), 2.36 (s, 3H), 3.75 (s, 2H), 4.20 (q, *J* = 7.3 Hz, 2H), 7.11 (d, *J* = 7.8 Hz, 1H), 7.14 (s, 1H), 7.28 (d, *J* = 7.8 Hz, 1H); MS *m/z* 203 (M⁺, 100). HR-MS. Calcd for C₁₂H₁₃NO₂: M, 203.0946. Found: *m/z* 203.0958.

Ethyl 2-(5-Chloro-2-isocyanophenyl)acetate (2c): a yellow oil; *R_f* 0.29 (1:7 THF–hexane); IR (neat) 2122, 1738 cm^{–1}; ¹H NMR δ 1.29 (t, *J* = 7.3 Hz, 3H), 3.76 (s, 2H), 4.21 (q, *J* = 7.3 Hz, 2H), 7.30 (dd, *J* = 8.2, 1.8 Hz, 1H), 7.35 (d, *J* = 8.2 Hz, 1H), 7.36 (d, *J* = 1.8 Hz, 1H); MS *m/z* 223 (M⁺, 100). HR-MS. Calcd for C₁₁H₁₀ClNO₂: M, 223.0400. Found: *m/z* 223.0391.

Ethyl 2-(2-Isocyano-5-methoxyphenyl)acetate (2d): a yellow oil; *R_f* 0.20 (1:7 Et₂O–hexane); IR (neat) 2118, 1738 cm^{–1}; ¹H NMR δ 1.28 (t, *J* = 7.3 Hz, 3H), 3.75 (s, 2H), 3.82 (s, 3H), 4.20 (q, *J* = 7.3 Hz, 2H), 6.80 (dd, *J* = 8.7, 2.7 Hz, 1H), 6.85 (d, *J* = 2.7 Hz, 1H), 7.33 (d, *J* = 8.7 Hz, 1H); MS *m/z* 219 (M⁺, 100). HR-MS. Calcd for C₁₂H₁₃NO₃: M, 219.0895. Found: *m/z* 219.0912.

Typical Procedure for the Preparation 1-Arylindole-3-carboxylates (5) and (7). Ethyl 1-(2-Nitrophenyl)indole-3-carboxylate (5a). To a stirred suspension of NaH (60% in oil; 21 mg, 0.53 mmol) in DMSO–THF (1:1 v/v, 2 mL) at rt was added a solution of **2** (0.10 g, 0.53 mmol) in THF (1 mL) dropwise; the mixture was stirred at the same temperature for 3 h. 1-Fluoro-2-nitrobenzene (**4a**) (75 mg, 0.53 mmol) was added, and stirring was continued for an additional 10 min. Saturated aqueous NH₄Cl (10 mL) was added, and organic materials were extracted with AcOEt three times (10 mL each). The combined extracted was washed with water three times and then brine once, and dried over

anhydrous Na_2SO_4 . After evaporation of the solvent, the residue was purified by preparative TLC on silica gel (1:3 THF–hexane) to give **5a** (0.14 g, 84%); a yellow solid; mp 110–111 °C (hexane– CH_2Cl_2); IR (KBr) 1692, 1528, 1346 cm^{-1} ; ^1H NMR δ 1.44 (t, $J = 7.3$ Hz, 3H), 4.42 (q, $J = 7.3$ Hz, 2H), 7.06 (d, $J = 7.8$ Hz, 1H), 7.25 (td, $J = 7.3, 1.4$ Hz, 1H), 7.33 (ddd, $J = 7.8, 7.3, 0.9$ Hz, 1H), 7.59 (dd, $J = 7.8, 1.4$ Hz, 1H), 7.69 (ddd, $J = 7.8, 7.3, 1.4$ Hz, 1H), 7.81 (ddd, $J = 7.8, 7.3, 1.4$ Hz, 1H), 7.90 (s, 1H), 8.13 (dd, $J = 7.8, 1.4$ Hz, 1H), 8.25 (d, $J = 7.3$ Hz, 1H); ^{13}C NMR δ 14.50, 60.05, 109.79, 110.80, 122.11, 122.80, 123.95, 125.79, 126.44, 129.69, 130.12, 131.64, 134.10 (two overlapped C's), 137.42, 146.19, 164.69; MS m/z 310 (M^+ , 100). Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$: C, 65.80; H, 4.55; N, 9.03. Found: C, 65.74; H, 4.35; N, 8.87.

Ethyl 1-(4-Methyl-2-nitrophenyl)indole-3-carboxylate (5b): a yellow solid; mp 144–145 °C (hexane– CH_2Cl_2); IR (KBr) 1699, 1539, 1348 cm^{-1} ; ^1H NMR δ 1.43 (t, $J = 7.3$ Hz, 3H), 2.57 (s, 3H), 4.41 (q, $J = 7.3$ Hz, 2H), 7.03 (d, $J = 8.2$ Hz, 1H), 7.24 (ddd, $J = 8.2, 7.3, 1.4$ Hz, 1H), 7.31 (ddd, $J = 7.8, 7.3, 0.9$ Hz, 1H), 7.44 (d, $J = 7.8$ Hz, 1H), 7.59 (ddd, $J = 7.8, 1.4$ Hz, 1H), 7.88 (s, 1H), 7.93 (s, 1H), 8.24 (d, $J = 7.8$ Hz, 1H); MS m/z 324 (M^+ , 100). Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_4$: C, 66.66; H, 4.97; N, 8.64. Found: C, 66.68; H, 5.08; N, 8.39.

Ethyl 1-(4-Nitrophenyl)indole-3-carboxylate (5c): pale-yellow needles; mp 150–152 °C (hexane– CH_2Cl_2); IR (KBr) 1686, 1545, 1346 cm^{-1} ; ^1H NMR δ 1.45 (t, $J = 7.3$ Hz, 3H), 4.44 (q, $J = 7.3$ Hz, 2H), 7.35–7.40 (m, 2H), 7.58 (d, $J = 7.3$ Hz, 1H), 7.74 (d, $J = 9.2$ Hz, 2H), 8.07 (s, 1H), 8.29 (dd, $J = 7.3, 1.4$ Hz, 1H), 8.45 (d, $J = 9.2$ Hz, 2H); MS m/z 310 (M^+ , 100). Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$: C, 65.80; H, 4.55; N, 9.03. Found: C, 65.74; H, 4.72; N, 9.24.

Ethyl 1-(2,4-Dinitrophenyl)indole-3-carboxylate (5d): a yellow solid; mp 165–167 °C (hexane– CH_2Cl_2); IR (KBr) 1697, 1543, 1344 cm^{-1} ; ^1H NMR δ 1.44 (t, $J = 7.3$ Hz, 3H), 4.43 (q, $J = 7.3$ Hz, 2H), 7.10 (d, $J = 8.2$ Hz, 1H), 7.32 (ddd, $J = 8.2, 7.3, 1.4$ Hz, 1H), 7.38 (ddd, $J = 8.2, 7.3, 0.9$ Hz, 1H), 7.86 (d, $J = 8.7$ Hz, 1H), 7.88 (s, 1H), 8.28 (d, $J = 8.2$ Hz, 1H), 8.65 (dd, $J = 8.7, 2.3$ Hz, 1H), 8.98 (d, $J = 2.3$ Hz, 1H); MS m/z 355 (M^+ , 100). Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{O}_6$: C, 57.47; H, 3.69; N, 11.83. Found: C, 56.43; H, 3.56; N, 11.44

Ethyl 5-Methyl-1-(2-nitrophenyl)indole-3-carboxylate (5e): a yellow oil; R_f 0.39 (1:4 THF–hexane); IR (neat) 1699, 1533, 1350 cm^{-1} ; ^1H NMR δ 1.43 (t, $J = 7.3$ Hz, 3H), 2.49 (s, 3H), 4.41 (q, $J = 7.3$ Hz, 2H), 6.94 (d, $J = 8.2$ Hz, 1H), 7.07 (dd, $J = 8.2, 1.4$ Hz, 1H), 7.58 (dd, $J = 8.2, 1.4$ Hz, 1H), 7.67 (ddd, $J = 8.2, 7.3, 1.4$ Hz, 1H), 7.80 (ddd, $J = 8.2, 7.3, 1.4$ Hz, 1H), 7.85 (s, 1H), 8.05 (s, 1H), 8.11 (dd, $J = 8.2, 1.4$ Hz, 1H); MS m/z 324 (M^+ , 100). Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_4$: C, 66.66; H, 4.97; N, 8.64. Found: C, 66.64; H, 4.94; N, 8.64.

Ethyl 5-Methyl-1-(4-nitrophenyl)indole-3-carboxylate (5f): a yellow solid; mp 165–167 °C (hexane– CH_2Cl_2); IR (KBr) 1684, 1549, 1342 cm^{-1} ; ^1H NMR δ 1.45 (t, $J = 7.3$ Hz, 3H), 2.52 (s, 3H), 4.43 (q, $J = 7.3$ Hz, 2H), 7.18 (d, $J = 8.2$ Hz, 1H), 7.47 (d, $J = 8.2$ Hz, 1H), 7.72 (d, $J = 8.7$ Hz, 2H), 8.02

(s, 1H), 8.08 (s, 1H), 8.44 (d, $J = 8.7$ Hz, 2H); MS m/z 324 (M^+ , 100). Anal. Calcd for $C_{18}H_{16}N_2O_4$: C, 66.66; H, 4.97; N, 8.64. Found: C, 66.39; H, 4.90; N, 8.41.

Ethyl 1-(2,4-Dinitrophenyl)-5-methylindole-3-carboxylate (5g): a yellow solid; mp 132–134 °C (hexane–CH₂Cl₂); IR (KBr) 1678, 1541, 1346 cm⁻¹; ¹H NMR δ 1.44 (t, $J = 7.3$ Hz, 3H), 2.50 (s, 3H), 4.43 (q, $J = 7.3$ Hz, 2H), 6.99 (d, $J = 8.2$ Hz, 1H), 7.12 (d, $J = 8.2$ Hz, 1H), 7.83 (s, 1H), 7.84 (d, $J = 8.7$ Hz, 1H), 8.07 (s, 1H), 8.63 (dd, $J = 8.7, 2.7$ Hz, 1H), 8.97 (d, $J = 2.7$ Hz, 1H); ¹³C NMR δ 14.51, 21.52, 60.34, 109.07, 112.44, 121.84, 122.33, 126.24, 127.12, 128.39, 130.52, 132.94, 133.49, 134.97, 137.02, 145.21, 146.54, 164.25; MS m/z 369 (M^+ , 100). Anal. Calcd for $C_{18}H_{15}N_3O_6$: C, 58.54; H, 4.09; N, 11.38. Found: C, 58.66; H, 4.17; N, 11.07.

Ethyl 5-Chloro-1-(2-nitrophenyl)indole-3-carboxylate (5h): a yellow solid; mp 128–130 °C (hexane–CH₂Cl₂); IR (KBr) 1690, 1535, 1364 cm⁻¹; ¹H NMR δ 1.44 (t, $J = 7.3$ Hz, 3H), 4.42 (q, $J = 7.3$ Hz, 2H), 6.96 (d, $J = 8.7$ Hz, 1H), 7.21 (dd, $J = 8.7, 2.3$ Hz, 1H), 7.57 (d, $J = 7.8$ Hz, 1H), 7.71 (ddd, $J = 7.8, 7.3, 1.4$ Hz, 1H), 7.82 (ddd, $J = 7.8, 7.3, 1.4$ Hz, 1H), 7.90 (s, 1H), 8.14 (d, $J = 7.8$ Hz, 1H), 8.23 (d, $J = 2.3$ Hz, 1H); MS m/z 344 (M^+ , 100). Anal. Calcd for $C_{17}H_{13}ClN_2O_4$: C, 59.23; H, 3.80; N, 8.13. Found: C, 59.22; H, 3.83; N, 7.84.

Ethyl 5-Chloro-1-(2,4-dinitrophenyl)indole-3-carboxylate (5i): an orange solid; mp 182–183 °C (hexane–CH₂Cl₂); IR (KBr) 1607, 1535, 1366 cm⁻¹; ¹H NMR δ 1.46 (t, $J = 7.3$ Hz, 3H), 4.44 (q, $J = 7.3$ Hz, 2H), 7.00 (d, $J = 9.2$ Hz, 1H), 7.27 (dd, $J = 9.2, 1.8$ Hz, 1H), 7.85 (d, $J = 8.7$ Hz, 1H), 7.88 (s, 1H), 8.26 (d, $J = 1.8$ Hz, 1H), 8.67 (dd, $J = 8.7, 2.7$ Hz, 1H), 8.99 (d, $J = 2.7$ Hz, 1H); MS m/z 389 (M^+ , 100). Anal. Calcd for $C_{17}H_{12}ClN_3O_6$: C, 52.39; H, 3.10; N, 10.78. Found: C, 52.34; H, 3.20; N, 10.68.

Ethyl 1-(2,4-Dinitrophenyl)-5-methoxyindole-3-carboxylate (5j): an orange solid; mp 190–191 °C (hexane–THF); IR (KBr) 1682, 1607, 1541, 1346 cm⁻¹; ¹H NMR δ 1.43 (t, $J = 7.3$ Hz, 3H), 3.91 (s, 3H), 4.42 (q, $J = 7.3$ Hz, 2H), 6.93 (dd, $J = 8.7, 2.3$ Hz, 1H), 7.00 (d, $J = 8.7$ Hz, 1H), 7.75 (d, $J = 2.3$ Hz, 1H), 7.81 (s, 1H), 7.85 (d, $J = 8.7$ Hz, 1H), 8.63 (dd, $J = 8.7, 2.3$ Hz, 1H), 8.96 (d, $J = 2.3$ Hz, 1H); MS m/z 385 (M^+ , 100). Anal. Calcd for $C_{18}H_{15}N_2O_7$: C, 56.11; H, 3.92; N, 10.91. Found: C, 56.07; H, 4.02; N, 10.85.

Ethyl 1-[2,4-Dinitrophenyl-5-(3-ethoxycarbonylindol-1-yl)]indole-3-carboxylate (7a): an orange-yellow solid; mp 103–105 °C (hexane–CH₂Cl₂); IR (KBr) 1699, 1539, 1339 cm⁻¹; ¹H NMR δ 1.44 (t, $J = 7.3$ Hz, 6H), 4.43 (q, $J = 7.3$ Hz, 4H), 7.18 (d, $J = 8.2$ Hz, 2H), 7.36 (ddd, $J = 8.2, 7.3, 1.4$ Hz, 2H), 7.40 (ddd, $J = 7.8, 7.3, 1.4$ Hz, 2H), 7.90 (s, 2H), 7.91 (s, 1H), 8.28 (d, $J = 7.8$ Hz, 2H), 8.95 (s, 1H); ¹³C NMR δ 14.45, 60.49, 109.24, 113.40, 122.84, 123.92, 124.85, 125.05, 126.86, 130.15, 132.73, 136.45, 136.59, 143.15, 163.93; MS m/z 542 (M^+ , 100). Anal. Calcd for $C_{28}H_{22}N_4O_8$: C, 61.99; H, 4.09; N, 10.33. Found: C, 61.66; H, 4.36; N, 10.06.

Ethyl 1-[2,4-Dinitrophenyl-5-(3-ethoxycarbonyl-5-methoxyindol-1-yl)]-5-methoxyindole-3-carboxylate (7b): a yellow solid; mp 212–214 °C (hexane–CH₂Cl₂); IR (KBr) 1695, 1616, 1541, 1346

cm⁻¹; ¹H NMR δ 1.43 (t, *J* = 7.3 Hz, 6H), 3.91 (s, 6H), 4.42 (q, *J* = 7.3 Hz, 4H), 6.97 (dd, *J* = 8.7, 2.7 Hz, 2H), 7.06 (d, *J* = 8.7 Hz, 2H), 7.74 (d, *J* = 2.7 Hz, 2H), 7.84 (s, 2H), 7.88 (s, 1H), 8.90 (s, 1H); MS *m/z* 602 (M⁺, 100). Anal. Calcd for C₃₀H₂₆N₄O₁₀: C, 59.80; H, 4.35; N, 9.30. Found: C, 59.68; H, 4.40; N, 9.13.

ACKNOWLEDGEMENTS

We are grateful to Mrs. Miyuki Tanmatsu of this University for her support in determining mass spectra and performing combustion analyses.

REFERENCES AND NOTES

1. Y. Ito, K. Kobayashi, N. Seko, and T. Saegusa, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 73 and pertinent references cited in this paper.
2. Y. Ito, K. Kobayashi, and T. Saegusa, *J. Am. Chem. Soc.*, 1977, **99**, 3532.
3. Y. Ito, Y. Inubushi, T. Sugaya, K. Kobayashi, and T. Saegusa, *Bull. Chem. Soc. Jpn.*, 1978, **51**, 1186.
4. For recent reports on the biological activity of indole-3-carboxylates: (a) Anti-hepatitis B virus activities; H. Chai, Y. Zhao, C. Zhao, and P. Gong, *Bioorg. Med. Chem.*, 2006, **14**, 911. (b) Anti-hepatitis B virus activities; C. Zhao, Y. Zhao, H. Chai, and P. Gong, *Bioorg. Med. Chem.*, 2006, **14**, 2552. Indole-3-carboxylates have recently utilized for the synthesis of more complex molecules: (c) C. R. Hopkins, M. Czekaj, S. S. Kaye, Z. Gao, J. Pribish, H. Pauls, G. Liang, K. Sides, D. Cramer, K. Cairns, Y. Luo, H.-K. Lim, R. Vaz, S. Rebello, S. Maignan, A. Dupuy, M. Mathieu, and J. Levell, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2734.
5. For recent syntheses of indole-3-carboxylates: (a) K. Yamazaki and Y. Kondo, *J. Comb. Chem.*, 2002, **4**, 191. (b) K. Yamazaki, Y. Nakamura, and Y. Kondo, *J. Org. Chem.*, 2003, **68**, 6011. (c) M. S. Islam, C. Brennan, Q. Wang, and M. M. Hossain, *J. Org. Chem.*, 2006, **71**, 4675.
6. (a) Using Nenitzescu reaction; U. Kuckländer and W. Hühnermann, *Arch. Pharm.*, 1979, **312**, 515. (b) Via PIFA-mediated intramolecular cyclization of 2-aryl-3-arylaminoacrylates: Y. Du, R. Liu, G. Linn, and K. Zhao, *Org. Lett.*, 2006, **8**, 5919. (c) Via a reaction of *N,N*-diphenylhydrazine with diethyl acetylenedicarboxylate: I. A. Sayyed, K. Alex, A. Tillack, N. Schwarz, D. Michalik, and M. Beller, *Eur. J. Org. Chem.*, 2007, 4525. (d) Via a copper-catalyzed reaction of methyl 2-(2-bromophenyl)acetate with methyl formate and arylamines: F. S. Melkonyan, A. V. Karchava, and M. A. Yurovskaya, *J. Org. Chem.*, 2008, **73**, 4275.
7. (a) J. C. Antilla, A. Klapars, and S. L. Buchwald, *J. Am. Chem. Soc.*, 2002, **124**, 11684. (b) F. Bellina, C. Calandri, S. Cauteruccio, and R. Rossi, *Eur. J. Org. Chem.*, 2007, 2147.
8. *N*-4-Nitrophenylation of indole with 4-iodonitrobenzene has been achieved using copper catalysts

- at higher temperatures (90–160 °C): (a) A. Klapars, J. C. Antilla, X. Huang, and S. L. Buchwald, *J. Am. Chem. Soc.*, 2001, **123**, 7727. (b) K. Okano, H. Tokuyama, and T. Fukuyama, *Org. Lett.*, 2003, **5**, 4987. See also ref. 7b.
9. 1-(2-Nitrophenyl)- and 1-(2,4-dinitrophenyl)indoles have been prepared by *N*-arylation of indoline-2-carboxylic acid followed by decarboxylation and dehydrogenation; R. Chicharro, S. de Castro, J. L. Reino, and V. J. Arán, *Eur. J. Org. Chem.*, 2003, 2314.
10. I. Ugi and R. Meyr, *Org. Synth., Coll. Vol. V*, 1973, 1060.