

HETEROCYCLES, Vol. 78, No. 5, 2009, pp. 1177 - 1181. © The Japan Institute of Heterocyclic Chemistry
 Received, 8th December, 2008, Accepted, 23rd January, 2009, Published online, 27th January, 2009
 DOI: 10.3987/COM-08-11621


FACILE SYNTHESIS OF 1-ARYL-2,3-DIHYDRO-1H-ISOINDOLES BY CYCLIZATION OF *N*-FORMYLIMINIUM ION VIA GEOMETRICALLY DISFAVORED 5-*ENDO*-TRIG PROCESS

Michikazu Kitabatake, Toshiaki Saitoh, Takehiro Sano, and Yoshie Horiguchi*

Showa Pharmaceutical University, 3-3165 Higashi-tamagawagakuen, Machida, Tokyo 194-8543, Japan; e-mail: horiguti@ac.shoyaku.ac.jp

Abstract — Synthesis of 1-aryl-2,3-dihydro-1*H*-isoindoles (isoindolines) (**10**) was achieved in a highly effective manner *via* acid catalyzed cyclization of *N*-formyliminium ion (**8**) obtained from 2,3-dimethoxybenzylamine and carbonyl compounds with acetic-formic anhydride under a one pot procedure. This Pictet-Spengler type reaction provides a convenient method for preparing 1-arylisoindolines.

Compounds containing 2,3-dihydro-1*H*-isoindoles (isoindoline) (**1**) such as 1-methyl-1-phenyl-(**3**) and 1-phenyl-derivative (**4**) are known to be a non-competitive NMDA antagonist^{1, 2} and a phencyclidine agonist³ respectively. However, isoindoline derivatives are very rare in spite of biological interest, when




Scheme 1 2,3-Dihydro-1*H*-isoindoles and related compounds as a NMDA antagonist and a phencyclidine agonist

compared to 1,2,3,4-tetrahydroisoquinoline (**2**) derivatives that constitute the fundamental ring system of various medicines. Although there are several syntheses of isoindoline in the literature,⁴⁻¹⁰ only one paper which deals with a general methodology leading to the isoindoline ring system is reported.¹⁰

In this paper we report the synthesis of 1-aryl-2,3-dihydro-1*H*-isoindole (**10**) utilizing a simple and convenient method *via* a Pictet-Spengler type reaction, that is, the intramolecular cyclization of *N*-formyliminium ion (**8**). We have recently developed this methodology for the synthesis of tetrahydroisoquinoline (**2**) producing not only 1-substituted but also 1,1-disubstituted derivatives.¹¹

The *N*-formyliminium ions (**8a~e**), substrates of the reaction, were readily prepared by formylation of imines (**7a~e**) with acetic-formic anhydride. The imines (**7a~e**) were obtained by heating a mixture of 3,4-dimethoxybenzylamine (**5**) and arylaldehydes (**6a~d**) or acetophenone (**6e**). The procedures are shown in Table 1.

The intramolecular cyclization of *N*-formyliminium ions (**8**) leading to the isoindoline ring system *via* 5-*endo*-trig process was anticipated to be difficult to achieve when compared with the Pictet-Spengler reaction to 1,2,3,4-tetrahydroisoquinoline *via* 6-*endo*-trig; according to the Baldwin rule¹² the former reaction of 5-*endo*-trig process is geometrically disfavored while the latter 6-*endo*-trig one is favored. This disadvantage was overcome by finding reaction conditions using a large excess of trifluoroacetic acid (TFA) described in Table 1. However, when the process of removing acetic-formic anhydride from the reaction mixture was omitted or inadequate, the cyclization did not occur in a practical sense. This fact revealed that the acidity of the reaction solution is critically important to produce this intramolecular cyclization. Thus, the 1-aryl-2,3-dihydro-2-formyl-1*H*-isoindoles (**9**) were obtained in fairly good yields, as shown in Table 1.¹³ Hydrolysis of **9** with hydrochloric acid yielded the corresponding 1-aryl-2,3-dihydro-1*H*-isoindoles (**10**) in a quantitative yield.¹³ The direct cyclization of the imine (**7**) to the isoindoline (**10**) did not occur at all even when treated with TFA even under more forced conditions, indicating that the cyclization is the reaction of *N*-formyliminium ions (**8**) activated by *N*-formyl group. These experiments clearly demonstrated that the cyclization of the *N*-formyliminium ion to isoindoline



Scheme 2 Synthesis of 1-aryl-2,3-dihydro-1H-isoindoles

Table 1. Synthesis of 1-Aryl-2,3-dihydro-2-formyl-1H-isoindoles (**9**) byAcid Catalyzed Cyclization of *N*-Formyliminium Ions (**8**).

Run	<i>N</i> -Formyliminium ion (8)* ¹			Cyclization* ²				Yields of 9 (%) ^{*3}
	R ¹	R ²	R ³	Reagent	Temp (°)	Time(h)		
1	8a	H	H	H	TFA	90	3	9a (65)
2	8b	OMe	OMe	H	TFA	90	3	9b (80)
3	8c	OMe	H	H	TFA	90	3	9c (83)
4	8d	H	OMe	H	TFA	90	3	9d (38)
5	8e	H	H	Me	TFA	90	5	9e (86)

*1a) **8a~d** were prepared by heating the mixture of **5** (1.2 mol eq.) and **6** (1 mol eq.) without solvent at 80 °C for 1 h, followed by heating at 70 °C for 1 h in acetic-formic anhydride (100 mol eq.) which was prepared from acetic anhydride (100 mol eq.) and formic acid (100 mol eq.)


*1b) **8e** was prepared as follows: the mixture of **5** (1.2 mol eq.) and **6e** (1 mol eq.) was heated in Ti(*i*-OPr)₄ (1.5 mol eq.) at 80 °C for 1 h, and the resulting imine **7e** was formylated by acetic-formic anhydride (100 mol eq.) at 70 °C for 1 h without removing the Ti catalyst.

*2) The reaction was carried out in large excess of TFA (100 mol eq.) after eliminating excess acetic-formic anhydride by heating *in vacuo*.

*3) The yields of **9** were calculated based on the quantity of the carbonyl compounds (**6**).

occurs through the geometrically disfavored 5-*endo* trigonal process. It is especially noteworthy that the *N*-formyliminium ion (**8e**), in spite of having the congested tetra-substituted C=N⁺ system, readily caused the cyclization to give **9e** in 86% yield.

The occurrence of this cyclization may be explained in terms of mesomeric effect. The C=N⁺ bond is conjugated to the benzene ring. Therefore, the energy of the transition state should be lowered by the contribution of mesomeric enamino-structure **8B** featuring 5-*exo* trigonal cyclization. Thus, the 5-*endo* trig process of the imino-structure **8A** to **9**, although geometrically disfavored, will be facilitated by the contribution of **8B** that is geometrically favored. The relatively low yield of isoindoline **9d** probably is attributable to the instability of the cation of **8B** which is induced by the electron attractive *meta* OMe group (R²).



Scheme 3 Mesomeric structure of *N*-formyliminium ion

Thus, the Pictet-Spengler type reaction of *N*-formyliminium ions obtained from 3,4-dimethoxybenzylamine and carbonyl compounds under a one pot procedure provides a convenient method for preparing 1-arylisoidolines. Investigation concerning the scope and limitation of this methodology is underway.

REFERENCES AND NOTES

1. J. P. Hodgkiss, H. J. Sherriffs, D. A. Contrell, K. Shirakawa, J. S. Kelly, A. Kuno, M. Ohkubo, S. P. Bucher, and H. J. Olverman, *Eur. J. Pharmacol.*, 1993, **240**, 219.
2. M. Ohkubo, A. Kuno, K. Katsita, Y. Ueda, K. Shirakawa, H. Nakanishi, I. Nakanishi, T. Kinoshita, and H. Takasugi, *Chem. Pharm. Bull.*, 1996, **44**, 95.

3. N. M. Gray, B. K. Cheng, J. M. Stephen, C. M. Lair, and P. C. Contreras, *J. Med. Chem.*, 1989, **32**, 1242.
4. J. Bornstein, S. F. Bedell, P. E. Drummond, and C. L. Kosloski, *J. Am. Chem. Soc.*, 1956, **78**, 83.
5. D. F. Veber and W. Lwowski, *J. Am. Chem. Soc.*, 1964, **86**, 4152.
6. G. Gast, J. Schmutz, and D. Sorg, *Helvetica Chim. Acta*, 1977, **60**, 1644.
7. A. I. Meyers and B. Santiago, *Tetrahedron Lett.*, 1995, **36**, 5877.
8. H. Ishibashi, I. Kato, Y. Takeda, M. Kogure, and O. Tamura, *Chem. Commun.*, **2000**, 1527.
9. D.-R. Hou, M.-S. Wang, M.-W. Chung., Y.-D. Hsieh, and H.-H. G. Tsai, *J. Org. Chem.*, 2007, **72**, 9231.
10. K. Kobayashi, S. Kondo, K. Hashimoto, S. Fukamachi, and O. Konishi, *Heterocycles*, 2007, **71**, 1827.
11. Y. Horiguchi, H. Kodama, M. Nakamura, T. Yoshimura, K. Hanezi, H. Hamada, T. Saitoh, and T. Sano, *Chem. Pharm. Bull.*, 2002, **50**, 253.
12. J. E. Baldwin, *J. Chem. Soc., Chem. Commun.*, 1976, **18**, 738.
13. The structures of **9** and **10** were determined by elementary and spectral analyses (Mass, ¹H- and ¹³C-NMR data).