

HETEROCYCLES, Vol. 80, No. 1, 2010, pp. 613 - 621. © The Japan Institute of Heterocyclic Chemistry
Received, 19th May, 2009, Accepted, 25th June, 2009, Published online, 26th June, 2009
DOI: 10.3987/COM-09-S(S)28

BIYOUXANTHONES A – D, PRENYLATED XANTHONES FROM ROOTS OF *HYPERICUM CHINENSE*

Naonobu Tanaka,^a Takuji Mamemura,^a Shuhei Abe,^a Kiyoshi Imabayashi,^b
Yoshiki Kashiwada,^b Yoshihisa Takaishi,^b Tetsuro Suzuki,^c
Yutaka Takebe,^d Takaaki Kubota,^a and Jun'ichi Kobayashi^{a,*}

^aGraduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

^bGraduate School of Pharmaceutical Sciences, University of Tokushima, Tokushima 770-8505, Japan


^cDepartment of Virology II, National Institute of Infectious Disease, Tokyo, 162-8640, Japan

^dAIDS Research Center, National Institute of Infectious Disease, Tokyo, 162-8640, Japan

E-mail: jkobay@pharm.hokudai.ac.jp

Abstract - Four new prenylated xanthones, biyouxanthones A – D (**1** – **4**), were isolated from roots of *Hypericum chinense*. The gross structures of **1** – **4** were elucidated from spectroscopic data, and **2** was assigned as a racemate by chiral HPLC analysis. Biyouxanthones A (**1**) and B (**2**) exhibited anti-viral activity against hepatitis C virus (HCV).

The plants, belonging to the genus *Hypericum* (family Clusiaceae), are known to be a traditional medicine for the treatment of burns, bruises, swelling, inflammation, and anxiety as well as bacterial and viral infections.¹⁻⁴ In our continuing search for bioactive compounds from *Hypericum* spp,⁵ three new prenylated xanthones possessing 2-hydroxycyclohexa-2,4-dienone moiety, biyouxanthones A – C (**1** – **3**), and one new prenylated xanthone having a 5-methyl-2-(prop-1-en-2-yl)hex-4-enyl group, biyouxanthone D (**4**), were isolated from roots of *H. chinense*. In this paper, we describe the isolation, structure elucidation of **1** – **4** and anti-HCV activity of **1** and **2**.



The dried roots of *H. chinense* were extracted with MeOH, and the extracts were partitioned successively with *n*-hexane, EtOAc, and H₂O. *n*-Hexane-soluble portions were separated by a silica gel column (*n*-hexane/EtOAc) and then a Sephadex LH-20 column (MeOH) to afford a mixture of xanthone derivatives, which was purified by C₁₈ reversed-phase HPLC (MeOH/H₂O) to yield biyouxanthones A (1, 0.00017%), B (2, 0.00016%), and C (3, 0.00087%). EtOAc-soluble portions were purified by a silica gel column (CHCl₃/MeOH) and a Sephadex LH-20 (MeOH) column chromatographies, and C₁₈ reversed-phase HPLC (MeOH/H₂O, 0.1% TFA) to give biyouxanthone D (4, 0.00012%).

The molecular formula of biyouxanthone A (**1**), C₃₈H₄₈O₆, was revealed by HRESIMS (*m/z* 599.3366 [M-H]⁻, Δ -0.6 mmu). IR absorptions at 1671 and 1636 cm⁻¹ implied the presence of carbonyl




Figure 1. Selected 2D NMR correlations for bijouxanthone A (1).

functionalities. ^1H and ^{13}C NMR data of **1** (Table 1) revealed the presence of three carbonyl groups, seven trisubstituted olefins, two tetrasubstituted olefins, two sp^3 quaternary carbons, six sp^3 methylenes, nine tertiary methyls, and one hydrogen-bonded hydroxy group. ^1H – ^1H COSY correlations for H-11 and H-12, H-16 and H-17, and H-31 and H-32, and HMBC cross-peaks of H₃-14 to C-12, C-13, and C-15, H₃-19 to C-17, C-18, and C-20, H₃-34 to C-32, C-33, and C-35 suggested the presence of three prenyl groups. The

Table 1. ^1H and ^{13}C NMR data for biyouxanthone A (**1**) in CDCl_3

position	δ_{C}	δ_{H} (<i>J</i> in Hz)
1	170.0	-
2	103.6	5.62 (1H, s)
3	195.9	-
4	57.6	-
4a	173.5	-
5	106.5	6.47 (1H, s)
6	160.1	-
7	200.6	-
8	56.3	-
8a	121.3	-
9	177.0	-
9a	112.2	-
10a	152.5	-
11	38.4	2.81 (2H, m)
12	117.0	4.82 (1H, t, <i>J</i> = 6.9)
13	135.5	-
14	25.7	1.55 (3H, s)
15	17.9	1.47 (3H, s)
16	38.6	2.70 (2H, m)
17	117.1	4.80 (1H, t, <i>J</i> = 6.9)
18	135.6	-
19	25.7	1.55 (3H, s)
20	17.7	1.48 (3H, s)
21	37.5	3.40 (1H, dd, <i>J</i> = 14.5, 7.5) 2.79 (1H, m)
22	116.9	4.60 (1H, t, <i>J</i> = 7.5)
23	139.8	-
24	39.8	1.79 (2H, m)
25	26.5	1.91 (2H, m)
26	123.5	4.90 (1H, t, <i>J</i> = 7.2)
27	131.7	-
28	25.6	1.63 (3H, s)
29	17.6	1.51 (3H, s)
30	16.1	1.48 (3H, s)
31	37.5	3.41 (1H, dd, <i>J</i> = 14.5, 7.5) 2.79 (1H, m)
32	117.1	4.55 (1H, t, <i>J</i> = 7.5)
33	136.1	-
34	25.7	1.51 (3H, s)
35	17.7	1.48 (3H, s)
1-OH	-	13.53 (1H, s)

presence of a geranyl group was implied by ^1H - ^1H COSY correlations for H-21 and H-22, H-24 and H-25, H-25 and H-26, and HMBC cross-peaks of H₃-30 to C-22, C-23, and C-24, and H₃-28 to C-26, C-27, and C-29. NOESY correlations for H-21 to H₃-30 and H-22 to H₂-24 suggested the 22*E* configuration for the geranyl group. From these data, **1** was presumed to be a xanthone derivative possessing one geranyl and three prenyl groups. The chemical shift of OH-1 (δ_{H} 13.53) revealed the presence of hydrogen-bonded hydroxy group at C-1. The substitution pattern of A-ring (C-1 to C-4, C-4a, and C-9a) was deduced from HMBC cross-peaks of H-2 to C-1, C-3, and C-9a, and H₂-16 to C-3, C-4, C-4a, and C-11. HMBC correlations for H₂-31 to C-7, C-8, C-8a, and C-21 indicated that C-7, C-8a, a prenyl group, and a geranyl group were attached to an sp^3 quaternary carbon (C-8). Connectivities of C-8a to C-10a, C-10a to C-5, and C-5 to C-7 were deduced from HMBC correlations for H-5 to C-6, C-7, C-8a, and C-10a. Thus, the gross structure of biyouxanthone A was elucidated to be **1** (Figure 1).

The molecular formula of biyouxanthone B (**2**), $\text{C}_{34}\text{H}_{42}\text{O}_6$, was established by HRESIMS (m/z 545.2892 [$\text{M}-\text{H}$]⁻, Δ -1.1 mmu). IR absorptions at 1668 and 1645 cm^{-1} implied the presence of carbonyl functionalities. ^1H and ^{13}C NMR data of **2** (Table 2) suggested that **2** was a xanthone derivative possessing one geranyl, two prenyl, one methoxy, and two hydroxy groups. Carbon chemical shifts of ring B (C-5 to C-8, C-8a, and C-10a) and C-16 to C-30 were similar to corresponding carbon chemical shifts of **1**. The substitution pattern of A-ring (C-1 to C-4, C-4a, and C-9a) as shown in Figure 2 were deduced by HMBC cross-peaks of a hydrogen-bonded hydroxy proton (OH-1) to C-1, C-2, and C-9a, H₂-11 to C-3, C-4, and C-4a, H-2 and 3-OMe to C-3, and the NOESY correlation between H-2 and 3-OMe. Thus, the gross structure of biyouxanthone B was elucidated to be **2**. Biyouxanthone B (**2**) showed no specific rotation. Chiral HPLC analysis of biyouxanthone B (**2**) gave two peaks with almost same peak areas due to (-)-bonyouxanthone B (t_{R} 19.0 min) and (+)-bonyouxanthone B (t_{R} 24.0 min). In




Figure 2. Selected 2D NMR correlations for biyouxanthone B (**2**)

addition, patterns of Cotton effects for respective enantiomers in the CD spectra were symmetrical each other. From these data, biyouxanthone B (**2**) was assigned as a racemate.

Biyouxanthone C (**3**) showed the pseudomolecular ion peak at *m/z* 531 [M–H][–] in the ESIMS, and the HRESIMS analysis revealed that the molecular formula of **3** was C₃₃H₄₀O₆, whose molecular weight is

Table 2. ¹H and ¹³C NMR data for biyouxanthones B (**2**) and C (**3**) in CDCl₃

position	δ_{C}	δ_{H} (<i>J</i> in Hz)	δ_{C}	δ_{H} (<i>J</i> in Hz)
1	161.0	-	160.4	-
2	95.2	6.41 (1H, s)	99.6	6.33 (1H, s)
3	162.6	-	160.6	-
4	107.2	-	104.6	-
4a	153.2	-	153.7	-
5	108.8	6.52 (1H, s)	108.7	6.52 (1H, s)
6	159.2	-	159.0	-
7	201.4	-	201.4	-
8	55.9	-	55.9	-
8a	116.2	-	116.4	-
9	179.7	-	179.6	-
9a	104.6	-	105.1	-
10a	151.7	-	151.8	-
11	21.5	3.44 (2H, m)	21.7	3.51 (2H, m)
12	121.8	5.21 (1H, t, <i>J</i> = 7.2)	121.1	5.28 (1H, t, <i>J</i> = 6.8)
13	131.9	-	135.1	-
14	25.7	1.70 (3H, s)	25.8	1.78 (3H, s)
15	17.9	1.83 (3H, s)	17.9	1.87 (3H, s)
16	37.7	3.48 (1H, m)	37.7	3.44 (1H, m)
		2.81 (1H, dd, <i>J</i> = 13.2, 7.8)		2.80 (1H, dd, <i>J</i> = 14.8, 7.8)
17	117.8	4.69 (1H, t, <i>J</i> = 7.8)	117.7	4.68 (1H, t, <i>J</i> = 7.8)
18	139.0	-	139.0	-
19	39.6	1.80 (2H, m)	39.6	1.83 (2H, m)
20	26.6	1.86 (2H, m)	26.5	1.80 (2H, m)
21	123.9	4.89 (1H, t, <i>J</i> = 6.7)	123.9	4.88 (1H, t, <i>J</i> = 6.8)
22	131.3	-	131.4	-
23	25.5	1.60 (3H, s)	25.5	1.60 (3H, s)
24	17.8	1.49 (3H, s)	17.9	1.50 (3H, s)
25	16.2	1.48 (3H, s)	16.2	1.48 (3H, s)
26	38.0	3.48 (1H, m)	37.9	3.44 (1H, m)
		2.83 (1H, dd, <i>J</i> = 12.5, 7.5)		2.83 (1H, dd, <i>J</i> = 13.9, 7.1)
27	117.7	4.66 (1H, t, <i>J</i> = 7.5)	117.6	4.65 (1H, t, <i>J</i> = 7.1)
28	135.2	-	135.3	-
29	25.7	1.50 (3H, s)	25.7	1.50 (3H, s)
30	17.5	1.50 (3H, s)	17.5	1.50 (3H, s)
1-OH	-	13.25 (1H, s)	-	13.23 (1H, s)
3-OH	-	-	-	-
6-OH	-	6.98 (1H, brs)	-	-
3-OMe	56.0	3.91 (3H, s)	-	-

smaller by 14 as compared with that of biyouxanthone B (**2**). ^1H and ^{13}C NMR data of biyouxanthone C (**3**) (Table 2) were similar to those of **2**, except for the absence of the signal for a methoxy group at C-3. These data suggested that **3** had a hydroxy group at C-3 in place of a methoxy group of **2**. It was supported by detailed analyses of 2D NMR spectra of **3** (Figure 3). Thus, the gross structure of biyouxanthone C was elucidated to be **3**.




Figure 3. Selected 2D NMR correlations for biyouxanthone C (**3**)

Biyouxanthone D (**4**) had a molecular formula of $\text{C}_{28}\text{H}_{32}\text{O}_6$ based on HRESIMS. ^1H and ^{13}C NMR data of **4** (Table 3) revealed that **4** was a xanthone derivative possessing one prenyl group, four hydroxy groups, and one monoterpene substituent. The substituent was elucidated to be a 5-methyl-2-(prop-1-en-2-yl)hex-4-enyl group by ^1H - ^1H COSY correlations for $\text{H}_2\text{-}16$ and $\text{H}\text{-}17$, $\text{H}\text{-}17$ and $\text{H}_2\text{-}18$, and $\text{H}_2\text{-}18$ and $\text{H}\text{-}19$, and HMBC cross-peaks of $\text{H}_3\text{-}21$ to C-19, C-20, and C-22, and $\text{H}_3\text{-}25$ to C-17, C-23, and C-24. The locations of these substituents were deduced from the HMBC correlations as shown in Figure 4. Thus, the gross structure of biyouxanthone D was elucidated to be **4**. Biyouxanthone D (**4**) showed optical rotation, $[\alpha]^{23}\text{D} +37.9$, while the stereochemistry of C-17 was not assigned.




Figure 4. Selected 2D NMR correlations for biyouxanthone D (**4**)

Biyouxanthones A (**1**) and B (**2**) inhibited the HCV core protein level in the culture of HCV-infected human hepatoma Huh7 cells (89% and 61%, respectively) at 10 μ M.

Table 3. ^1H and ^{13}C NMR data for biyouxanthone D (**4**) in CDCl_3

position	δ_{C}	δ_{H} (<i>J</i> in Hz)
1	158.4	-
2	108.6	-
3	160.5	-
4	105.8	-
4a	152.7	-
5	130.2	-
6	149.0	-
7	112.3	6.98 (1H, d, <i>J</i> = 8.8)
8	118.1	7.77 (1H, d, <i>J</i> = 8.8)
8a	114.0	-
9	180.5	-
9a	102.6	-
10a	144.7	-
11	21.5	3.49 (2H, d, <i>J</i> = 7.2)
12	121.0	5.27 (1H, t, <i>J</i> = 7.2)
13	136.4	-
14	25.8	1.80 (3H, s)
15	17.9	1.87 (3H, s)
16	27.2	2.93 (1H, dd, <i>J</i> = 14.0, 6.4) 2.84 (1H, dd, <i>J</i> = 14.0, 7.3)
17	47.7	2.39 (1H, quint, <i>J</i> = 7.0)
18	31.8	2.23, 2.15 (each 1H, dt, <i>J</i> = 15.2, 6.7)
19	122.6	5.20 (1H, t, <i>J</i> = 6.7)
20	133.8	-
21	25.8	1.75 (3H, s)
22	17.9	1.62 (3H, s)
23	150.0	-
24	110.6	4.76, 4.69 (each 1H, brs)
25	19.6	1.75 (3H, s)
1-OH	-	13.32 (1H, s)
3-OH	-	6.34 (1H, s)

EXPERIMENTAL

General Experimental Procedures

Optical rotations were recorded on a JASCO P-1030 digital polarimeter. IR, UV, and CD spectra were recorded on JASCO FT/IR-230, Shimadzu UV-1600PC, and JASCO J-720 spectrophotometers, respectively. NMR spectra were measured by a JEOL ECA 500 spectrometer. The 7.26 and 77.0 ppm resonances of residual CHCl_3 were used as internal references for ^1H and ^{13}C NMR spectra, respectively. ESIMS spectra were recorded on a JEOL JMS 700-TZ spectrometer.

Plant Material

Hypericum chinense was cultivated at the botanical garden of the University of Tokushima and collected in January 2006. Herbarium specimens were deposited in Experimental Station for Medicinal Plants Studies, Hokkaido University (specimen number: UTP98014).

Extraction and Isolation

Roots of *H. chinense* (2.52 kg, dry) were extracted with MeOH (10 L x 3), and the extracts were partitioned between *n*-hexane (1 L x 3) and H₂O (1 L). The *n*-hexane-soluble portions were subjected to a silica gel column (*n*-hexane / EtOAc), a Sephadex LH-20 column (MeOH), and then C₁₈ reversed-phase HPLC [LUNA 5μ C18(2), Phenomenex, 10 x 250 mm; flow rate 3.0 mL/min; UV detection at 254 nm; eluent MeOH/H₂O, 95:5] to afford biyouxanthones A (**1**, 4.2 mg, 0.00017%), B (**2**, 3.9 mg, 0.00016%), and C (**3**, 21.8 mg, 0.00087%). EtOAc-soluble portions were purified by a silica gel column (CHCl₃/MeOH) and a Sephadex LH-20 (MeOH) column chromatographies, and C₁₈ reversed-phase HPLC (Mighty sil RP-18, Kanto Chemical Co. Ltd, 10 x 250 mm; flow rate 2.0 mL/min; UV detection at 254 nm; eluent MeOH/H₂O, 9:1, 0.1% TFA) to give biyouxanthone D (**4**, 2.9 mg, 0.00012%).

Biyouxanthone A (1): yellow amorphous solid; $[\alpha]^{23}_D \pm 0$ (*c* 0.1 CHCl₃); UV (MeOH) λ_{\max} 306 (ε 3100) and 418 (4400) nm; IR (film) ν_{\max} 3347, 1671, and 1636 cm⁻¹; ¹H and ¹³C NMR data (Table 1); ESIMS *m/z* 599 [M-H]⁻; HRESIMS: *m/z* 599.3366 [M-H]⁻ (calcd for C₃₈H₄₇O₆, 599.3372).

Biyouxanthone B (2): yellow amorphous solid; $[\alpha]^{23}_D \pm 0$ (*c* 0.3 CHCl₃); UV (MeOH) λ_{\max} 260 (ε 12900), 280 (10100), 314 (13400), and 421 (9200) nm; IR (film) ν_{\max} 3367, 1668, and 1645 cm⁻¹; ¹H and ¹³C NMR data (Table 2); ESIMS *m/z* 545 [M-H]⁻; HRESIMS: *m/z* 545.2892 [M-H]⁻ (calcd for C₃₄H₄₁O₆, 545.2903).

Biyouxanthone C (3): yellow amorphous solid; $[\alpha]^{23}_D \pm 0$ (*c* 0.3 CHCl₃); UV (MeOH) λ_{\max} 250 (ε 10800), 277 (10900), and 419 (5600) nm; IR (film) ν_{\max} 3347, 1670, and 1645 cm⁻¹; ¹H and ¹³C NMR data (Table 2); ESIMS *m/z* 531 [M-H]⁻; HRESIMS: *m/z* 531.2747 [M-H]⁻ (calcd for C₃₃H₃₉O₆, 531.2750).

Biyouxanthone D (4): yellow amorphous solid; $[\alpha]^{23}_D +37.9$ (*c* 0.5 CHCl₃); UV (MeOH) λ_{\max} 254 (ε 33600), 286 (11300), and 331 (16000) nm; IR (KBr) ν_{\max} 3430, 2923, 1617, and 1591 cm⁻¹; ¹H and ¹³C NMR data (Table 3); ESIMS *m/z* 487 [M+Na]⁺; HRESIMS: *m/z* 487.2088 [M+Na]⁺ (calcd for C₂₈H₃₂O₆Na, 487.2097).

Chiral HPLC analysis of biyouxanthone B (2)

Biyouxanthone B (**2**) was subjected to a chiral HPLC [Chiralpak AD; Daicel Chemical Industry, Ltd., 4.6 x 250 mm; *n*-hexane / *i*-PrOH (99:1); flow rate 0.5 mL/min; UV detection at 254 nm] to afford (−)-biyouxanthone B (*t*_R 19.0) and (+)-biyouxanthone B (*t*_R 24.0 min). (−)-Biyouxanthone B: $[\alpha]^{21}_D$ −9.6 (*c* 0.1 CHCl₃); CD (MeOH) λ_{ext} 280 (Δε −0.96), 257 (+0.28), 237 (−0.88), 226 (−0.23), 220 (−0.49), and 211 (+4.52) nm. (+)-Biyouxanthone B: $[\alpha]^{21}_D$ +13.7 (*c* 0.1 CHCl₃); CD (MeOH) λ_{ext} 280 (Δε +0.79), 257 (−0.55), 237 (+0.63), 226 (+0.14), 220 (+0.60), and 211 (−3.67) nm.

Measurement of anti-HCV activity

Huh7 cells infected with HCV JFH-1 isolate⁶ were treated with biyouxanthones A – D (**1** – **4**) for 3 days. HCV core protein in cell culture supernatants was quantified using an enzyme immunoassay (Ortho HCV antigen ELISA Kit; Ortho Clinical Diagnostics, Tokyo, Japan), following the manufacturer's instructions.

ACKNOWLEDGMENTS

We thank S. Oka, Center for Instrumental Analysis, Hokkaido University, for measurements of HRESIMS. This work was partly supported by a grant from the Kuribayashi Foundation and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

1. A. Nahrstedt and V. Butterweck, *Pharmacopsychiatry*, 1997, **30**, 129.
2. M. Dostalek, J. Pistovcakova, J. Jurica, J. Tomandl, I. Linhart, A. Sulcavá, and E. Hadasova, *Life Sciences*, 2005, **78**, 239.
3. M. A. Medina, B. Mrtínez-Proveda, M. I. Amores-Sánchez, and A. R. Quesada, *Life Sciences*, 2006, **79**, 105.
4. L. Beerhues, *Phytochemistry*, 2006, **67**, 2201.
5. N. Tanaka, T. Kubota, H. Ishiyama, Y. Kashiwada, Y. Takaishi, J. Ito, Y. Mikami, M. Shiro, and J. Kobayashi, *Heterocycles*, 2009, **79**, 917.
6. T. Wakita, T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H. G. Krausslich, M. Mizokami, R. Bartenschlager, and T. J. Liang, *Nat. Med.*, 2005, **11**, 791.