


INTRODUCTION OF ALL FUNCTIONAL GROUPS FOUND IN THE BENZENE PART OF MITOMYCIN C TO PYRROLO[1,2-*a*]INDOLE DERIVATIVE

Satoshi Hirano, Yoshihiko Shinoda and Shin-ichi Nakatsuka*

The United Graduate School of Agricultural Science, Gifu University,
Yanagido, Gifu 501-11, Japan

Abstract : All functional groups on the benzene part of mitomycin C were directly introduced on pyrrolo[1,2-*a*]indole 2 by Friedel-Crafts alkylation, oxidative introduction of *p*-quinone moiety etc.

Introduction of substituents on the benzene part (4~7 position) of indole ring is one of the most difficult problems in the organic syntheses (1). We have developed several useful methods to resolve the problem and applied those to the syntheses of natural products (2~4). We have been reported efficient method for the synthesis of indoloquinone (4b) and applied them for the synthesis 7-methoxymitosene starting from 6-methylindole (4d, e). In this paper, we report the synthesis of pyrroloindoloquinone 3 by direct introduction of the functional groups found in mitomycin C 1 (5~7) from simple pyrrolo[1,2-*a*]indole 2.




Although Friedel-Crafts methylation [$\text{CH}_3\text{Br}/\text{AlCl}_3$, $(\text{CH}_3)_2\text{SO}_4/\text{AlCl}_3$ etc.] of pyrrolo[1,2-*a*]indole 2 gave no methylated product, alkylation with $\text{ClCH}_2\text{SCH}_3/\text{AlCl}_3$ in CH_2Cl_2 at -20°C for 30min gave monomethylthiomethyl derivatives 4 and 5 in 83% yield(1:1). Those could not be separated on silica gel TLC or column, but were separated with HPLC[Namsil NA-10 (8), 3% AcOEt in Hex] in small scale and those structures were determined as 7-substituted 4 and 6-substituted 5 by comparison of those $^1\text{H-NMR}$ -spectra. 4; ^1H NMR (CDCl_3) δ (ppm) 2.00 (3H, s), 2.66 (2H, m), 2.29 (2H, t, $J=7.6\text{Hz}$), 3.83 (2H, br.s), 3.91 (3H, s), 4.11 (2H, t, $J=7.2\text{Hz}$), 7.22 (2H, m), 7.97 (1H, s). 5; ^1H NMR (CDCl_3) δ (ppm) 2.00 (3H, s), 2.66 (2H, m), 2.29 (2H, t, $J=7.6\text{Hz}$), 3.82 (2H, br.s), 3.90 (3H, s), 4.11 (2H, t, $J=7.2\text{Hz}$), 7.16 (1H, d, $J=7.3\text{Hz}$), 7.22 (1H, s), 8.02 (1H, d, $J=7.3\text{Hz}$).

Reduction of a mixture of 4 and 5 with Raney Ni in methanol at 25°C gave corresponding methyl derivatives 6 and 7, in 98% yield. Separation of 6 (9) and 7 (4d) were also difficult in large scale, but each structure was determined after separation with HPLC(ODS-3, 50% MeOH in H_2O) and comparison with authentic samples.

A mixture of 6 and 7 were nitrated with sodium nitrate in the presence of 2.5% conc. H_2SO_4 in acetic acid at 40°C for 50min to afford mono-nitro-derivatives 8 and 9, which were easily separated on silica gel column (14% and 22% yields respectively). 8; mp 192°C , ^1H NMR (CDCl_3) δ (ppm) 2.42 (3H, br.s), 2.68 (2H, m), 3.32 (2H, t, $J=7.6\text{Hz}$), 3.82 (3H, s), 4.16 (2H, t, $J=7.3\text{Hz}$), 7.07 (1H, d, $J=8.5\text{Hz}$), 7.26 (1H, d, $J=8.3\text{Hz}$). 9; mp $178\text{--}179^\circ\text{C}$, ^1H NMR (CDCl_3) δ (ppm) 2.50 (3H, br.s), 2.69 (2H, m), 3.31 (2H, t, $J=7.6\text{Hz}$), 3.82 (3H, s), 4.15 (2H, t, $J=7.2\text{Hz}$), 7.26 (1H, s), 7.39 (1H, s). Compounds 8 and 9 were also obtained by nitration of purified 7-methyl derivative 6 and 6-methyl derivative 7, respectively, and those structures were easily determined by their $^1\text{H-NMR}$ -spectra. In both case, the nitrating position of 6 and 7 was favoured at the 8-position (3a, d, e).

Hydrogenation of compound 9 with $\text{H}_2/\text{Pd-C}$ in methanol gave a 8-amino derivative in 90% yield and subsequent oxidation of the amino-derivative with Fremy's salt [$\bullet\text{ON}(\text{SO}_3\text{K})_2$] gave *p*-quinone 10 (10) in 56% yield. Amination of 10 with benzylamine afforded 3 (11) which contains all functional groups on the benzene part of mitomycin C 1. Thus we developed novel method directly introducing all substituents, found in benzene part of mitomycin C, on the benzene part of pyrrolo[1,2-*a*]indole 2 in 6 steps(4.8% over all yield).



Reagents: a) $\text{ClCH}_2\text{SCH}_3$, AlCl_3 (4 and 5, 83%); b) Raney Ni (6 and 7, 98%); c) NaNO_3 , H_2SO_4 (8, 22%); d) H_2 , Pd-C (90%); e) $\text{ON}(\text{SO}_3\text{K})_2$, Phosphate Buffer (56%); f) BnNH_2 , pyridine (53%).

REFERENCES AND FOOTNOTES

- (1) R. J. Sundberg, 'The Chemistry of Indole,' published by Academic press, 1970
- (2) a) S. Nakatsuka, H. Miyazaki and T. Goto, *Tetrahedron Lett.* 21, 2817 (1980) ; b) S. Nakatsuka, H. Miyazaki and T. Goto, *Chemistry Lett.* 407 (1981) ; c) S. Nakatsuka, K. Yamada and T. Goto, *Tetrahedron Lett.* 27, 4757 (1986)
- (3) a) S. Nakatsuka, T. Masuda, O. Asano, T. Teramae and T. Goto, *Tetrahedron Lett.* 27, 4327 (1986); b) S. Nakatsuka, T. Masuda, K. Sakai and T. Goto, *ibid.* 27, 5735 (1986) ; c) S. Nakatsuka, T. Masuda and T. Goto, *ibid.* 27, 6245 (1986) ; d) T. Masuda, K. Ueda, O. Asano, S. Nakatsuka and T. Goto, *Heterocycles* 26, 1475 (1987) ; e) S. Nakatsuka, T. Masuda and T. Goto, *Tetrahedron Lett.* 28, 3671 (1987)
- (4) a) S. Nakatsuka, O. Asano and T. Goto, *Heterocycles* 24, 2109 (1986) ; b) S. Nakatsuka, K. Ueda, O. Asano and T. Goto, *ibid.* 26, 65 (1987) ; c) S. Nakatsuka, O. Asano, K. Ueda and T. Goto, *ibid.*, 26, 1471 (1987) ; d) O. Asano, S. Nakatsuka and T. Goto, *ibid.* 26, 1207 (1987) ; e) S. Nakatsuka, O. Asano and T. Goto, *Chemistry Lett.* 1225 (1987) ; f) S. Nakatsuka, O. Asano and T. Goto, *Heterocycles* 26, 2603 (1987)
- (5) a) T. Hata, Y. Sano, R. Sugiura, A. Matsumae, K. Kanamori, Y. Shima and T. Hoshi, *J. Antibiotics* 9, 141 and 146 (1956) ; b) S. Wakaki, K. Marumo, G. Tomioka, E. Shimazu, H. Kato, H. Kamata, S. Kudo and Y. Fujimoto, *Antibiotics and Chemotherapy* 8, 228 (1958) ; c) J. S. Webb, D. B. Cosulich, J. H. Mowat, J. B. Patrick, R. W. Broshard, W. E. Meyer, R. P. Williams, C. F. Wolf, W. Fulmor, C. Pidacks and J. E. Lancaster, *J. Am. Chem. Soc.* 84, 3185 (1962) ; d) A. Tulinsky, *ibid.* 84, 3188 (1962) ; e) K. Shirahata and N. Hirayama, *J. Am. Chem. Soc.* 105, 7199 (1983) ; f) M. Kasai and M. Kono, *SynLett* 778 (1992)
- (6) a) Y. Kishi, *J. Natural Prod.* 42, 549 (1979) and references cited therein b) T. Fukuyama and L. Yang, *J. Am. Chem. Soc.* 111, 8303 (1989)
- (7) References cited in the previous paper^{4c} and a) P. A. Wender and C. B. Cooper, *Tetrahedron Lett.* 28, 6125 (1987) ; b) Y. Naruta, N. Nagai and K. Maruyama, *J. Chem. Soc. Perkin Trans. I* 1143 (1988) ; c) J. W. Benbow, K. F. McClure and S. J. Danishefsky, *J. Am. Chem. Soc.* 115, 12305 (1993)
- (8) Packed column of 10 μ normal phase silicagel, Nam Res. Lab.
- (9) 6; ^1H NMR (CDCl₃) δ (ppm) 2.47 (3H, br.s), 2.64 (2H, m), 3.28 (2H, t, J=7.6Hz), 3.89 (3H, s), 4.09 (2H, t, J=7.0), 7.05 (1H, d, J=8.5Hz), 7.14 (1H, d, J=8.2Hz), 7.90 (1H, s)
- (10) 10; mp 172~173°C(decomp.), ^1H NMR (CDCl₃) δ (ppm) 2.05 (3H, d, J=1.8Hz), 2.61 (2H, m), 3.14 (2H, t, J=7.6Hz), 3.88 (3H, s), 4.33 (2H, t, J=7.5Hz), 6.46 (1H, d, J=1.3Hz)
- (11) 3; ^1H NMR (CDCl₃) δ (ppm) 2.08 (3H, br.s), 2.56 (2H, m), 3.09 (2H, t, J=7.6Hz), 3.85 (3H, s), 4.31 (2H, t, J=7.3Hz), 4.71 (2H, d, J=6.1Hz), 6.30 (1H, br.s), 7.26~7.35 (5H, m)

Received November 24, 1994