

REACTIVITY OF INDOLE AS AN ELECTROPHILE AND ITS APPLICATION FOR THE SYNTHESIS OF 2-PHENYLINDOLE AND 3-PHENYLINDOLE

Nobumitsu Tajima and Shin-ichi Nakatsuka*


United Graduate School of Agricultural Science, Gifu University,
1-1 Yanagido, Gifu 501-1193, Japan

Abstract: Reactivity of *N*-acylindole **3a~c** as an electrophile was studied and phenyl group was introduced at the 2 and 3-position of indole nucleus to afford phenylindoline derivatives **4a~c** and **5a~c**, which were derived to 2-phenylindole **8** and 3-phenylindole **9**.

Indole is one of the most fascinating compounds and its chemistry had been studied for a century (1). Since the pyrrole part of indole nucleus is electron rich, nucleophilic reactivity of the 1~3 positions plays the major part of indole chemistry (1a). On the other hand, reactivity of indole as an electrophile has also been reported in polymerization of indole to give dimer *etc.* reported by Hodson and Smith (2) as shown in Scheme 1. The reaction mechanism could be realized by the key step that electron deficient 2-position of protonated cation **I** is attacked by electron rich 3-position of non-protonated **1**. Here we report novel route to introduce phenyl group at 2 and 3-position of *N*-acylindole.

We started our synthetic study on electron deficient indole (3) such as **I** type and chose *N*-acylindole **3a~c** as the model compounds. Nucleophilic addition of benzene to electron deficient **3a~c**-AlCl₃ complex **II** prepared in CH₂Cl₂ was easily achieved as summarized in Table 1. When *N*-acetylindole **3a** was treated with 3~5 equiv. AlCl₃ in CH₂Cl₂ solution


Scheme 1.




containing 1.2 equiv. benzene at r.t., 3-phenylindoline derivative **5a** (80%) and 2-phenylindole derivative **4a** (8%) were obtained. In the case of *N*-trimethylacetylindole **3b**, no 2-phenylindoline derivative **4b** was produced in the reaction mixture. In the same reaction condition, *N*-benzoylindole **3c** provided about 1 : 1 mixture of **4c** (4) and **5c** (5) in 82% yield. Thus we could readily introduce a phenyl group at the 2 and 3-position of *N*-acylindole **3a-c** to give phenylindoline **4a-c** and **5a-c** in good yields.

2-phenylindole **8** and 3-phenylindole **9** are usually prepared by E. Fischer's indole synthesis (6), or other methods (7) starting from simple benzene derivative such as aniline. Recently, Yamanaka and Sakamoto (8) reported new route for the synthesis of **8** and **9** starting from protected indoles *via* its carbanion of 2 or 3-position, but no synthetic method of **8** and **9** utilizing electrophilic reactivity of indole nucleus has been reported.

A 1 : 1 mixture of 2 and 3-phenylindoline derivatives **4c** and **5c** was hydrolyzed with 6N HCl to give 2-phenylindoline **6** and 3-phenylindoline **7** in 89% yield. Separation of **6** and **7** was easily carried out by chromatography on silica gel PTLC plate (solvent: 10% EtOAc in hexane). Oxidation of **6** or **7** in dichloroethane with MnO_2 at 60 °C afforded 2-phenylindole **8** (9) and 3-phenylindole **9** (10) in 87% and 72% yield, respectively. Their m.p. and spectroscopic data were completely identical with reported data. Consequently, we could establish a novel and convenient synthetic route of 2-phenylindole **8** and 3-phenylindole **9** from indole.

Table 1. Electrophile addition of **3a-c**

Starting Materials 3	Products 4 5	
3a : $\text{R} = \text{CH}_3$	4a (8%)	5a (80%)
3b : $\text{R} = \text{C}(\text{CH}_3)_3$	4b (0%)	5b (81%)
3c : $\text{R} = \text{C}_6\text{H}_5$	4c (39%)	5c (43%)



REFERENS AND FOOTNOTES

- (1) a) R. J. Sundberg, *The Chemistry of Indoles*, published by Academic press, 1970.
b) H. G. Floss, *Tetrahedron*, **32**, 873 (1976).
- (2) H. F. Hodson and G. F. Smith, *J. Chem. Soc.*, 3544 (1957).
- (3) a) K. Teranishi, S. Hayashi, S. Nakatsuka and T. Goto, *Tetrahedron Lett.*, **35**, 8173 (1994) and *Synthesis*, 506 (1995).
c) S. Nakatsuka, T. Hayashi, S. Adachi, Y. Harada and N. Tajima, *Heterocyclic Commun.*, **3**, 47, (1997).
- (4) **4c**; MS m/z 299 (M^+), 105. 1H -nmr ($CDCl_3$) δ ppm 2.98 (1H, d.d., $J=2$ & 16 Hz), 3.74 (1H, d.d., $J=10$ & 16 Hz), 5.44 (1H, br.s), 6.99~7.08 (3H, m), 7.17~7.43(10H,m), 8.26 (1H, extremely br.s).
- (5) **5c**; MS m/z 299 (M^+), 105. 1H -nmr ($CDCl_3$) δ ppm 4.01 (1H, br.s), 4.45 (1H, br.s), 4.56(1H, br.t, $J=9$ Hz), 7.01 (2H,m), 7.18~7.20 (3H,m), 7.26 (1H,m), 7.32 (2H,m), 7.41~7.46 (2H, m), 8.27 (1H, extremely br.s).
- (6) a) E. Fischer and T. Schmidt, *Ber.*, **21**, 1071 (1888).
b) E. Fischer and T. Schmidt, *Ber.*, **21**, 1811 (1888).
- (7) a) R. L. Schiner, W. C. Ashley and E. Welch, *Org. Synth.*, **22**, 98 (1942).
b) G. Smolinsky and C. A. Pryde, *J. Org. Chem.*, **33**, 2411 (1968).
c) P. G. Gassman and T. J. van Bergen, *J. Am. Chem. Soc.*, **95**, 590 (1973)
d) I. Fleming and M. Woolias, *J. Chem. Soc. Perkin Trans. 1*, 829 (1979).
e) K. Iritani, S. Matsubara and K. Utimoto, *Tetrahedron Lett.*, **29**, 1799 (1988).
f) Y. Tsuji, S. Kotachi, K-T Huh and Y. Watanabe, *J. Org. Chem.*, **55**, 580 (1990).
g) N. Kato and N. Miyaura, *Tetrahedron*, **52**, 13347 (1996).
- (8) T. Sakamoto, Y. Kondo, N. Takazawa and H. Yamanaka, *J. Chem. Soc. Perkin Trans. 1*, 1927 (1996).
- (9) **8**; m.p. 189~190°C (lit.,^{7a} 188~189°C), MS m/z 193 (M^+). 1H -nmr ($CDCl_3$) δ ppm 6.83 (1H, br.s), 7.13 (1H, br.t, $J=8$ Hz), 7.20 (1H, br.t, $J=8$ Hz), 7.34 (1H, br.t, $J=8$ Hz), 7.44~7.50 (3H, m), 7.63 (1H, br.d, $J=8$ Hz), 7.68 (2H, m), 8.36 (1H, br.s).
- (10) **9**; m.p. 86.5~87°C (lit.,^{6b} 88~89°C), MS m/z 193 (M^+). 1H -nmr ($CDCl_3$) δ ppm 7.20 (1H, br.t, $J=8$ Hz), 7.26 (1H, br.t, $J=8$ Hz), 7.29 (1H, br.t, $J=8$ Hz), 7.38 (1H, d, $J=3$ Hz), 7.43~7.47 (3H, m), 7.68 (2H, br.d, $J=8$ Hz), 7.95 (1H, br.d, $J=8$ Hz), 8.25 (1H, br.s).

Received on October 30, 1999