

# HETEROCYCLIC COMMUNICATIONS

## An International Journal in Heterocyclic Chemistry

Published bimonthly by Freund Publishing House Ltd.,  
Suite 500, Chesham House, 150 Regent Street, London W1R 5FA, England

### Notes for the preparation of Camera-Ready Manuscripts

**Heterocyclic Communications** seeks to publish preliminary communications and full length research papers on all phases of heterocyclic chemistry including inorganic ring systems. All papers to the journal will be reproduced from author's typescripts (camera-ready manuscript) by a reduction of 25%.

#### Guidelines for Authors

##### 1. Preparation and Mailing of Manuscripts

Author(s) should mail manuscript to respective editor in triplicate. All manuscript should be made on a computer and printed out by a laser printer preferably at 600 dpi or higher resolution. Author(s) should send the editor a floppy disc (31/2" or 51/4") in DOS format (1.4MB for 31/2" and 1.25MB for 51/4") which contains a combined manuscript file including computer drawn structural formulas, figures, tables etc. For the all type setting, Times Roman 10 Pt. font or corresponding font set should be used. Preferably, file format of the manuscript is Adobe Page Maker 6.5/6.0/5.0 or MS Word 97 versions for MS Windows. Structures may be drawn by programs like Chem Draw, ISIS Draw, Chem Win, Chem Window/Coral Draw, Chem Text. Apple Macintosh and other word processor users should convert the file format to those readable by these softwares. Some times RTF (Rich Text Format) and PDF (Portable Document Format) are effective. Details of the program(s) used should be also provided. If structures/diagrams have been prepared using program(s) other than those stated, then structures/diagrams should be converted into windows meta file format. Details of the program(s) used to draw structures/diagrams should be also provided. Manuscript should end with even numbered page (viz 2,4,6....). Normally manuscript not exceeding 6 pages (excluding graphical abstract) will be accepted. Editor can accept even manuscript exceeding 6 pages, if they feel that it is appropriate to do so.

All manuscripts except from North America, European countries and Japan should be mailed to Dr. R.R. Gupta, Editor-in-Chief, Heterocyclic Communications, 10A, Vasundhara Colony, Tonk Road, Jaipur-302018, India. Authors from Europe should mail manuscripts to Prof. Jacques Barbe, Editor for European Countries, Faculte de Pharmacie, 27, Bd. Jean-Moulin, 13385 Marseille Cedex 5, France. Authors from North America should mail manuscripts to Prof. L. Strekowski, Department of Chemistry, Georgia State University, University Plaza, Atlanta, Georgia 30303-3083, USA. Authors from Japan should mail manuscripts to Prof. S. Eguchi, Department of Molecular Design and Engineering, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8603, Japan.

##### 2. Organization of the Manuscript

**2.1 Title :** It should be clear, concise, and informative. It should be typed with words all in bold face capitals, single spaced, centered on the width of the first page, 2 cm below the top of typing area.

**2.2 Author(s) :** After 4 single spaces from the title, authors names and addresses should be given on separate lines, typed single spaced, centered on the width of the first page. The name of the author to whom correspondence should be directed should be marked with an asterisk as superscript.

**2.3. Abstract :** After 4 single spaces from authors address, a short abstract (50-100 words) describing results, methodology employed, conclusions etc., should be placed. It should be labelled as the abstract with the word "Abstract" in bold face and should be typed single spaced. Total typing area on the first page should be 17x22 cm. Typing area for other pages should be 17x24 cm. Each manuscript should be accompanied by a Graphical Abstract, which should contain the title in bold face single spaced, authors names and addresses. It should summarise concisely the contents of the communications in pictorial form designed to capture the attention of a wide readership and to facilitate compilation of database. Chemical Structures designed to illustrate the theme of research work presented should be included. It should be prepared on a separate sheet of paper sized 19x7 cm. An example of graphical presentation is shown as :

## Graphical Abstract

|       |             |      |
|-------|-------------|------|
| Space | Leave blank | 1 cm |
|       | 10 cm       |      |

For

Abstract

**2.4 Text :** Text should be divided into sections like **Introduction, Experimental**, including spectra screening, instruments specifications, **Results, Discussions, Conclusions, Acknowledgements** (if applicable) and **References**. However, the experimental section should be excluded in preliminary Communications as far as possible. Between the end of one section and the beginning of another, there should be a gap of three single spaces. Text should be typed in one and a half spacing throughout.

**2.5 References:** These should be numbered consecutively throughout the text as they appear and should be superscripted. All compound number in text should be bold e.g., phenothiazines **7** can be prepared... . Abbreviations for journals should be those used in Chemical Abstracts. Abbreviations for other terms should also be taken from Chemical Abstracts/ACS journal. Each reference should be typed single spaced. However, there should be one empty line between two consecutive references. Some examples for journal and books references are given:

- 1 R.R. Gupta, V. Saraswat, A. Gupta, M. Jain and V. Gupta, J. Heterocycl. Chem. **29**, 1703 (1992)
- 2 R.R. Gupta, Diamagnetic Susceptibility, Springer Verlag, Berlin, 1986
- 3 R.R. Gupta (Ed.), Physical Methods in Heterocyclic Chemistry, John Wiley, New York, 1984
- 4 R.R. Gupta and M. Kumar, Synthesis, Properties and Reactions of Phenothiazines, in R.R. Gupta (Ed.), Phenothiazines and 1,4-Benzothiazines-Chemical and Biomedical Aspects, Elsevier, Amsterdam, 1988, pp.1

### 2.6 Illustrations

Both line drawing and photographs should be used sparingly. All illustrations must be numbered in sequence using arabic numbers, and each illustration must be referred to in the text. Each illustration must be provided with a caption which is complete enough for the figure to be appreciated without referring to the text. Line drawings must be submitted as the perfectly drawn original or as a glossy photograph print. Photocopies, Multiliths, Verifax or Xerox copies are not acceptable substitutes. If you are using desktop publishing software with high quality graphics capability, which is recommended, it is desirable that the graphics be sized and printed along with the text. A laser printer should be used. Photographs must be supplied as glossy prints in which any lettering must be part of the photographs. No lettering should be applied to the surface of the photograph. Micrographs and similar material in which linear dimensions are important should have a scale of length.

### 2.7 Tables

Tabular presentation of data is an economic way of condensing many items. Refer to tables by using Latin numerals in text and to head the table itself. Provide each table with a caption which will allow it to be read without reference to the text. The table caption should be typed to the width of the table itself. Exceptionally large tables may be placed landscape on the page with the top of table at the left-hand margin.

### 2.8 Page Numbers

When your typescript is complete and in its final form, number each page sequentially from 1, using a light blue pencil, in the top right-hand corner of each sheet. Final page numbers will be inserted by the editor or the publisher.

### 2.9 Layout

The typing area should be 17x24 cm for all pages except the first. On the first page typing should start 2 cm below top of the typing area, e.g., its typing area should be 17x22 cm. After abstract text should be typed in one and a half spacing throughout on all pages. Photographic reduction will be by 25%.

### 2.10 Displayed Equations and Formulas

Leave a blank line above and below all display. Break equations, where possible, only at an equals sign (=) or equivalent (>, <, ≥, ≤). Do not repeat equations sign at the end of one line and the beginning of the next, but carry it over to the beginning of the following line.

# **REPRINT ORDER FORM FOR HETEROCYCLIC COMMUNICATIONS**

**FREUND PUBLISHING HOUSE LTD.**  
P.O.Box 35010, Tel Aviv, Isreal, 61350

**Telephone: 972-3-562-8540**  
**Fax: 972-3-562-8538**  
**E-mail: h\_freund@netvision.net.il**

Author of the manuscript: .....

Title of the manuscript: .....

**TWENTY-FIVE** reprints will be supplied free of charge for each **ARTICLE**.  
**(PLEASE NOTE: NO FREE REPRINTS ARE SUPPLIED FOR BOOKS.)**  
Reprints may be purchased in units of 100 only.

## **COST OF REPRINTS**

| No. of copies | Number of Pages |        |        |         |          |                            |
|---------------|-----------------|--------|--------|---------|----------|----------------------------|
|               | 1-2 pp          | 3-4 pp | 5-8 pp | 9-12 pp | 13-16 pp | Each Add.<br>Unit (1-4 pp) |
| 100           | \$ 83           | 122    | 159    | 253     | 341      | 111                        |
| 200           | \$ 109          | 150    | 259    | 380     | 516      | 136                        |
| 300           | \$ 128          | 245    | 380    | 569     | 759      | 172                        |

Orders for quantities of over 300 copies are subject to special quotation.

- \* **IMPORTANT: No reprints will be sent unless this form is filled in and signed.**
- \* Type may be destroyed after the printing of the issue.
- \* Please note that payment is to be received in U.S. Dollars.
- \* Co-author do not receive copies of this form.
- \* Author copies of the journal and reprints will be sent by airmail.

## **\*\*PLEASE TYPE\*\***

Please supply me with ..... reprints, in the addition to the 25 free copies to which I am entitled.

NAME.....

ADDRESS.....

TELEPHONE..... FAX.....

E-MAIL.....

INVOICE TO:.....

PRICE OF REPRINTS:.....

- ☐ I enclosed a check (made payable to FREUND PUBLISHING HOUSE LTD.) in the amount of \$.....
- ☐ Please debit my: VISA/AMERICAN EXPRESS/MASTER CARD for the amount of \$.....

Number..... Expiry Date.....

# Index of Vol. 6 (Issues 1-6), 2000

## Heterocyclic Communications, Vol. 6, No. 1, 2000

### Vol. 6, Issue 1 (2000)

|                                                                                                              |    |                                                                                                                                                      |
|--------------------------------------------------------------------------------------------------------------|----|------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                              | 1  | Contributors to this issue.                                                                                                                          |
|                                                                                                              | 2  | Graphical Abstracts.                                                                                                                                 |
| L. Strckowski, Y. Gulevich and K.V. Aken                                                                     | 9  | A novel synthesis of 3-[(methylamino)methyl]-6-oxo-3,4,5,6-tetrahydro-1 <i>H</i> -azepino[5,4,3- <i>cd</i> ]indole.                                  |
| G.M. Coppola, Y.L. Zhang and H.F. Schuster                                                                   | 13 | Isoquinoline-based amino acid derivatives.                                                                                                           |
| D. Geffken, C. Hoist and I. Willrodt                                                                         | 21 | Synthesis of oxazolo[4,3- <i>c</i> ]1,2,4-triazol-5-ones.                                                                                            |
| E. Barbu, J.J. Wolff, I. Bolocan and F. Cuiban                                                               | 25 | Valerian alkaloids: First total synthesis of a naphthyridyl methyl ketone.                                                                           |
| J.C. DiCesare and B.M. White                                                                                 | 29 | Large scale reductive cleavage of dibenzothiophene.                                                                                                  |
| K. Matsumoto and S. Kimura                                                                                   | 31 | Diels-Alder reaction of Ni(II) $\beta$ -vinyl-meso-tetraphenylporphyrin with dimethyl acetylenedicarboxylate and <i>N</i> -phenyl maleimide.         |
| I.V. Tselinsky, S.F. Melnikova, S.V. Pirogov, S.D. Shaposhnikov, C. Näther, T. Trauisen and F. Friedrichsen  | 35 | On the interaction of 3,4-dibenzoylfuroxan with hydroxylamine.                                                                                       |
| R.M. Srivastava, L.P.F. Morais, M.T.J.A. Catanho, G.M.L. Souza, G.M. Seabra, A.M. Simas and M.A.L. Rodrigues | 41 | Synthesis and antiinflammatory activity of 3-aryl-5-isopropyl-1,2,4-oxadiazoles.                                                                     |
| V.L.M. Guarda, M. Perrissin, F. Thomasson, E.A. Ximenes, S.L. Galdino, I.R. Pitta and C. Luu-Duc             | 49 | Synthesis and microbiological activity of some 2 <i>H</i> -benzothiazin-3-one derivatives.                                                           |
| D. Geffken and S. Zilz                                                                                       | 55 | 4-Aminosubstituted 1,2,4-oxadiazolidin-3,5-diones.                                                                                                   |
| N. Tajima and S.-i. Nakatsuka                                                                                | 59 | Reactivity of indole as an electrophile and its application for the synthesis of 2-phenylindole and 3-phenylindole.                                  |
| G.A. Prakash and S.P. Rajendran                                                                              | 63 | Synthesis of 1,2,3,4-tetrahydrodibenzo[ <i>b,g</i> ][1,8]naphthyridines.                                                                             |
| M. Eckert-Maksić, V. Rapić, D. Margetić and V. Matonićkin                                                    | 67 | Ionic hydrogenation of 4-hydroxycoumarin derivatives.                                                                                                |
| O.N. Burchak, A.M. Chibiryayev and A.V. Tkachev                                                              | 73 | Terpenic methyl ketones from limonene, (+)-3-carene and $\alpha$ -pinene in Fischer's indole synthesis.                                              |
| L. Wenjian, S. Xuejun, Y. Jinmao and O. Qingyu                                                               | 81 | Microwave-assisted preparation of substituted carbazole-9-acetic/propionic acid and their absorption and fluorescence spectroscopic characteristics. |
| V. Peesapati and K. Anuradha                                                                                 | 87 | Studies of Vilsmeier-Haack reaction : Formation of some heterocyclic annelated compounds from 6,7,8,9-tetrahydrobenzocyclohepten-5-ones.             |
| K. Saidi, H. Shaterian and D. Aghaei                                                                         | 93 | A one-pot synthesis of 1,3-dioxazine-4-one derivatives.                                                                                              |
|                                                                                                              | 97 | Notes for the preparation of manuscript.                                                                                                             |

# Heterocyclic Communications, Vol. 6, No. 2, 2000

## Vol. 6, Issue 2 (2000)

|                                                                                               |     |                                                                                                                                                                                          |
|-----------------------------------------------------------------------------------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                               | 99  | Contributors to this Issue                                                                                                                                                               |
|                                                                                               | 100 | Graphical Abstracts                                                                                                                                                                      |
| J. Tierny, D. Sheridan, K. Kovalsky,<br>L.M. Mascavage, K. Schneider and<br>C. Sheely         | 105 | Substituent effects for some substituted 3-benzyl-2-phenyl-1,3-thiazolidin-4-ones using $^1\text{H}$ and $^{13}\text{C}$ NMR.                                                            |
| M.J. Uddin, T. Fujimoto, A. Kakehi,<br>H. Shirai and I. Yamamoto                              | 113 | Diastereoselective synthesis of bridgehead heterocyclic spiro compounds derived from Tandem Michael intramolecular 1,3-dipolar cycloaddition of nitrones.                                |
| E. Michelena-Baez,<br>A.M. Navarro-Eisenstein, H.D. Banks,<br>P.C. Vasquez and A.L. Baumstark | 119 | Epoxidation of aliphatic alkenes by dimethyldioxirane: Steric effects                                                                                                                    |
| C. Locher                                                                                     | 123 | Synthesis of some bis( <i>N</i> -1,2,3,4-tetrahydroisoquinoliny)methanes using benzotriazole as auxiliary.                                                                               |
| V. Rogers, S. Mendonca, T.W. Packham<br>and M.J. Davies                                       | 133 | Synthetic approaches to diazabiphenylenes: Synthesis of the carbon-nitrogen skeleton of 1,5- and 2,6-diazabiphenylene.                                                                   |
| O.V. Mikhailov, A.I. Khamitova and<br>V.I. Morozov                                            | 137 | Template synthesis of cobalt(II,III), nickel(II) and copper(II) macrocyclic compounds with 2,8-dithio-3,7-diaza-5-oxanonandithioamide-1,9 in gelatin-immobilized matrix.                 |
| N. Tajima, Y. Shinoda and<br>S.-i. Nakatsuka                                                  | 143 | Biomimetic intramolecular cyclization of 1-trimethylacetyl-3-(3-methanesulfonylbutyl) indole at the 4-position of indole nucleus.                                                        |
| D. Csanyi, G. Hajos, Z. Riedl,<br>O. Egyed, C. Kos and T. Kappe                               | 147 | Synthesis of new tricyclic heteroaromatic ring system: Pyridazo[3,2- <i>b</i> ]-quinazoline.                                                                                             |
| B. Balamurali and K.J.R. Prasad                                                               | 153 | A new synthesis of 3,4,5-trihydroazepino[2,3- <i>b</i> ]indoles.                                                                                                                         |
| M. Petroligi, O. Iglessi-Markopoulou,<br>and J. Markopoulos                                   | 157 | Complexation and spectroscopic studies of 5-benzylidene-3-hexanoyl tetramic acid with magnesium (II), zinc (II) and barium (II) ions.                                                    |
| G. Yang and H. Yang                                                                           | 165 | Regioselective addition of 5-amino-1,2,4-triazoles and its analogues with arylsulfonyl isocyanates.                                                                                      |
| K. Shanmugasundaram and<br>K.J.R. Prasad                                                      | 169 | A free radical initiated dimerisation in the synthesis of 4,5-bis(2,4-diacetylcarbazol-1-yl) 1,4,5-oxdiazepanes.                                                                         |
| F.H. Osman and F.A. El-Samahy                                                                 | 175 | The reaction of isatin with alkoxycarbonylmethylene(triphenyl)-phosphoranes in acetic anhydride.                                                                                         |
| A. Dandia, H. Sachdeva, N. Ahmed<br>and K.C. Joshi                                            | 181 | One pot synthesis of fluorine containing diastereoisomeric spiro [3H-indol-3,2'-oxiran]-2(1H)-ones and their conversion to 5a,10b-dihydro-5H,6H-indolo-[2,3- <i>b</i> ]quinolin-11-ones. |
|                                                                                               | 189 | Notes for the preparation of manuscript.                                                                                                                                                 |

# Heterocyclic Communications, Vol. 6, No. 3, 2000

## Vol. 6, Issue 3 (2000)

|                                                                                    |     |                                                                                                                                                                                                                                                                             |
|------------------------------------------------------------------------------------|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                    | 191 | Contributors to this Issue                                                                                                                                                                                                                                                  |
|                                                                                    | 192 | Graphical Abstracts                                                                                                                                                                                                                                                         |
| Y. Satyanarayana and J.W. Lown                                                     | 199 | Design and synthesis of a new class of pyrrolobenzimidazole based agents to target human tumor helicases.                                                                                                                                                                   |
| J.B. Bremner, A.A. Monteith and R.J. Smith                                         | 209 | Free-radical cyclization of 2-( <i>o</i> -bromobenzoyl)-1-(2-propenyl)-1,2,3,4-tetrahydroisoquinoline: Synthesis of 7-methyl-6,7,8,9-tetrahydro-5 <i>H</i> -benzocyclohepten-5,8-imine.                                                                                     |
| H. Konda, T. Oritani and H. Kiyota                                                 | 211 | Synthesis and antimicrobial activity of unnatural (-)-antimycin A <sub>3</sub> and its analog.                                                                                                                                                                              |
| N. Roue and R. Barret                                                              | 215 | Approach of pyrrolo[4,3,2- <i>de</i> ]quinoline.                                                                                                                                                                                                                            |
| C. Socaci, I. Grosu, G. Pię, H.W. Zwanziger, E. Mesaros, D. Marginean and S. Mager | 219 | Synthesis and stereochemistry of some new chiral brominated 1,3-dioxane derivatives.                                                                                                                                                                                        |
| E. Gavrielatos, C. Mitsos, J. Markopoulos, M. Petroligi and O. Iglessi-Markopoulou | 225 | Synthesis and nmr spectroscopic studies of novel N-acetyl-3-hydrazonoalkyl tetramic acids.                                                                                                                                                                                  |
| B. Insuasty, R. Rodriguez, J. Quiroga, R. Abonia, C. Saitz and C. Jullian          | 231 | Synthesis of new 1-benzyl-4,6-diaryl-2,3-dihydropyrazolo[3,4- <i>b</i> ][1,4]-diazepines.                                                                                                                                                                                   |
| A. Krauze and G. Duburs                                                            | 239 | Synthesis, dehydration and oxidation of 3-cyano-4,6-diaryl-5-ethoxycarbonyl-6-hydroxypiperidine-2-thiones.                                                                                                                                                                  |
| K. Matsumoto, S. Goto, M. Toda and A. Kakehi                                       | 245 | An extremely unusual reaction of 2,3,4-triphenyl-3-azabicyclo-[3.2.0]hepta-1,4-diene with dimethyl acetylenedicarboxylate.                                                                                                                                                  |
| M.F. Aly, A.H. Atta and M.I. Younes                                                | 249 | Reaction of 1-acetyl-3-dicyanomethylene-1,3-dihydro-2 <i>H</i> -indol-2-one with some nucleophilic reagents : Synthesis of some indol and quinoline derivatives.                                                                                                            |
| M. Venugopal, C.D. Reddy, C.N. Raju and K.D. Berlin                                | 253 | Synthesis of 2-substituted-2,3-dihydro-3-(2-nitrophenyl)-1 <i>H</i> -naphth[1,2- <i>e</i> ][1,3,2]oxazaphosphorine 2-oxides.                                                                                                                                                |
| E. Barbu and F. Cuiban                                                             | 259 | A method for the synthesis of 3-methyl-2,7-naphthyridine derivatives.                                                                                                                                                                                                       |
| A. Krauze and G. Duburs                                                            | 265 | Synthesis and properties of 4,5-trans-4-aryl-3-cyano-6-hydroxy-6-methyl-5-pyridino-1,4,5,6-tetrahydropyridine-2-thiolates.                                                                                                                                                  |
| D.B. Reddy, A.S. Reddy, R.V. Padmavathi, and N.C. Babu                             | 271 | Synthesis of 5,7-diphenyl(-4-alkyl)trihydro pyrano[3,4- <i>d</i> ][1,2,3]selenadiazoles and 5,7-diphenyl(-4-alkyl)trihydropyrano[3,4- <i>d</i> ][1,2,3]thiadiazoles.                                                                                                        |
| J. Quiroga, S. Cruz, B. Insuasty and R. Abonia                                     | 275 | Synthesis of 5-cyano-4,7-dihydropyrazolo[3,4- <i>b</i> ]pyridin-4-ones and 5-cyano-pyrazolo[3,4- <i>b</i> ]pyridin-4-ones in one-step by the reaction of 5-aminopyrazolone with benzaldehyde and benzoylacetone nitrile in ethanol and by microwave radiation in dry media. |
|                                                                                    | 283 | Notes for the preparation of manuscript.                                                                                                                                                                                                                                    |

# Heterocyclic Communications, Vol. 6, No. 4, 2000

## Vol. 6, Issue 4 (2000)

|                                                                                                  |     |                                                                                                                                                                                                    |
|--------------------------------------------------------------------------------------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                  | 285 | Contributors to this Issue                                                                                                                                                                         |
|                                                                                                  | 286 | Graphical Abstracts                                                                                                                                                                                |
| M. Yamashita, V.K. Reddy, T. Oshikawa and M. Takahashi                                           | 291 | Synthesis of novel nitro-substituted 1-phenoxy-2-phospholenes and phospholanes.                                                                                                                    |
| M.d.M. Blanco, A. Marcos, M. A. Ruiz, C. Avendaño, J.C. Menéndez, C. Pedregal and J.M. Pingarrón | 295 | Electrochemical evidence for oxo-enol tautomerism of azaanthracene-4,9,10-triones.                                                                                                                 |
| R.A. Mekheimer, H.A.A. El-Nabi, A.M. Abd-Elhameed and K.U. Sadek                                 | 301 | A convenient one pot synthesis of pyrido[2,3-d]pyrimidines and pyrimido-[2,3-c]pyrimidines (N <sub>5</sub> -deazapetrins).                                                                         |
| L. Kiss and S. Antus                                                                             | 309 | A convenient synthesis of pterocarpanes.                                                                                                                                                           |
| V. Ravikanth, P. Ramesh, P.V. Diwan and Y. Venkateswarlu                                         | 315 | Microwave irradiation of embelin and evaluation of antibacterial activity.                                                                                                                         |
| S.-M. Lu, W.-J. Xiao and W.-F. Huang                                                             | 319 | A novel and efficient method for the synthesis of $\beta$ -triazolylalkenes via the phase transfer catalyzed Wittig reaction.                                                                      |
| C. Sousa, J.E. Borges, C. Freire, C. Romão and B. Castro                                         | 323 | Synthesis and characterisation of dibenzo-18-crown-6 ethers with appended schiff bases.                                                                                                            |
| S.A. Popov and A.V. Tkachev                                                                      | 327 | Heteroannulations with pinane-derived $\beta$ -enaminaldehyde.                                                                                                                                     |
| H.V.S. Krishnan, S.K. Chowdary, P.K. Dubey, A. Naidu and V. Sunkara                              | 333 | A facile synthesis of mono and dialkoxyquinoxalines.                                                                                                                                               |
| M.A. Barsy, A.K. Khalafallah, M.E. Hassan and A.A. Rezk                                          | 339 | Successful approach for the synthesis of newly fused heterocyclic compounds incorporating phenylperinaphthenone and naphrydine derivatives.                                                        |
| J. Quiroga, B. Insuasty, R. Abonia, P. Hernandez, M. Noguera and A. Sánchez                      | 345 | Synthesis of pyrazolo[3,4-b]pyridines and pyrido[2,3-d]pyrimidinones by hetero-Diels-Alder reaction of pyrazolyl- and pyrimidinilimines under irradiation in dry media.                            |
| Y.T. Reddy, M.K. Rao and B. Rajitha                                                              | 351 | Facile synthesis of 2-cinnamoyl-4-methyl / 4,6-dimethyl-3-phenyl-furo[3,2-C]-quinolines as marked anti microbial agents.                                                                           |
| O.V. Mikhailov, M.A. Kazymova, T.A. Shumilova and L.R. Vafina                                    | 357 | Ni(II) and Cu(II)- containing heterocyclic compounds with 4,4',6-trimethyl-2,8-dithio-3,7-diazanonen-6-dithioamide-1,9 obtained in gelatin-immobilized matrices in the template synthesis process. |
| G.H. Elgemeie, H.Z. Shams, Y.M. Elkholy and N.S. Abbas                                           | 363 | Novel synthesis of N-amino-2-pyridones and cycloalkane ring-fused pyridines containing benzothiazole moiety.                                                                                       |
| N. Gautam, R. Gupta, D.C. Gautam and R.R. Gupta                                                  | 369 | Synthesis of 3-bromo-1-methylphenothiazines by Smiles rearrangement.                                                                                                                               |
|                                                                                                  | 375 | Notes for the preparation of manuscript.                                                                                                                                                           |

# Heterocyclic Communications, Vol. 6, No. 5, 2000

## Vol. 6, Issue 5 (2000)

|                                                                 |     |                                                                                                                                                                  |
|-----------------------------------------------------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                 | 377 | Contributors to this Issue                                                                                                                                       |
|                                                                 | 378 | Graphical Abstracts                                                                                                                                              |
| H. Göker, S. Süzen, C. Kus and D.W. Boykin                      | 385 | Synthesis of new 2-aryl-4H-4-oxo-1-benzopyran oxime ethers.                                                                                                      |
| K. Ito, T. Sato and Y. Ohba                                     | 391 | Synthesis of bowl-shaped chiral homotriazacalixarenes: Chiral transmission from the point chirality of the bridge to the cyclophane moiety.                      |
| F. Lepifre, C. Buon, P. Bouyssou and G. Coudert                 | 397 | Synthesis of 4 <i>H</i> -pyrido[3,2- <i>b</i> ][1,4]oxazine and 3-substituted-4 <i>H</i> -pyrido[3,2- <i>b</i> ][1,4]oxazines via palladium-catalysed reactions. |
| J.S. Yadav, E.J. Reddy, G. Madhavi and T. Ramalingam            | 403 | Facile synthesis of 3-(triphenoxy/phenoxy)-4-phenyl-1,2-dihydro-2-quinolinones.                                                                                  |
| Y.V. Bilokin and S.M. Kovalenko                                 | 409 | Solution and solid-phase approaches to quinconazole and fluquinconazole – inhibitors of fungal ergosterol biosynthesis.                                          |
| J.J.V. Eynde, L. Estievenart and Y.V. Haverbeke                 | 415 | A simple route to novel 3-(5-anuno-1 <i>H</i> -pyrazol-4-yl)-5-methyl-4 <i>H</i> -1,2,4-triazoles.                                                               |
| M. Chen, S. Lu, G. Yuan, S. Yang and X. Du                      | 421 | Synthesis and antibacterial activity of some heterocyclic $\beta$ -enamino esters with 1,2,3-triazole.                                                           |
| Y. Furuya, H. Kiyota and T. Oritani                             | 427 | Synthesis of the southern part of pamamycin-607, an aerial mycelium-inducing substance of streptomyces alboniger.                                                |
| L.N. Rao, C.D. Reddy and C.N. Raju                              | 431 | Synthesis and characterization of 2-alkylcarbamato-1,2,3,4-tetrahydro-1,3,2-benzodiazaphosphorine 2-oxides.                                                      |
| Z. Odabas and N. Talinli                                        | 437 | Synthesis of dinaphthofuranofuran and dinaphthopyranopyran derivatives.                                                                                          |
| S.T. Abu-Orabi, R. Al-Hamdany, S. Shahateet and K. Abu-Shandi   | 443 | Synthesis of substituted benzyl-1 <i>H</i> -1,2,3-triazolo[4,5- <i>d</i> ]pyridazine-4,7-diones.                                                                 |
| S. Malzogu, I. Susnik, J. Vorkapic-Furac, V. Rapić and V. Kovac | 451 | Synthesis of some 2 <i>H</i> -pyrone and ferrocene containing heterocyclic systems.                                                                              |
| R.N. Kumar, S.T. Selvi and P.S. Mohan                           | 457 | Synthesis of dimeric barbiturates.                                                                                                                               |
| A.M.R. Bernardino, M.A. Khan and A.O. Gomes                     | 463 | A new tetracyclic system-3 <i>H</i> ,10 <i>H</i> -pyrazolo[4',3':5,6]pyrido[4,3- <i>b</i> ]indole.                                                               |
| O.S. Moustafa                                                   | 465 | Synthesis and reactions of some new 1,2,4-triazolyl quinoxaline derivatives.                                                                                     |
| J. Lari, D.A. Afshar, H. Sheibani and K. Saidi                  | 473 | Synthesis of a pseudo-cross-conjugated heterocyclic mesomeric betaine (PCCMB).                                                                                   |
| K.S. Chowdary, A. Prashant, S. S. Rao and V.S.H. Krishnan       | 477 | Synthesis of 6-(1,2-dihydro-1-oxo-phthalazin-4-yl)-2 <i>H</i> -1,4-benzoxazin-3(4 <i>H</i> )-ones.                                                               |
|                                                                 | 483 | Notes for the preparation of manuscript.                                                                                                                         |

# Heterocyclic Communications, Vol. 6, No. 6, 2000

## Vol. 6, Issue 6 (2000)

|                                                                                                |     |                                                                                                                                                                                |
|------------------------------------------------------------------------------------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                | 487 | Contributors to this Issue                                                                                                                                                     |
|                                                                                                | 488 | Graphical Abstracts                                                                                                                                                            |
| H. Shao, Q. Ding and J.W. Lown                                                                 | 495 | Studies on the reactions of 2,3-diketopyrido[4,3,2- <i>de</i> ]quinolines with amino acids and amino esters.                                                                   |
| A.L. Baumstark, H.Chen, S.N. Singh and P.C. Vasquez                                            | 501 | Kinetics of epoxidation of $\alpha,\beta$ -unsaturated carbonyl compounds and enol esters by dimethyldioxirane.                                                                |
| M.J. Uddin, A. Shinooka, T. Fujimoto, H. Shirai and I. Yamamoto                                | 505 | Isoxazolidine based new chiral auxiliary for asymmetric synthesis.                                                                                                             |
| A.S.D.d. Matta, C.D. d. Oliveira and G.A. Romeiro                                              | 511 | Synthesis of a new pyranoquinolonic derivative from coumarin.                                                                                                                  |
| N. Nagarapu and N. Ravirala                                                                    | 515 | New 1,4-benzoxazine fused heterocycles II: Synthesis of 5-methyl-2-phenyl-5 <i>H</i> -benzo[b]imidazo[2',1':2,3][1,3]thiazolo[4,5- <i>e</i> ]oxazine.                          |
| V. Kouznetsov, A. Palma, S. Salas, E. Stashenko, G.N. Montenegro and A.G. Fontela              | 519 | Synthesis of new 4-allyl-4-N-benzylaminopiperidines and their spirocyclic products.                                                                                            |
| V.V. Rozhkov, A.M. Kuvshinov and S.A. Shevelev                                                 | 525 | Transformation of 2-aryl-4,6-dinitroindoles.                                                                                                                                   |
| S. Onitsuka, H. Nishino and K. Kurosawa                                                        | 529 | Selective transformation of 3-acetyl-5-aryl-2-methylfurans using photooxygenation.                                                                                             |
| M. Venugopal, C.D. Reddy, M.F.S. Babu and C.S. Reddy                                           | 533 | Novel phosphorus derivatives of sugars.                                                                                                                                        |
| R.A. Tapia, Y. Prieto, G. Zamora, A. Morella and Y. Repetto                                    | 539 | Synthesis and evaluation of the trypanocidal activity of 4-alkylamino-6-nitroquinolines.                                                                                       |
| F.M. Abd El Latif, M.A. Barsy, E.A. Elrady and M.E. Hassan                                     | 545 | A facile synthesis of new pyrazolo- and triazolo[5,1- <i>c</i> ][1,2,4]triazepines derivatives via intermolecular Wittig ring-closure reaction.                                |
| J. Vela-Becerra, P. Sharma, A. Cabrera, C. Alvarez, A. Toscano and G. Penieres                 | 553 | Antimony trichloride mediated trimerization of benzothiazole: Crystal structure of 2,2-bis[benzothiazolyl]benzothiazole.                                                       |
| H. Agirbas and S. Güner                                                                        | 557 | 1,3-Dipolar cycloaddition reactions of N-methyl- <i>c</i> -substituted-phenylnitrones with N-methylmaleimide.                                                                  |
| V. Srivastav, R. Gupta and R.R. Gupta                                                          | 563 | Synthesis of 3-bromo-1-methylphenothiazines by Smiles rearrangement.                                                                                                           |
| G.A. Romeiro, V.F. Ferreira, M.d.S. Costa and M.G.d. Fonseca                                   | 567 | Fischer indole synthesis with alkyl group elimination: A Three-step preparation of a pyrido[2,3- <i>c</i> ]carbazole derivative.                                               |
| J. Charris, J. Dominguez, M. Cordero, L. Orfila, F. Riggione, S. López, D. Enriz and F. Suvire | 571 | Synthesis of 3-amino-2-ethoxycarbonyl pyrido[1,2- <i>a</i> ]thieno-[2,3- <i>d</i> ]-4-pyrimidone derivatives and their antimalarial and cytotoxic activities <i>in vitro</i> . |
| S. Vijayalakshmi and S.P. Rajendran                                                            | 579 | Synthesis of 6-chloro-7-phenyldibenzo[b,h][1,6]naphthyridines.                                                                                                                 |
| R.F. Fandy                                                                                     | 585 | Synthesis of heterocyclic quinones containing bridgehead nitrogen atom from 2-aminonaphtho[2,3- <i>d</i> ]thiazolo-4,9-dione.                                                  |
| R.T. Pardasani, P. Pardasani, M.M. Agrawal, R. Ghosh, G. Mathur, S. Yadav and T. Mukherjee     | 591 | [1,5]-Rearrangement of 4a-heterocycl-4a,5,8,8a-tetrahydro-1,4-naphthoquinones.                                                                                                 |
|                                                                                                | 597 | Notes for the preparation of manuscript.                                                                                                                                       |