

HETEROCYCLIC COMMUNICATIONS

An International Journal in Heterocyclic Chemistry

Published bimonthly by **Freund Publishing House Ltd.**,
Suite 500, Chesham House, 150 Regent Street, London W1R 5FA, England

Notes for the preparation of Camera-Ready Manuscripts

Heterocyclic Communications seeks to publish preliminary communications and full length research papers on all phases of heterocyclic chemistry including inorganic ring systems. All papers to the journal will be reproduced from author's typescripts (camera-ready manuscript) by a reduction of 25%.

Guidelines for Authors

1. Preparation and Mailing of Manuscripts

Author(s) should mail manuscript to respective editor in triplicate. All manuscript should be made on a computer and printed out by a laser printer preferably at 600 dpi or higher resolution. Author(s) should send the editor a floppy disc (31/2" or 51/4") in DOS format (1.4MB for 31/2" and 1.25MB for 51/4") which contains a combined manuscript file including computer drawn structural formulas, figures, tables etc. For the all type setting, Times Roman 10 Pt. font or corresponding font set should be used. Preferably, file format of the manuscript is Adobe Page Maker 6.5/6.0/5.0 or MS Word 97 versions for MS Windows. Structures may be drawn by programs like Chem Draw, ISIS Draw, Chem Win, Chem Window/Coral Draw, Chem Text. Apple Macintosh and other word processor users should convert the file format to those readable by these softwares. Some times RTF (Rich Text Format) and PDF (Portable Document Format) are effective. Details of the program(s) used should be also provided. If structures/diagrams have been prepared using program(s) other than those stated, then structures/diagrams should be converted into windows meta file format. Details of the program(s) used to draw structures/diagrams should be also provided. Manuscript should end with even numbered page (viz 2,4,6....). Normally manuscript not exceeding 6 pages (excluding graphical abstract) will be accepted. Editor can accept even manuscript exceeding 6 pages, if they feel that it is appropriate to do so.

All manuscripts except from North America, European countries and Japan should be mailed to Dr. R.R. Gupta, Editor-in-Chief, Heterocyclic Communications, 10A, Vasundhara Colony, Tonk Road, Jaipur-302018, India. Authors from Europe should mail manuscripts to Prof. Jacques Barbe, Editor for European Countries, Faculte de Pharmacie, 27, Bd. Jean-Moulin, 13385 Marseille Cedex 5, France. Authors from North America should mail manuscripts to Prof. L. Strekowski, Department of Chemistry, Georgia State University, University Plaza, Atlanta, Georgia 30303-3083, USA. Authors from Japan should mail manuscripts to Prof. S. Eguchi, Department of Molecular Design and Engineering, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8603, Japan.

2. Organization of the Manuscript

2.1 Title : It should be clear, concise, and informative. It should be typed with words all in bold face capitals, single spaced, centered on the width of the first page, 2 cm below the top of typing area.

2.2 Author(s) : After 4 single spaces from the title, authors names and addresses should be given on separate lines, typed single spaced, centered on the width of the first page. The name of the author to whom correspondence should be directed should be marked with an asterisk as superscript.

2.3 Abstract : After 4 single spaces from authors address, a short abstract (50-100 words) describing results, methodology employed, conclusions etc., should be placed. It should be labelled as the abstract with the word "Abstract" in bold face and should be typed single spaced. Total typing area on the first page should be 17x22 cm. Typing area for other pages should be 17x24 cm. Each manuscript should be accompanied by a Graphical Abstract, which should contain the title in bold face single spaced, authors names and addresses. It should summarise concisely the contents of the communications in pictorial form designed to capture the attention of a wide readership and to facilitate compilation of database. Chemical Structures designed to illustrate the theme of research work presented should be included. It should be prepared on a separate sheet of paper sized 19x7 cm. An example of graphical presentation is shown as :

Space

Leave blank

1 cm

10 cm

For

Abstract

2.4 Text : Text should be divided into sections like **Introduction, Experimental**, including spectra screening, instruments specifications, **Results, Discussions, Conclusions, Acknowledgements** (if applicable) and **References**. However, the experimental section should be excluded in preliminary Communications as far as possible. Between the end of one section and the beginning of another, there should be a gap of three single spaces. Text should be typed in one and a half spacing throughout.

2.5 References: These should be numbered consecutively throughout the text as they appear and should be superscripted. All compound number in text should be bold e.g., phenothiazines **7** can be prepared... Abbreviations for journals should be those used in Chemical Abstracts. Abbreviations for other terms should also be taken from Chemical Abstracts/ACS journal. Each reference should be typed single spaced. However, there should be one empty line between two consecutive references. Some examples for journal and books references are given:

- 1 R.R. Gupta, V. Saraswat, A. Gupta, M. Jain and V. Gupta, J. Heterocycl. Chem. **29**, 1703 (1992)
- 2 R.R. Gupta, Diamagnetic Susceptibility, Springer Verlag, Berlin, 1986
- 3 R.R. Gupta (Ed.), Physical Methods in Heterocyclic Chemistry, John Wiley, New York, 1984
- 4 R.R. Gupta and M. Kumar, Synthesis, Properties and Reactions of Phenothiazines, in R.R. Gupta (Ed.), Phenothiazines and 1,4-Benzothiazines-Chemical and Biomedical Aspects, Elsevier, Amsterdam, 1988, pp. 1

2.6 Illustrations

Both line drawing and photographs should be used sparingly. All illustrations must be numbered in sequence using arabic numbers, and each illustration must be referred to in the text. Each illustration must be provided with a caption which is complete enough for the figure to be appreciated without referring to the text. Line drawings must be submitted as the perfectly drawn original or as a glossy photograph print. Photocopies, Multiliths, Verifax or Xerox copies are not acceptable substitutes. If you are using desktop publishing software with high quality graphics capability, which is recommended, it is desirable that the graphics be sized and printed along with the text. A laser printer should be used. Photographs must be supplied as glossy prints in which any lettering must be part of the photographs. No lettering should be applied to the surface of the photograph. Micrographs and similar material in which linear dimensions are important should have a scale of length.

2.7 Tables

Tabular presentation of data is an economic way of condensing many items. Refer to tables by using Latin numerals in text and to head the table itself. Provide each table with a caption which will allow it to be read without reference to the text. The table caption should be typed to the width of the table itself. Exceptionally large tables may be placed landscape on the page with the top of table at the left-hand margin.

2.8 Page Numbers

When your typescript is complete and in its final form, number each page sequentially from 1, using a light blue pencil, in the top right-hand corner of each sheet. Final page numbers will be inserted by the editor or the publisher.

2.9 Layout

The typing area should be 17x24 cm for all pages except the first. On the first page typing should start 2 cm below top of the typing area, e.g., its typing area should be 17x22 cm. After abstract text should be typed in one and a half spacing throughout on all pages. Photographic reduction will be by 25%.

2.10 Displayed Equations and Formulas

Leave a blank line above and below all display. Break equations, where possible, only at an equals sign (=) or equivalent (>, <, ≥, ≤). Do not repeat equations sign at the end of one line and the beginning of the next, but carry it over to the beginning of the following line.

REPRINT ORDER FORM FOR HETEROCYCLIC COMMUNICATIONS

FREUND PUBLISHING HOUSE LTD.
P.O.Box 35010, Tel Aviv, Isreal, 61350

Telephone: 972-3-562-8540
Fax: 972-3-562-8538
E-mail: h_freund@netvision.net.il

Author of the manuscript:

Title of the manuscript:

TWENTY-FIVE reprints will be supplied free of charge for each **ARTICLE**.
(PLEASE NOTE: **NO FREE REPRINTS ARE SUPPLIED FOR BOOKS.**)
Reprints may be purchased in units of 100 only.

COST OF REPRINTS

No. of copies	Number of Pages					Each Add. Unit (1-4 pp)
	1-2 pp	3-4 pp	5-8 pp	9-12 pp	13-16 pp	
100	\$ 83	122	159	253	341	111
200	\$ 109	150	259	380	516	136
300	\$ 128	245	380	569	759	172

Orders for quantities of over 300 copies are subject to special quotation.

- * **IMPORTANT: No reprints will be sent unless this form is filled in and signed.**
- * Type may be destroyed after the printing of the issue.
- * Please note that payment is to be received in U.S. Dollars.
- * Co-author do not receive copies of this form.
- * Author copies of the journal and reprints will be sent by airmail.

****PLEASE TYPE****

Please supply me with reprints, in the addition to the 25 free copies to which I am entitled.

NAME.....

ADDRESS.....

TELEPHONE..... FAX.....

E-MAIL.....

INVOICE TO:.....

PRICE OF REPRINTS:.....

☐ I enclosed a check (made payable to FREUND PUBLISHING HOUSE LTD.) in the amount of \$.....

☐ Please debit my: **VISA/AMERICAN EXPRESS/MASTER CARD** for the amount of \$.....

Number..... Expiry Date.....

Index of Vol. 7 (Issues 1-6), 2001

Heterocyclic Communications, Vol. 7, No. 1, 2001

Vol. 7, Issue 1 (2001)

	1	Contributors to this Issue
	2	Graphical Abstracts
S.A. Jennings, J.L. Toth, S.G. Roller, N. Brooks, C. O'Hara, K. Kiakos, J.A. Hartley, P.J. Burke and M. Lee	7	Efficient synthesis of (+)-seco-cyclopropaneindoline analogs of CC-1065.
F.A. Chowdhury, H. Nishino and K. Kurosawa	17	Synthesis of azabicyclic peroxides connected by methylene chain using manganese(III)-catalyzed simultaneous cyclization on both ends.
M. Dupuy, F. Pinguet, O. Chavignon, J.-C. Teulade, J.-P. Chapat and Y. Blache	23	Reactivity of 6,7,8,9-tetrahydropyrido[1,2-a]benzimidazol-9-one : Synthesis of pyrrolo pyridobenzimidazoles.
R. Augusti	29	A mechanistic proposal for the formation of unexpected pyrroles in reactions of of carboethoxycarbene with enaminones.
L. Nagarapu and K. Sambaru	33	Synthesis of 5-(4-methyl phenyl sulfonyl)-10-phenyl-6,7-dihydro-5H-benzo[b]imidazo[2',1':2,3][1,3]thiazolo[4,5-d]azepin-5-yl.
Y. Luo, L. He, M. Ding, G. Yang, A. Luo, X. Liu and T. Wu	37	A convenient synthesis of derivatives of 1,3,2-dioxaphosphocane-2-sulfide with bioactivity via Lawesson's reagent.
Z. Tokic-Vujosevic and Z. Cekovic	43	Synthesis of new stereoisomeric nitrate esters derived from isosorbide-mononitrates.
S.P. Singh, J.K. Kapoor and A.K. Sadana	49	Synthesis of 4-acetyl or (carbethoxy)-5-methyl-1-heterocyclylpyrazoles.
I. Wiedermannová and J. Slouka	55	Cyclization reactions of hydrazones XXVII : Synthesis and oxidative cyclization of some formazans of quinoxaline group.
A. Prashant, S.S. Rao, K.S. Chowdary and V.S.H. Krishnan	61	Synthesis of some new 1,5-benzothiazepins containing 2H-1-benzopyran-2-one heterocycle.
S. Balalaie and N. Nemati	67	One pot preparation of coumarins by Knoevenagel condensation in solvent free condition under microwave irradiation.
S.H. Mashraqui, S. Kumar and K.R. Nivalkar	73	Attempt towards syn-benzothiazolophane: Synthesis and conformational study of syn-dithia-[3.3](2,6)-benzothiazolophane.
O.V. Mikhailov	79	Low-temperature template synthesis of nickel-containing heterocyclic Compound with 2,8-dithio-3,5,7-triazanone and dithioamide-1,9 in gelatin-immobilized matrix.
S.B. Desai, P.B. Desai and K.R. Desai	83	Synthesis of some schiff bases, thiazolidinones and azetidinones derived from 2,6-diamino benzo[1,2-d:4,5-d']bisthiazole and their anticancer activities.
	91	Notes for the preparation of manuscript.
	94	Index of Vol. 6 (Issue 1-6), 2000.

Heterocyclic Communications, Vol. 7, No. 2, 2001

Vol. 7, Issue 2 (2001)

	101	Contributors to this Issue
	102	Graphical Abstracts
H. Shao, Q. Ding and J.W. Lown	107	Studies on the reactions of 2,3-diketopyrido[4,3,2- <i>de</i>]quinolines with aliphatic amines.
T. Oshikawa, K. Seo, M. Yamashita, M. Takahashi and Y. Hamazu	113	Crystal structure analysis of a 1,2,3,4-tetra- <i>o</i> -acetyl-5-deoxy-5-phenylphosphinyl-d- <i>ribo</i> -pyranose derivative.
L. Strekowski, T. Gorecki, J.C. Mason, H. Lee and G. Patonay	117	New heptamethine cyanine reagents for labeling of biomolecules with a near-infrared chromophore.
P.S. Goud, P.M. Goud and D. Ashok	123	A facile syntheses of ethyl 5-substituted allyloxy 11 <i>H</i> -benzo[<i>e</i>]thieno-[2',3':4,5]pyrrolo[1,2- <i>a</i>][1,4]diazepine-2-carboxylates.
L.Y.V. Méndez, V. Kouznetsov, J.C. Poveda, C. Yolaçan, N. Öcal and F. Aydoğan	129	Transformation of 4- <i>N</i> -arylamino-4-(8-quinolinyl)-1-butenes and 3-aryl-2-(8-quinolinyl)-4-thiazolidinones.
A.S. Capilla, N. Hernández and M.D. Pujol	135	NIS or NBS mediated formation of halo acetals. A convenient method for the one pot disubstitution of 1,4-benzodioxin.
N.A. Ganoub	143	A facile approach to <i>N</i> -heterocycles. The reactions of ylide phosphoranes with hydrazones.
R. Badamurali and K.J.R. Prasad	149	Synthesis of tetracyclic carbazole derivatives.
A.N. Andin, V.A. Kaminskii and S.V. Dubovitskii	155	The synthesis of 3-acetyl-2-(4,4-dimethyl-2,6-dioxocyclohexyl)-1-phenylpentanedione-1,4 and its reactions with <i>N</i> -nucleophiles.
A.M.E. Massry, M.M.A. Rahman, S.E. Sayed and E.S.H.E. Ashry	159	Construction of some pyridazine derivatives and their annelated ring systems from keto acids.
H. Gaveriya, B. Desai, V. Vora and A. Shah	167	Synthesis of some new unsymmetrical 1,4-dihydropyridine derivatives as potent antitubercular agents.
Y.Y. Morzherin, T.V. Glukhareva, V.S. Mokrushin, A.V. Tkachev and V.A. Bakulev	173	Application of the Hurd-Mori reaction for the synthesis of chiral 1,2,3-thia-(seleno)diazole derivatives from (+)-3-carene and α -pinene.
S. Vijayalakshmi and S.P. Rajendran	177	A new approach to the synthesis of benzo[<i>b</i>][1,8]naphthyridin-4(1 <i>H</i>)ones.
F.A. Abu-Shanab, Y.M. Elkholy and M.H. Elnagdi	183	Enaminones as building blocks in organic synthesis: Synthesis of new poly-functional pyridines, condensed pyridines and pentasubstituted benzene.
R.N. Kumar, T. Suresh, A. Mythili and P.S. Mohan	193	Facile entry to pyrimido[4,5]quinolines and its thio analogues.
	199	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 7, No. 3, 2001

Vol. 7, Issue 3 (2001)

	203	Contributors to this Issue
	204	Graphical Abstracts
H. Salgado-Zamora, E. Campos, R. Jiménez, R. Ruiz, T. Castaneda and S. Turiján	209	Oxidative degradation of arylfuro-1,2-oxazoles to arynitriles by potassium permanganate.
M. Trávníček, M. Potáček and A. Mannschreck	213	Stereochemistry of 1,3-dipolar cycloaddition reaction of azomethine ylides derived from <i>N</i> -alkyl- <i>N</i> -(4-toluenesulphonyl) carbamoylmethyl phenanthridinium with olefinic dipolarophiles.
H. Ishida, H. Asaji, K. Itoh and M. Ohno	223	Cycloaddition reaction of C ₆₀ with 2-diazo-1,3-dithiane and oxidation of the cycloadduct to sulfoxide derivatives.
G.A. Romeiro and P.R.C. Martins	227	Analogues of antidepressants-Synthesis of new 4-amine derivatives of 10,11-dihydro-5H-dibenzo[a,d]cycloheptane.
J.E. Charris, J.N. Domínguez and S.E. López	233	A practical synthesis of ureas from 2-aminopicolines using microwave irradiation.
L. Nagarapu, N. Ravirala and D. Akkewar	237	Benzothiazepine fused heterocycles IV : A convenient synthesis of benzo[b][1,5]thiazepines using MCH-41(H) zeolite.
S. Man, J. Dostal, M. Necas, Z. Zák and M. Potáček	243	Berberine and coptisine in liquid ammonia.
M.C.S. de Mattos, S.P.L. de Souza and S.M. Elias	249	Transition metal salts mediated oxidative radical addition of ethyl acetoacetate to limonene.
K.A. Kumar, C.S. Reddy, L.N. Rao and C.N. Raju	253	Synthesis and characterization of 6-substituted dibenzo[d,f][1,3,2]dioxaphosphin 6-oxides.
E. Tóth, Z. Dinya, L. Szilágyi and S. Antus	257	Synthesis of deuterium labeled pterocarpanes.
H.H. Abdel-Razik, H.M. Refat and M.E.A. Zaki	263	Synthesis of some new 2,6-diamino-4-(<i>p</i> -aryloxy)anilinopyrimidine and some related 5-aryloxy pyrimidine derivatives for dyeing synthetic fibres.
P.N. Kumar, B. Prasanna and G.V.P. Chandramouli	271	Facile synthesis of substituted 4-hydroxy coumarinbenzothiophenes.
M. Tosa, C. Paizs, C. Majdik, L. Poppe, P. Kolonits, I. Silberg, L. Novák and F-D. Irimie	277	Selective oxidation methods for preparation of <i>N</i> -alkylphenothiazine sulfoxides and sulfones.
V.I. Gulevskaya, A.M. Kuvshinov and S.A. Shevelev	283	Synthesis of 3-arylamino-4,6-dinitrobenzo[b]thiophene-2-methylcarboxylates. Smooth dehydrogenation of 2,3-dihydrobenzo[b]thiophene derivatives.
N. Dodia and A. Shah	289	Synthesis of some tricyclic and tetracyclic ring systems built of 4-hydroxy-2-quinolones.
	295	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 7, No. 4, 2001

Vol. 7, Issue 4 (2001)

	299	Contributors to this Issue
	300	Graphical Abstracts
S.E. McKay, J.A. Sooter, S.G. Bodige and S.C. Blackstoch	307	Oxidation methods for aromatic diazines: Substituted pyrazine- <i>N</i> -oxides, pyrazine- <i>N,N'</i> -dioxides, and 2,2':6',2''-terpyridine-1,1''-dioxide.
T. Watanabe, H. Kiyota, R. Takeuchi, K. Enari and T. Oritani	313	Stereochemistry of enacyloxins 2. Structure elucidation of decarbamoyl enacyloxin IIa and IVa, new members of enacyloxin antibiotics from <i>Frateruria</i> sp. W-315.
G. Pastuch, I. Wandzik, W. Szeja, G. Gryniewicz, J. Ramza, P. Priebe and W. Pucko	317	New synthesis of O- and S-glycosyl derivatives of 2-chloro-3-cyano-5-nitropyridine.
L.Y.V. Méndez, V. Kouznetsov, E. Stashenko, A. Bahsas and J. Amaro-Luis	323	4- <i>N</i> -Aryl(benzyl)amino-4-hetaryl-1-butenes as building blocks in heterocyclic synthesis. 1. New route to 4,6-dimethyl-2-pyridylquinolines from 4- <i>N</i> -p-methylphenylamino-4-pyridyl-1-butenes.
T. Fujimori, O. Nakayama, H. Kiyota, Y.-i. Kamijima, T. Watanabe and T. Oritani	327	Stereochemistry of enacyloxins 1. Absolute configuration of the cyclohexane ring part of enacyloxins, a series of antibiotics from <i>Frateruria</i> sp. W-315.
R. Buffa, P. Zahradnik and P. Foltinova	331	Computer projected benzothiazole derivatives. Synthesis structure and biological study of new push-pull conjugated benzothiazolium salts.
S. Balalaie, A. Sharifi, B. Ahangarian and E. Kowsari	337	Microwave enhanced synthesis of quinoxolines in solvent-free condition.
M.H.N. Arsanious	341	The behaviour of diphenylmethylenetriphenylphosphorane and phosphites toward 5-substituted-1,3,4-thiadiazol derivatives.
G. Zedníková, K. Nálepá and T. Gucky	349	Synthesis of some new 3,5,7-trisubstituted pyrazolo[4,3- <i>e</i>]1,2,4-triazines.
G.A. Prakash and S.P. Rajendran	353	A convenient synthesis of 8,9,10,11-tetrahydridibenzo[<i>b,h</i>][1,6]naphthyridin-6(5H)ones.
O.V. Mikhailov, A.I. Khamitova and L.S. Mingalieva	359	Co(II,III), Ni(II) and Cu(II)- containing macrocyclic compounds with 2,7-dithio-3,6-diazoctadien-3,5-dithioamide-1,8 obtained in gelatin-immobilized matrix by template synthesis process.
G. Keglevich, H. Szelke and L. Töke	365	Novel bridged P-heterocycles: The first 2,3,5-diazaphosphabicyclo[2.2.2]oct-7-ene 5-oxides.
I. Saramet, C. Draghici, C. Barcutean, V. Radulescu, T. Loloiu and M.D. Banciu	369	Synthesis of new 4-substituted-1-aroylethiosemicarbazides and their cyclization to mercaptotriazoles and aminothiadiazoles.
A. Krauze, L. Sile and G. Duburs	375	Convenient one-pot synthesis of 2-carbamoylmethylthio-3-cyano-4,6-diaryl-5-ethoxycarbonyl-1,4-dihydropyridines.
R.M. Dinica and C. Pettinari	381	Synthesis of substituted 7,7'-bis-indolizines via 1,3-dipolar cycloaddition under microwave irradiation.
H.A. Radhy, G.F. Fadhil, A. Perjéssy, E. Kolehmaincn, W.M.F. Fabian, M. Samaliková, K. Laihai and Z. Susteková	387	Substituent effects and vibrational coupling in indole-2,3-diones: An ir, nmr and theoretical study.
K. Sharma, R. Swaroop and R.V. Singh	393	Heterocyclic complexes of palladium (II) : Template synthesis, spectroscopic studies and biochemical aspects.
	399	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 7, No. 5, 2001

Vol. 7, Issue 5 (2001)

	403	Contributors to this Issue
	404	Graphical Abstracts
L. Janda	411	Synthesis of 5-aryl-1,3,4-oxadiazolyl-2-acetic acids.
L. Kiss, G. Papp, F. Joó and S. Antus	417	Efficient synthesis of pterocarpanes by heck-oxyarylation in ionic liquids.
S. Deprets and G. Kirsch	421	A first synthesis of isothiocomestanes and heterocyclic analogues.
A. Krauze, V. Garalene, R. Vitolina and G. Duburs	427	Synthesis of 4,5-dipyridylpyridin-2(1H)-ones, pyridine-2(1H)-thiones and related derivatives as analogues of cardiotonic drug milrinone.
L. Nagarapu and N. Ravirala	433	New 1,4-benzothiazine fused heterocycles-V : Synthesis of 9H-thieno[3,2- <i>b</i>]-benzothiazine and 4H-thiazolo[2,3- <i>b</i>][1,4]benzothiazine derivatives.
M. Saoul, F.B. Benabdelouahad, F.E. Guemmout and A.M. Romerosa	439	Convenient synthesis of 3-(1-carboxyalkyl)pyrido[2,3- <i>d</i>]pyrimidine-2,4-diones.
K.P. da Silva, I.A. de Souza, A.R. de Faria, D.J. Brondani and A.C.L. Leite	445	Synthesis of peptidyl benzodioxole derivatives from safrole, a potential class of antitumor drugs.
M.S.R. Murty, T. Ramalingam, G. Sabitha and J.S. Yadav	449	Facile synthesis of 9,10-dihydro-4H-benzo [4,5] cyclohepta [1,2- <i>b</i>] thiophene-4-one: A crucial drug intermediate-application of Wittig-Horner reaction.
Z. Ates-Alagoz and E. Buyukbingol	455	Synthesis of some novel tetrahydronaphthalene benzimidazole derivatives.
A.V. Varlamov, T.N. Borisova, L.G. Voskressensky, B. Nsabimana and A.I. Chernyshev	461	Tandem Michael addition - Hoffman elimination sequence of DMAD on tetrahydropyrrolo[3,2- <i>c</i>]pyridines. New route to vinylpyrroles.
V. Padmvathi, B.J.M. Reddy, M.R. Sarma, A. Padmaja and D.B. Reddy	467	Cyclohexanone derivatives : Synthons for substituted quinolines.
S.E. López, M.E. Rosales, C.E. Canelón, E.A. Valverde, R.C. Narváez, J.E. Charris, F.A. Giannini, R.D. Enriz, M. Carrasco and S. Zacchino	473	Synthesis and preliminary cytotoxic and antifungal evaluation of some 6- <i>N,N</i> -dialkyl 2-aryl-4(3 <i>H</i>)-quinazolinone derivatives.
H. Gevariya, B. Desai, V. Vora and A. Shah	481	Synthesis of some new unsymmetrical 1,4-dihydropyridine derivatives as potent antitubercular agents.
F.M.A.E. Latif, M.A. Kahlil, I. Helmy and H.A. Solieman	485	A mild and efficient synthesis of new benzimidazole derivatives via a one-pot reaction. An addition versus condensation reaction.
A.A.F. Wasfy, A.L. El-Shenawy and S.A. Nassar	493	Synthesis and biological activity of some 4-(5-alkyl or 5-aryl-2-furfurylidene-1,3-disubstituted-2-pyrazolin-5-ones.
A. Nasreen and S.R. Adapa	501	An efficient facile and selective hydroxylation of nitrogen heterocycles.
	507	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 7, No. 6, 2001

Vol. 7, Issue 6 (2001)

	511	Contributors to this Issue.
	512	Graphical Abstracts.
M.D. Rosa, M. LaRue, L Sellitto and M.D. Timken	519	Reaction of 2-aminopyrrole and 1-alkyl-2-aminopyrroles with DMAD: Michael addition vs Diels-Alder reaction.
L. Streckowski and H. Lee	521	Acridine formation by the reaction of 2-nonafluorobutylaniline with 2-mesitylmagnesium bromide.
T.G. Silva, F.S.V. Barbosa, S.S.F. Brandao, M.C.A. Lima, S.L. Galdino, I.R. Pitta and J. Barbe	523	Synthesis and structural elucidation of new benzylidene imidazolidines and acridinylidene thiazolidines.
D.W. Boykin and H. Goker	529	Ring cleavage of dicyano-substituted benzimidazoles by DIBAL.
L. Nagarapu and N.V. Rao	535	Synthesis of phenylimidazo thiazolo benzocycloheptene derivatives as potential antiinflammatory agent-IV.
A.-F.E. Mourad, A.A. Hassan, N.K. Mohamed, A.A. Aly and B.A. Ali	541	Novel heterocycles from creatinine.
L. Gaina, T. Lovasz, L.A. Silberg, C. Cristea and S. Udrea	549	New Schiff bases derived from 3-formyl-10-alkyl-phenothiazine I. NMR and uv-vis structural assignments.
A.C.L. Leite, K.P. da Silva and D.J. Brondani	555	Synthesis of 1,3-benzodioxole derivatives containing a amino acid moiety in side chain.
M.A.M. Taha	559	Synthesis of 7H-1,2,4-triazolo[1,5-d] tetrazolo acyclo C-nucleosides.
I. Panea, A. Ghirisan, I. Cristea, R. Gropeanu and I.A. Silberg	563	Azocoupling products. II. Synthesis and structural study of azocoupling products of 1-(5,6-dimethyl-4-X-pyrimidin-2-yl)-3-methyl-pyrazolin-5-ones with aromatic diazonium salts.
A. Dandia, R. Singh, G. Kumar, K. Arya and H. Sachdeva	571	Facile one pot microwave induced synthesis of spiro[indole-pyrazoles] and spiro [indol-isoxazoles].
M. Lazarevic, V. Dimova, D.M. Gabor, V. Kakurinov and K.C. Ragenovic	577	Synthesis of some N ¹ -aryl/heteroarylaminomethyl/ethyl-1,2,4-triazoles and their antibacterial and antifungal activities.
A.A. Abdel-Hafez and M.F. El-Zohry	583	Synthesis of some new spiropyrazolo[4,5-c]-benzodiazepines and spiropyrazolo-[4,5-c]-benzoxazepines.
A. Nagpal, R. Unny, P. Joshi and Y.C. Joshi	589	Synthesis of 1,3-diketone and its reaction with different N-nucleophiles (part I).
M. Darabantu, C. Maieranu, G. Plé, C. Berghian, E. Condamine and Y. Ramondenc	593	Convenient synthesis of chiral 5-amino-1,3-dioxanes built on some <i>l-p</i> -nitro-phenylserinols skeletons.
V.H. Patel, M.P. Patel and R.G. Patel	599	Synthesis and application of novel heterocyclic dyes based on 11-amino-3-bromo-13H-acenaphtho[1,2-e]-pyridazino[3,2-b]quinazoline-13-one.
	607	Notes for the preparation of manuscript.