

Subject index Volume 254

Alkaline-earth metal cations

Exploration of the catalytic oxidation of ethylene with N_2O mediated by atomic alkaline-earth metal cations, 155

Alkanium ions

C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189

Atmospheric chemistry

A gas-phase solvent effect: The role of water molecules in the conversion of the $HOMg^+/CO_2$ adduct ion to the magnesium bicarbonate ion, Mg^+O_2COH , 136

Benzene

Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152

Bicarbonate formation

A gas-phase solvent effect: The role of water molecules in the conversion of the $HOMg^+/CO_2$ adduct ion to the magnesium bicarbonate ion, Mg^+O_2COH , 136

Binomial and multinomial distributions

Random protein association analyses of MALDI-TOF mass spectra of two- and three-component protein systems using binomial and multinomial distribution, 101

Bond dissociation energies

Why is hafnium so unreactive? Experimental and theoretical studies of the reaction of Hf^+ with methane, 168

Carbocations

C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189

Carbon

TOF-SIMS characterisation of spark-generated nanoparticles made from pairs of Ir–Ir and Ir–C electrodes, 70

Catalysis

Activation of methane and methane- d_4 by ionic platinum clusters, 183
Formation, deposition and examination of size selected metal clusters on semiconductor surfaces: An experimental setup, 202

Catalysis by metal hydrides

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, $HMgL_2^-$ ($L = Cl$ and HCO_2), 145

CBS-QB3 model chemistry

The loss of NH_2O^\bullet from the *N*-hydroxyacetamide radical cation $CH_3C(=O)NHOH^{+\bullet}$: An ion-catalysed rearrangement, 127

CF-IRMS

Memory effect in deuterium analysis by continuous flow isotope ratio measurement, 44

C–H activation

Activation of methane and methane- d_4 by ionic platinum clusters, 183

C–H bond activation

Why is hafnium so unreactive? Experimental and theoretical studies of the reaction of Hf^+ with methane, 168

Chemical ionization

Chemical ionization reaction time-of-flight mass spectrometry: Multi-reagent analysis for determination of trace gas composition, 85

Chemical reaction

Reactions in nanofilms of trifluoroacetic acid (CF_3COOH) driven by low energy electrons, 63

CID

Ion chemistry of a series of cluster compounds with Mo_3Q_4 and $Mo_3M'Q_4$ ($Q = S, Se$; $M' = Cu, Co, Ni$) cores containing 1,2 diphosphanes as ancillary ligands: New insights on the gas-phase stability from electrospray tandem mass spectrometry, 28

CIRMS

Chemical ionization reaction time-of-flight mass spectrometry: Multi-reagent analysis for determination of trace gas composition, 85

Cluster reaction

Interactions of CO with $Au_nO_m^-$ ($n \geq 4$)

CO oxidation

Interactions of CO with $Au_nO_m^-$ ($n \geq 4$)

CO poisoning

Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152

Collision induced dissociation

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, $HMgL_2^-$ ($L = Cl$ and HCO_2), 145

Correction

Memory effect in deuterium analysis by continuous flow isotope ratio measurement, 44

Cuboidal complexes

Ion chemistry of a series of cluster compounds with Mo_3Q_4 and $Mo_3M'Q_4$ ($Q = S, Se$; $M' = Cu, Co, Ni$) cores containing 1,2 diphosphanes as ancillary ligands: New insights on the gas-phase stability from electrospray tandem mass spectrometry, 28

δD deuterium measurements

Memory effect in deuterium analysis by continuous flow isotope ratio measurement, 44

DEA

Reactions in nanofilms of trifluoroacetic acid (CF_3COOH) driven by low energy electrons, 63

Decarboxylation

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, $HMgL_2^-$ ($L = Cl$ and HCO_2), 145

Dehydrogenation kinetics

Activation of methane and methane- d_4 by ionic platinum clusters, 183

Density-functional calculations

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, $HMgL_2^-$ ($L = Cl$ and HCO_2), 145

Diphosphane

Ion chemistry of a series of cluster compounds with Mo_3Q_4 and $Mo_3M'Q_4$ ($Q = S, Se$; $M' = Cu, Co, Ni$) cores containing 1,2 diphosphanes as ancillary ligands: New insights on the gas-phase stability from electrospray tandem mass spectrometry, 28

Dipolar ac

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Electron stimulated desorption

Reactions in nanofilms of trifluoroacetic acid (CF_3COOH) driven by low energy electrons, 63

Electrospray ionisation

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, HMgL_2^- ($\text{L} = \text{Cl}$ and HCO_2), 145

Electrospray ionization as a convenient new method for the generation of catalytically active iron-oxide ions in the gas phase, 197

ESI-MS

Ion chemistry of a series of cluster compounds with Mo_3Q_4 and $\text{Mo}_3\text{M}'\text{Q}_4$ ($\text{Q} = \text{S}, \text{Se}$; $\text{M}' = \text{Cu}, \text{Co}, \text{Ni}$) cores containing 1,2 diphosphanes as ancillary ligands: New insights on the gas-phase stability from electrospray tandem mass spectrometry, 28

Ethylene

Exploration of the catalytic oxidation of ethylene with N_2O mediated by atomic alkaline-earth metal cations, 155

Excitation

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Field solver

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Fourier transform

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Free fatty acid

Identification of molecular species of simple lipids by normal phase liquid chromatography–positive electrospray tandem mass spectrometry, and application of developed methods in comprehensive analysis of low erucic acid rapeseed oil lipids, 106

Gas phase

Interactions of CO with Au_nO_m^- ($n \geq 4$)

Gas-phase ions

Gas-phase infrared multiple photon dissociation spectroscopy of mass-selected molecular ions, 1

Gold cluster

Interactions of CO with Au_nO_m^- ($n \geq 4$)

Guided ion beam mass spectrometry

Why is hafnium so unreactive? Experimental and theoretical studies of the reaction of Hf^+ with methane, 168

Hafnium

Why is hafnium so unreactive? Experimental and theoretical studies of the reaction of Hf^+ with methane, 168

Harmonic motion

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Hydride abstraction

C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189

Hyperpolarizability

Effects of polarizability and hyperpolarizability on molecular spatial alignment in intense light fields, 37

Infrared

Gas-phase infrared multiple photon dissociation spectroscopy of mass-selected molecular ions, 1

Intense light fields

Effects of polarizability and hyperpolarizability on molecular spatial alignment in intense light fields, 37

Ion source

Formation, deposition and examination of size selected metal clusters on semiconductor surfaces: An experimental setup, 202

Ion trajectory simulation

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

Ion-catalysis

The loss of NH_2O^* from the *N*-hydroxyacetamide radical cation $\text{CH}_3\text{C}(=\text{O})\text{NHOH}^{+\bullet}$: An ion-catalysed rearrangement, 127

A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136

Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152

Exploration of the catalytic oxidation of ethylene with N_2O mediated by atomic alkaline-earth metal cations, 155

Ion–ligand binding

A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136

Ion–molecule reactions

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, HMgL_2^- ($\text{L} = \text{Cl}$ and HCO_2), 145

Iridium

TOF-SIMS characterisation of spark-generated nanoparticles made from pairs of Ir–Ir and Ir–C electrodes, 70

Iron cations

Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152

Iron hydroxide

Electrospray ionization as a convenient new method for the generation of catalytically active iron-oxide ions in the gas phase, 197

Iron oxide

Electrospray ionization as a convenient new method for the generation of catalytically active iron-oxide ions in the gas phase, 197

Isotope analysis

Application of non-resonant laser ionization mass spectrometry for a fast isotope analysis of metal microparticles, 122

Isotope ratio

Optimization of the conditions for simultaneous non-selective excitation of plutonium isotopes for isotope ratio measurements in resonance ionization mass spectrometry, 94

Laser ionization mass spectrometry

Application of non-resonant laser ionization mass spectrometry for a fast isotope analysis of metal microparticles, 122

Laser spectroscopy

Optimization of the conditions for simultaneous non-selective excitation of plutonium isotopes for isotope ratio measurements in resonance ionization mass spectrometry, 94

Low erucic acid rapeseed oil

Identification of molecular species of simple lipids by normal phase liquid chromatography–positive electrospray tandem mass spectrometry, and application of developed methods in comprehensive analysis of low erucic acid rapeseed oil lipids, 106

Magnesium formate anions

Dehydrogenation of formic acid catalyzed by magnesium hydride anions, HMgL_2^- ($\text{L} = \text{Cl}$ and HCO_2), 145

Main-group metal ion

A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136

MALDI-TOF MS

Random protein association analyses of MALDI-TOF mass spectra of two- and three-component protein systems using binomial and multinomial distribution, 101

Mass spectrometry

Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53

TOF-SIMS characterisation of spark-generated nanoparticles made from pairs of Ir–Ir and Ir–C electrodes, 70

Optimization of the conditions for simultaneous non-selective excitation of plutonium isotopes for isotope ratio measurements in resonance ionization mass spectrometry, 94

- C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189
Electrospray ionization as a convenient new method for the generation of catalytically active iron-oxide ions in the gas phase, 197
- Memory effect
Memory effect in deuterium analysis by continuous flow isotope ratio measurement, 44
- Metal clusters
Formation, deposition and examination of size selected metal clusters on semiconductor surfaces: An experimental setup, 202
- Metal nitrate
Electrospray ionization as a convenient new method for the generation of catalytically active iron-oxide ions in the gas phase, 197
- Meteoric ablation
A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136
- Methane
Why is hafnium so unreactive? Experimental and theoretical studies of the reaction of Hf^+ with methane, 168
Activation of methane and methane- d_4 by ionic platinum clusters, 183
- Microparticle
Application of non-resonant laser ionization mass spectrometry for a fast isotope analysis of metal microparticles, 122
- Molybdenum sulfide clusters
Ion chemistry of a series of cluster compounds with Mo_3Q_4 and $\text{Mo}_3\text{M}'\text{Q}_4$ ($\text{Q} = \text{S, Se}$; $\text{M}' = \text{Cu, Co, Ni}$) cores containing 1,2 diphosphanes as ancillary ligands: New insights on the gas-phase stability from electrospray tandem mass spectrometry, 28
- Mono-, Di- and Triacylglycerol
Identification of molecular species of simple lipids by normal phase liquid chromatography–positive electrospray tandem mass spectrometry, and application of developed methods in comprehensive analysis of low erucic acid rapeseed oil lipids, 106
- Multiple photon dissociation
Gas-phase infrared multiple photon dissociation spectroscopy of mass-selected molecular ions, 1
- Multistage mass spectrometry
Dehydrogenation of formic acid catalyzed by magnesium hydride anions, HMgL_2^- ($\text{L} = \text{Cl and HCO}_2$), 145
- N_2 and Cl_2 molecules
Effects of polarizability and hyperpolarizability on molecular spatial alignment in intense light fields, 37
- N_2O reduction
Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152
Exploration of the catalytic oxidation of ethylene with N_2O mediated by atomic alkaline-earth metal cations, 155
- Nanoparticle
TOF-SIMS characterisation of spark-generated nanoparticles made from pairs of Ir–Ir and Ir–C electrodes, 70
- Normal phase liquid chromatography – Positive electrospray ionization tandem mass spectrometry
Identification of molecular species of simple lipids by normal phase liquid chromatography–positive electrospray tandem mass spectrometry, and application of developed methods in comprehensive analysis of low erucic acid rapeseed oil lipids, 106
- O-atom transport
Catalytic reduction of N_2O by CO on benzene clusters of Fe^+ : Catalytic poisoning by CO, 152
Exploration of the catalytic oxidation of ethylene with N_2O mediated by atomic alkaline-earth metal cations, 155
- Orbitrap
Ion trajectory simulations of axial ac dipolar excitation in the Orbitrap, 53
- Orthogonal time-of-flight mass spectrometry
Correlated phase space distributions of ions in an orthogonal time-of-flight mass spectrometer, 20
- Platinum
Activation of methane and methane- d_4 by ionic platinum clusters, 183
- Polarizability
Effects of polarizability and hyperpolarizability on molecular spatial alignment in intense light fields, 37
- Proton-transfer
Chemical ionization reaction time-of-flight mass spectrometry: Multi-reagent analysis for determination of trace gas composition, 85
C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189
- PTR-MS
Chemical ionization reaction time-of-flight mass spectrometry: Multi-reagent analysis for determination of trace gas composition, 85
- Pu
Optimization of the conditions for simultaneous non-selective excitation of plutonium isotopes for isotope ratio measurements in resonance ionization mass spectrometry, 94
- Quantum chemical calculation
A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136
- Quantum chemistry
C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189
- Quid-pro-quo mechanism
The loss of $\text{NH}_2\text{O}^\bullet$ from the *N*-hydroxyacetamide radical cation $\text{CH}_3\text{C(=O)NHOH}^{+\bullet}$: An ion-catalysed rearrangement, 127
- Random protein association
Random protein association analyses of MALDI-TOF mass spectra of two- and three-component protein systems using binomial and multinomial distribution, 101
- Reaction mechanism
C–H activation in reactions of protonated hydrogen peroxide with propane. Remarkable regioselectivity for hydride abstraction and mechanism for acid catalyzed hydrogen exchange, 189
- Resonance ionization mass spectrometry
Optimization of the conditions for simultaneous non-selective excitation of plutonium isotopes for isotope ratio measurements in resonance ionization mass spectrometry, 94
- Size selected
Formation, deposition and examination of size selected metal clusters on semiconductor surfaces: An experimental setup, 202
- Solvent-assisted reaction
A gas-phase solvent effect: The role of water molecules in the conversion of the $\text{HOMg}^+/\text{CO}_2$ adduct ion to the magnesium bicarbonate ion, $\text{Mg}^+\text{O}_2\text{COH}$, 136
- Space-velocity correlation focusing
Correlated phase space distributions of ions in an orthogonal time-of-flight mass spectrometer, 20
- Spark discharge
TOF-SIMS characterisation of spark-generated nanoparticles made from pairs of Ir–Ir and Ir–C electrodes, 70
- Spectroscopy
Gas-phase infrared multiple photon dissociation spectroscopy of mass-selected molecular ions, 1
- Sterol
Identification of molecular species of simple lipids by normal phase liquid chromatography–positive electrospray tandem mass spectrometry, and application of developed methods in comprehensive analysis of low erucic acid rapeseed oil lipids, 106

Tandem mass spectrometry

The loss of $\text{NH}_2\text{O}^\bullet$ from the *N*-hydroxyacetamide radical cation $\text{CH}_3\text{C}(=\text{O})\text{NHOH}^{\bullet+}$: An ion-catalysed rearrangement, 127

Thermal desorption

Reactions in nanofilms of trifluoroacetic acid (CF_3COOH) driven by low energy electrons, 63