

Erratum

Erratum to “Interaction of liposome formulations with human skin in vitro” [Int. J. Pharm. 229 (2001) 117–129][☆]

Gabriele Betz ^a, Roger Imboden ^b, Georgios Imanidis ^{a,*}

^a Pharmaceutical Technology, Department of Pharmacy, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Im Link 9b, 4142 Munchenstein, Switzerland

The publisher regrets that in the above article, Fig. 3 parts (a) and (b) were incorrectly orientated. Please find the correct figure produced overleaf.

[☆] PII of original article: S0378-5173(01)00824-9.

* Corresponding author. Tel.: +41-61-267-1513; fax: +41-61-267-1516.

E-mail address: georgios.imanidis@unibas.ch (G. Imanidis).




Fig. 3. Optical cross sections perpendicular to the skin surface. (a): PL80/SM formulation with Heparin. (b): PL80 formulation with Heparin. (c): PL80 formulation without Heparin. Scale: (a): $1000 \times 1000 \mu\text{m}$; (b) and (c): $2000 \times 2000 \mu\text{m}$. The stratum corneum surface of the skin faces to the left hand side of the image. Three-dimensional stereo anaglyph generated by superimposed images shifted by -5° and $+5^\circ$ which are represented by red and green color and produce the yellow color when combined.